

**ES/ER/TM-85/R3**

**Toxicological Benchmarks for Screening  
Contaminants of Potential Concern  
for Effects on Terrestrial Plants:  
1997 Revision**

This document has been approved  
by the East Tennessee Technology Park  
Technical Information Office  
for release to the public. Date: \_\_\_\_\_

**Toxicological Benchmarks for Screening  
Contaminants of Potential Concern  
for Effects on Terrestrial Plants:  
1997 Revision**

R. A. Efroymson  
M. E. Will  
G. W. Suter II  
A. C. Wooten

Date Issued—November 1997

Prepared for the  
U.S. Department of Energy  
Office of Environmental Management  
under budget and reporting code EW 20

LOCKHEED MARTIN ENERGY SYSTEMS, INC.  
managing the  
Environmental Management Activities at the  
East Tennessee Technology Park  
Oak Ridge Y-12 Plant   Oak Ridge National Laboratory  
Paducah Gaseous Diffusion Plant   Portsmouth Gaseous Diffusion Plant  
under contract DE-AC05-84OR21400  
for the  
U.S. DEPARTMENT OF ENERGY

## **PREFACE**

The purpose of this technical memorandum is to present plant toxicity data and discuss their utility as benchmarks for determining the hazard to terrestrial plants caused by contaminants in soil. This work was performed under Work Breakdown Structure 1.4.12.2.3.04.07.02 (Activity Data Sheet 8304). This report presents a standard method for deriving benchmarks, a set of data concerning effects of chemicals in soil or soil solution on plants, and a set of phytotoxicity benchmarks for 38 chemicals potentially associated with United States Department of Energy sites. In addition, background information on the phytotoxicity and occurrence of the chemicals in soils is presented, and literature describing the experiments from which data were drawn for benchmark derivation is reviewed.

## **ACKNOWLEDGMENTS**

The authors would like to thank the following individuals for their helpful reviews of the document: Richard Bonczek, Jeffrey Duncan, Ruth Hull, David Kocher, Bobette Nourse, Dennis Opresko, Mark Stack, and Barbara Walton. Parts of the literature review for this report were conducted by Melanie Futrell and Christopher Evans.

# CONTENTS

PREFACE .....	iii
ACKNOWLEDGMENTS .....	v
TABLES .....	xi
ABBREVIATIONS .....	xiii
EXECUTIVE SUMMARY .....	xv
1. INTRODUCTION .....	1-1
1.1 SCREENING BENCHMARKS IN ECOLOGICAL RISK ASSESSMENT .....	1-1
1.2 CHEMICALS IN THE SOIL-PLANT SYSTEM .....	1-2
2. METHODS .....	2-1
2.1 DATA .....	2-1
2.2 SELECTION OF TYPES AND LEVELS OF EFFECTS .....	2-2
2.3 DERIVATION OF BENCHMARKS .....	2-3
3. TOXICITY DATA REVIEW .....	3-1
3.1 INORGANIC CHEMICALS .....	3-1
3.1.1 Aluminum .....	3-1
3.1.2 Antimony .....	3-3
3.1.3 Arsenic .....	3-4
3.1.4 Barium .....	3-5
3.1.5 Beryllium .....	3-5
3.1.6 Bismuth .....	3-6
3.1.7 Boron .....	3-6
3.1.8 Bromine .....	3-7
3.1.9 Cadmium .....	3-7
3.1.10 Chromium .....	3-13
3.1.11 Cobalt .....	3-14
3.1.12 Copper .....	3-15
3.1.13 Fluorine .....	3-16
3.1.14 Iodine .....	3-16
3.1.15 Iron .....	3-17
3.1.16 Lead .....	3-17
3.1.17 Lithium .....	3-20
3.1.18 Manganese .....	3-20
3.1.19 Mercury .....	3-21
3.1.20 Molybdenum .....	3-23
3.1.21 Nickel .....	3-24
3.1.22 Selenium .....	3-26
3.1.23 Silver .....	3-27
3.1.24 Technetium .....	3-28

3.1.25	Tellurium	3-28
3.1.26	Thallium	3-29
3.1.27	Tin	3-29
3.1.28	Titanium	3-30
3.1.29	Uranium	3-30
3.1.30	Vanadium	3-31
3.1.31	Zinc	3-31
3.2	ORGANIC COMPOUNDS	3-33
3.2.1	Acenaphthene	3-33
3.2.2	Aniline	3-34
3.2.3	Biphenyl	3-34
3.2.4	4-Bromoaniline	3-34
3.2.5	3-Chloroaniline	3-34
3.2.6	4-Chloroaniline	3-34
3.2.7	2-Chlorophenol	3-34
3.2.8	3-Chlorophenol	3-35
3.2.9	4-Chlorophenol	3-35
3.2.10	2-Cresol	3-35
3.2.11	Di-n-butyl phthalate	3-35
3.2.12	3,4-Dichloroaniline	3-35
3.2.13	2,4-Dichlorophenol	3-35
3.2.14	3,4-Dichlorophenol	3-36
3.2.15	Diethylphthalate	3-36
3.2.16	2,4-Dinitrophenol	3-36
3.2.17	Furan	3-36
3.2.18	Heptane	3-36
3.2.19	Hexachlorocyclopentadiene	3-37
3.2.20	Naphthalene	3-37
3.2.21	3-Nitroaniline	3-37
3.2.22	4-Nitroaniline	3-37
3.2.23	Nitrobenzene	3-37
3.2.24	4-Nitrophenol	3-37
3.2.25	Pentachlorophenol	3-38
3.2.26	Phenol	3-38
3.2.27	Polychlorinated biphenyls (PCBs)	3-38
3.2.28	Styrene	3-39
3.2.29	2,3,5,6-Tetrachloroaniline	3-39
3.2.30	Tetrachloroethene	3-39
3.2.31	Toluene	3-40
3.2.32	4-Toluidine	3-40
3.2.33	2,4,5-Trichloroaniline	3-40
3.2.34	Trichloroethane	3-40
3.2.35	2,4,5-Trichlorophenol	3-40
3.2.36	2,4,6-Trichlorophenol	3-40
3.2.37	Ortho-xylene	3-41
3.2.38	Xylene	3-41

4. RELATIONSHIP BETWEEN SOIL PHYTOTOXICITY BENCHMARKS AND OTHER ECOTOXICOLOGICAL CRITERIA .....	4-1
4.1 COMPARISON OF PHYTOTOXICITY BENCHMARKS FOR CONTAMINANTS IN SOIL TO CANADIAN ENVIRONMENTAL QUALITY CRITERIA FOR CONTAMINATED SITES .....	4-1
4.2 COMPARISON OF PHYTOTOXICITY BENCHMARKS FOR CONTAMINANTS IN SOIL TO RIVM (NETHERLANDS) ECOTOXICOLOGICAL INTERVENTION VALUES FOR CONTAMINANTS IN SOILS .....	4-1
5. COMPARISON OF PHYTOTOXICITY BENCHMARKS FOR CONTAMINANTS IN SOIL TO CONCENTRATIONS OF CHEMICALS IN UNPOLLUTED SOILS .....	5-1
5.1 COMPARISON TO USGS ELEMENT CONCENTRATIONS IN SOILS AND OTHER SURFICIAL MATERIALS OF THE EASTERN UNITED STATES ....	5-1
5.2 COMPARISON TO DOE OAK RIDGE RESERVATION BACKGROUND SOIL CHARACTERIZATION ELEMENT CONCENTRATIONS IN SOILS .....	5-1
6. RECOMMENDATIONS AND CONCLUSIONS .....	6-1
7. REFERENCES .....	7-1
APPENDIX A. PHYTOTOXICITY DATA DERIVED FROM EXPERIMENTS CONDUCTED IN SOIL .....	A-1
APPENDIX B. PHYTOTOXICITY DATA DERIVED FROM EXPERIMENTS CONDUCTED IN SOLUTION CULTURE .....	B-1

## TABLES

1	Screening benchmark concentrations for the phytotoxicity of chemicals in soil and soil solution . . . . .	2-5
2	Comparison of screening benchmark concentrations for the phytotoxicity of chemicals in soil to CCME remediation criteria (RC), RIVM ecotoxicological intervention values (EIVs), arithmetic means of elements in uncontaminated soils of the Oak Ridge Reservation (ORR), and geometric means of elements in soils and surficial material of the eastern U.S. . . . .	4-2



## ABBREVIATIONS

CCME	Canadian Council of Ministers of the Environment
CEC	Cation Exchange Capacity
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
DNA	Deoxyribonucleic Acid
DOE	United States Department of Energy
EIV	Ecotoxicological Intervention Value
EPA	United States Environmental Protection Agency
ER-L	Effects Range Low
LCT	Lowest Concentration Tested
LOEC	Lowest Observed Effect Concentration
NOEC	No Observed Effect Concentration
ORR	Oak Ridge Reservation
PAH	Polyaromatic Hydrocarbon
PCB	Polychlorinated Biphenyl
RC	Remediation Criteria
RIVM	National Institute of Public Health and Environmental Protection
USGS	United States Geologic Survey

## **EXECUTIVE SUMMARY**

One of the initial stages in ecological risk assessment for hazardous waste sites is screening contaminants to determine which of them are worthy of further consideration as contaminants of potential concern. This process is termed contaminant screening. It is performed by comparing measured ambient concentrations of chemicals to benchmark concentrations. Currently, no standard benchmark concentrations exist for assessing contaminants in soil with respect to their toxicity to plants.

This report presents a standard method for deriving benchmarks for this purpose (phytotoxicity benchmarks), a set of data concerning effects of chemicals in soil or soil solution on plants, and a set of phytotoxicity benchmarks for 38 chemicals potentially associated with United States Department of Energy (DOE) sites. In addition, background information on the phytotoxicity and occurrence of the chemicals in soils is presented, and literature describing the experiments from which data were drawn for benchmark derivation is reviewed. Chemicals that are found in soil at concentrations exceeding both the phytotoxicity benchmark and the background concentration for the soil type should be considered contaminants of potential concern.

# 1. INTRODUCTION

## 1.1 SCREENING BENCHMARKS IN ECOLOGICAL RISK ASSESSMENT

An important step in ecological risk assessment is screening the chemicals occurring on a site for contaminants of potential concern. Screening may be accomplished by comparing reported concentrations in media to a set of toxicological benchmarks. Multiple endpoints have been established for assessment of risks posed by soil-borne contaminants to organisms directly impacted by them. This report supersedes a prior report on screening benchmarks for phytotoxicity (Will and Suter 1995a). Benchmarks for toxic effects of contaminants on earthworms and soil microbial processes are presented in a companion report (Will and Suter 1995b), which will also be revised in 1997.

If a chemical concentration or the reported detection limit exceeds the screening benchmark, more analysis is needed to determine the hazards posed by that chemical (i.e., it is a contaminant of potential concern). If, however, the chemical concentration or its detection limit falls below the proposed benchmark, the chemical may be ignored during further study unless public concern or ancillary evidence suggest that it should be retained.

The purpose of this report is to present plant toxicity data and to use them to derive benchmarks for determining the hazard to terrestrial plants caused by contaminants in soil. Benchmarks are provided for soils and solutions.

Tests of the toxicity of chemicals in the rooting medium of plants are conducted using a variety of rooting media which have been divided into two categories for purposes of this report: soil and solution. In a previous version of this document, data from experiments conducted in other growth media were provided, such as vermiculite and quartz sand. However, these data were determined to be not applicable to field situations and were not used in benchmark derivation; therefore, these data have been omitted from the present revision of the document.

Tests conducted in natural soils (even when brought into the laboratory, dried, sieved, fertilized, etc.) are assumed to be representative of the exposure of plants to contaminants measured in field soils. Tests conducted in nutrient and mineral solutions are assumed to be representative of exposures of plants to contaminants measured in soil solutions (e.g., from lysimeter samples or possibly from aqueous extracts of soil) or in very shallow groundwater (e.g., plants in the vicinity of seeps and springs).

Soil benchmarks are based on data provided by toxicity studies in the field or more commonly in greenhouse and growth chamber settings. Most of the soil concentrations of metals reported from waste sites are from extractions with hydrochloric acid or other mineral acids which are intended to provide total concentrations. Similarly, concentrations of organic contaminants in waste site soils are total concentrations derived from rigorous solvent extractions. In some cases, toxicity tests report concentrations extracted from contaminated soils, but various extractants are used that may not yield total concentrations. More commonly, the concentrations reported are nominal concentrations of a soluble form (i.e., a highly bioavailable form) of the chemical added to soil. Most metals in natural soils and contaminants of waste sites are in poorly available forms.

Solution benchmarks include data from toxicity tests conducted using whole plants rooted in aqueous solutions. Tests are commonly conducted in this manner because plants are assumed to be

exposed to contaminants in the solution phase of soil, and the presence of soil in test systems reduces the experimenter's degree of control over exposure. Groundwater samples from waste sites are typically acidified before analysis to obtain total concentrations, but some samples are filtered before acidification. In general, the concentrations in filtered samples are likely to be more comparable to the concentrations reported from solution toxicity tests and should be used if available.

These benchmarks are to serve primarily for contaminant screening. An assessor must realize that the soil and plant characteristics discussed in the following sections play a large part in plant toxicity and incorporate these site-specific considerations in the evaluation of the potential hazards of a chemical. If chemical concentrations reported in field soils that support vigorous and diverse plant communities exceed one or more of the benchmarks presented in this report or if a benchmark is exceeded by background soil concentrations, it is generally safe to assume that the benchmark is a poor measure of risk to the plant community at that site.

## **1.2 CHEMICALS IN THE SOIL-PLANT SYSTEM**

Elements occur in the soil in a variety of forms more or less available for uptake by plants. Many of the contaminants of concern at waste sites are metals or metalloids. Availability is determined by characteristics of the elements, such as behavior of the ion as a Lewis acid (electron acceptor) which determines the predominant type and strength of bond created (ionic or covalent) and, therefore, the mobility of the metal in the soil environment. Soil characteristics (e.g., pH, clay and organic matter content and type, and moisture content) also determine availability to plants by controlling speciation of the element, temporary immobilization by particle surfaces (adsorption-desorption processes), precipitation reactions, and availability in soil solution. The most general sinks for metals are iron and manganese oxides and organic matter (Jenne and Luoma 1977). Although particulate soil organic matter serves to immobilize metals, soluble organic matter may act to keep metals in solution in a form absorbed and translocated by plants.

The final control on availability of metals and metalloids in soil to plants is the selective absorption from soil solution by the root. Metals may be bound to exterior exchange sites on the root and not actually taken up. They may enter the root passively in organic or inorganic complexes with the mass flow of water or actively by way of metabolically controlled membrane transport systems often meant to take up a nutrient which the "contaminant" metal mimics. At different soil solute concentrations, metals may be absorbed by both processes. Absorption mechanisms and quantity absorbed are influenced by plant species (and cultivar), growth stage, physiological state, and the presence of other elements.

Once in the plant, a metal can be sequestered in the roots in vacuoles or in association with cell walls and organelles or translocated to above ground parts in xylem as organic or inorganic complexes. Location and forms of metals in plants, as well as their toxic effects, depend on plant species, growth stage, physiological state, and presence of other metals.

Mechanisms of toxicity of metals tend to be dependent on the nature of the reactivity of the metal itself. They may alter or inhibit enzyme activity, interfere with deoxyribonucleic acid (DNA) synthesis or electron transport, or block uptake of essential elements. Variability in response to 'toxic' levels of metals by different plants is due to a number of defenses. These include exclusion from the root, translocation in nontoxic form, sequestering in nontoxic form in the root or other plant parts, and formation of unusable complexes containing metals that may otherwise be inserted into biomolecules instead of the proper element (e.g., As replacing P) (Peterson, 1983).

Organic compounds of environmental concern include nonionic compounds [pyrene, chlorinated benzenes, polychlorinated biphenyls (PCBs), toluene, and many pesticides], ionizable compounds (chlorophenols, carboxylic acids, surfactants, and amines), and weakly hydrophobic volatile organic compounds (trichloroethene). For the nonionic compounds, sorption in soil is mainly a function of degree of hydrophobicity and amount of sorbent hydrophobic phase (i.e., soil organic matter). Sorption of the compound by soil organic matter is reversible. The activities of these compounds in soil can be predicted by the organic matter-water coefficient,  $K_{om}$ , as estimated by the octanol-water coefficient,  $K_{ow}$ . Absorption onto colloidal organic matter in solution may alter the availability of these nonionic compounds. Ionizable compounds contain anionic or cationic moieties or both within their structure. These charged structures interact with organic and inorganic charged surfaces in the soil in a variety of reversible reactions. The extent and nature of the associations with charged surfaces depends on characteristics of the organic compound, solution pH and ionic strength, and mineral composition of the soil particulates (Schwarzenbach et al. 1993). Organic compounds may be degraded by microorganisms in the soil to metabolites with greater or lesser toxicity. Very stable compounds, like highly chlorinated PCBs, may persist in essentially unaltered form for many years.

Plant roots are not discriminating in uptake of small organic molecules (molecular weight less than 500) except on the basis of polarity. More water soluble molecules pass through the root epidermis and translocate throughout the plant. The less soluble compounds (like many polycyclic aromatic hydrocarbons) seem to have limited entry into the plant and minimal translocation once inside. Highly lipophilic compounds, such as PCBs, move into the plant root via the symplastic route (from cell to cell, as opposed to between the cells) and are translocated within the plant.

## 2. METHODS

### 2.1 DATA

References on the toxicity of selected chemicals to terrestrial plants were obtained from searches of bibliographic databases (BIOSIS, POL TOX I, current contents), a numeric database (PHYTOTOX), review articles, and conventional literature searches. Reports of toxicity tests of individual chemicals in laboratory, greenhouse, or field settings were obtained.

Data presented in this report were derived mainly from primary sources. Secondary sources were used if the primary source cited in the secondary source was unavailable, if only a small amount of data for a particular chemical was available, and if secondary sources suggested that a benchmark derived from limited primary source material was too high. The general criteria for inclusion of a study in the data set used to derive phytotoxicity benchmarks were:

1. if the methodology was clearly stated (especially concentrations of applied chemicals) and followed in the experiment,
2. if results were quantified as measures of plant growth or yield (e.g., weight, height) (measures of metabolic activity or tissue chemical concentration were used if measures of growth or yield were not available for a particular chemical of interest),
3. if results were presented in numeric form or graphical presentations of data were clearly interpretable, and
4. if an unambiguous reduction existed in the measured parameter within the range of applied concentrations of the chemical of interest.

The data selected for soil benchmarks are given in Appendix A. They were selected using these criteria and were assigned to the following categories for analysis:

1. Chemical—The effects of individual chemicals of interest were analyzed. In the case of metals, the metal is listed in the "Chemical" field. For organics, the compound is listed in the "Chemical" field.
2. Chemical Form—The form in which the chemical was added to the experimental medium.
3. Soil Type—Soil textural classification, if provided.
4. Cation exchange capacity (CEC) is the sum of the exchangeable cations that a soil can adsorb, expressed as milliequivalents per 100 g soil. Soil organic and inorganic constituents contain negatively charged sites that are the location of important interactions with positively charged ions in soil solution. These interactions partially control the effective toxicity of many contaminants.
5. Organic matter—Soil organic matter is important in reactions of many contaminants in the soil. Percentage organic carbon, if given, was converted to the more frequently cited measure of percentage organic matter by the equation (Nelson and Sommers 1982):

$$\% \text{organic carbon} \times 2 = \% \text{organic matter}$$

6. Soil pH—The pH of the soil system is a critical controller of the reactions occurring in the soil and therefore of the toxicity of any given quantity of chemical in the soil-plant system.
7. Plant Species—The analysis was limited to terrestrial vascular plants. Common names are given.
8. Exposure duration—The durations of exposure of the test plants to chemicals of interest ranged from 2 to 335 days, with trees generally being exposed longer than plants with shorter life spans.
9. NOEC Applied—The no observed effect concentration (NOEC) is defined herein as the highest applied concentration of the chemical of interest which gave a reduction of 20% or less in a measured response.
10. LOEC Applied—The lowest observed effect concentration (LOEC) is defined herein as the lowest applied concentration of the chemical of interest which gave a greater than 20% reduction in a measured response. In some cases, the LOEC for the test was the lowest concentration tested (LCT) or the only concentration reported, as when the EC<sub>50</sub> was reported.
11. Growth parameter—The majority of the growth responses were oven-dry weights of whole plants or their parts. Others included root length, plant height, relative growth rate, grain yield, seeds per plant, percent seed germination, and fresh and air-dry weights. Responses other than these growth and yield parameters were included only if growth or yield parameters were unavailable for a chemical.

The data selected for solution benchmarks using these criteria were assigned to the same categories for analysis with several exceptions. Categories relating to soil characteristics (type, CEC, % organic matter) were not applicable. These data are presented in Appendix B.

## **2.2 SELECTION OF TYPES AND LEVELS OF EFFECTS**

Growth and yield parameters were used for two reasons. First, they are the most common class of response parameters reported from phytotoxicity studies; thus, they permit derivation of reasonably consistent benchmarks for a large number of contaminants. Second, growth and yield are ecologically significant responses both in terms of the plant populations and the ability of the vegetation to support higher trophic levels.

Twenty percent reduction in growth or yield was used as the threshold for significant effects to be consistent with other screening benchmarks for ecological risk assessment and with current regulatory practice (Suter et al. 1995). In brief, most regulatory criteria are based on concentrations in toxicity tests that cause effects which are statistically significantly different from controls. On average, those concentrations correspond to greater than a 20% difference in effects. In addition, regulatory actions may be based on comparisons of biological parameters measured on contaminated sites to those from reference sites. Differences between parameters at sites generally must be greater than 20% to be reliably detected in such studies. Therefore, the 20% effects level is treated as a conservative approximation of the threshold for regulatory concern.

## **2.3 DERIVATION OF BENCHMARKS**

Because of the diversity of soils, plant species, chemical forms, and test procedures, it is not possible to estimate concentrations that would constitute thresholds for toxic effects on the plant

communities at particular sites from published toxicity data. This situation is analogous to the problem of deriving benchmarks for sediments. In this report, the method used for deriving soil benchmarks is based on the National Oceanographic and Atmospheric Administration's method for deriving the Effects Range Low (ER-L) (Long and Morgan 1990), which has been recommended as a sediment screening benchmark by the United States Environmental Protection Agency (EPA) Region IV. The ER-L is the 10th percentile of the distribution of various toxic effects thresholds for various organisms in sediments.

This approach can be justified by assuming that the phytotoxicity of a chemical in soil is a random variate, the toxicity of contaminated soil at a particular site is drawn from the same distribution, and the assessor should be 90% certain of protecting plants growing in the site soil. Any bias in the data set would mitigate against that assumption. In this implementation of the approach, the bias most likely to be significant is the use of soluble metal salts in the toxicity tests. These salts are likely to be more toxic than the mixture of forms encountered in field soils. That bias would result in conservative benchmark values. Other possible sources of bias include the exclusion of synergistic and antagonistic effects resulting from interactions between chemicals, the use of predominately domestic plant species that may not be representative of plant species in general, the use of predominately agricultural soils which may not be representative of soils in general, and the laboratory test conditions which may not be representative of field conditions. The direction and magnitude of these potential biases is unknown.

The phytotoxicity benchmarks were derived by rank-ordering the LOEC values and then picking a number that approximated the 10th percentile. As with the ER-Ls, statistical fitting was not used because there were seldom sufficient data and because these benchmarks are to be used as screening values and do not require the consistency and precision of regulatory criteria. If there were 10 or fewer values for a chemical, the lowest LOEC was used. If there were more than 10 values, the 10th percentile LOEC value was used. If the 10th percentile fell between LOEC values, a value was chosen by interpolation. Since these benchmarks are intended to be thresholds for significant effects on growth and production, test endpoints that indicate a high frequency of lethality are not appropriate. Therefore, when a benchmark is based on an  $LC_{50}$  or on some other endpoint that includes a 50% or greater reduction in survivorship, the value is divided by a factor of 5. This factor is based on the authors' expert judgment. Although there is not a body of data for comparison of lethal and sublethal effects concentrations in tests conducted with the same species and soils, it is the authors' impression that a factor of 5 approximates the ratio  $LC_{50}/EC_{20}$ .

In all cases, benchmark values were rounded down to one significant figure. This rounding was done for two reasons. First, it is not appropriate to ascribe greater precision to a number than it actually possesses; these benchmarks are very imprecise. Second, the rounding serves to emphasize the fact that the benchmarks are conceptually distinct from the test endpoint values from which they were derived. That is, a LOEC may be a precise estimate of the lowest toxic concentration for a particular plant variety in a particular test system, but when an LOEC is used as a benchmark for all plants in field soils, it is a qualitatively different and much more poorly defined value.

Another source of benchmark values was published reviews of the phytotoxicity literature. When primary literature was unavailable for a particular contaminant, concentrations identified in reviews as thresholds for phytotoxicity were used as benchmarks. In addition, when fewer than three LOEC values were found for a chemical in soil or solution and a toxicity threshold from a review was lower than the lowest LOEC, the toxicity threshold was used as the benchmark for that chemical. Proposed screening benchmarks for phytotoxic effects of contaminants in soils and solutions are presented in Table 1.

This method of deriving screening benchmarks for soil organisms may strike some readers as insufficiently conservative. That impression could result from the fact that the derivation of the



benchmark (like the derivation of the ER-L values) implies a significant effect on approximately 10% of the species. However, the authors believe that the method described in this report is sufficiently conservative for the following reasons. First, these benchmarks were derived for a community-level assessment endpoint. Given the water, nutrient, or physical limitations of most soil and litter-dwelling communities, a reduction in growth, reproduction, or functioning of 10% of component species is likely to be acceptable. Second, the benchmarks derived by these methods have proved to be conservative in practice. In some locations for some elements, they are lower than background concentrations (Section 4). This is believed to be caused by the fact that they are based on toxicity tests which dose growth substrates with soluble salts of metals. Therefore, they are much more available than most naturally occurring metals, and even metals at many, if not most, waste sites.

In this report, the authors have attempted to assign levels of confidence to the benchmarks. The criteria that best reflect that confidence are as follows:

1. Low Confidence—Benchmarks based on fewer than 10 literature values.
2. Moderate Confidence—Benchmarks based on 10 to 20 literature values.
3. High Confidence—Benchmarks based on over 20 literature values.

Confidence in a benchmark based on more than 20 reported toxic concentrations may be reduced to moderate if the range of plant species tested is narrow, i.e., no tree species or only one family of plants were tested. Moderate or high confidence benchmarks may be demoted one level if the value approximating the 10th percentile was the lowest concentration tested and caused a greater than 30% reduction in the measured growth parameter. Although these criteria may seem arbitrary, the result is a confidence classification that fairly reflects the authors' professional judgment.

Any scheme for deriving a set of standard ecotoxicological benchmarks is based on assumptions that may be questioned by readers. The procedure used herein is one that is consistent with current regulatory practice and contains a minimum of assumptions or factors. Those who care to make other assumptions or to add safety factors may make use of the data presented herein to calculate their own benchmarks.

**Table 1. Screening benchmark concentrations for the phytotoxicity of chemicals in soil and soil solution**

Chemical	Soil (mg/kg)	Solution (mg/L)	Chemical	Soil (mg/kg)	Solution (mg/L)
Aluminum	50	0.3	4-Bromoaniline	—	100
Antimony	5	—	3-Chloroaniline	20	—
Arsenic	10	0.001	4-Chloroaniline	—	40
Barium	500	—	2-Chlorophenol	—	60
Beryllium	10	0.5	3-Chlorophenol	7	—
Bismuth	—	20	4-Chlorophenol	—	50
Boron	0.5	1	2-Cresol	—	50
Bromine	10	10	3,4-Dichloroaniline	—	10
Cadmium	4	0.1	2,4-Dichlorophenol	—	20
Chromium	1	0.05	3,4-Dichlorophenol	20	—
Cobalt	20	0.06	2,4-Dinitrophenol	20	—
Copper	100	0.06	Di-n-butyl phthalate	200	—
Fluorine	200	5	Diethylphthalate	100	20
Iodine	4	0.5	Furan	600	100
Iron	—	10	Heptane	—	1
Lead	50	0.02	Hexachlorocyclopentadiene	10	0.1
Lithium	2	3	Naphthalene	—	10
Manganese	500	4	3-Nitroaniline	—	70
Methyl mercury	—	0.0002	4-Nitroaniline	—	40
Mercury	0.3	0.005	Nitrobenzene	—	8
Molybdenum	2	0.5	4-Nitrophenol	—	10
Nickel	30	0.5	Pentachlorophenol	3	0.03
Selenium	1	0.7	Phenol	70	10
Silver	2	0.1	PCBs	40	—
Technetium	0.2	0.2	Styrene	300	10
Tellurium	—	2	2,3,5,6-Tetrachloroaniline	20	—
Thallium	1	0.05	Tetrachloroethene	—	10
Tin	50	100	Toluene	200	10
Titanium	—	0.06	4-Toluidine	—	100
Uranium	5	40	2,4,5-Trichloroaniline	20	—
Vanadium	2	0.2	Trichloroethane	—	100
Zinc	50	0.4	2,4,5-Trichlorophenol	4	—
Acenaphthene	20	0.1	2,4,6-Trichlorophenol	—	10
Aniline	—	200	Ortho-xylene	—	1
Biphenyl	60	2	Xylene	—	100

### 3. TOXICITY DATA REVIEW

Results of the literature review are summarized in Appendixes A and B. A short noncritical review of the literature from which data were derived for the calculation of benchmarks is presented in the following text. All soil experiments were conducted in pots in a greenhouse (glass or screen) unless otherwise noted. Experiments conducted in solution culture were generally conducted in growth chambers although some experimental setups were contained in greenhouses. Confidence in the benchmark for a particular chemical is also discussed in the following text. The criteria used to establish confidence levels are given in Section 2. The units of ppm are equivalent to mg/kg for chemicals in soil and mg/L for chemicals in solution.

Information is also given on the mechanisms of phytotoxicity of the chemicals. The mechanisms of growth reductions measured are seldom discussed in the literature from which toxicity data are extracted for benchmark calculations. This information is offered to allow a better understanding of the potential mechanisms of toxicity of these and related contaminants.

#### 3.1 INORGANIC CHEMICALS

##### 3.1.1 Aluminum

**Experiments conducted in soil.** Seedling establishment of white clover (*Trifolium repens* L.) in a silt loam soil (pH 5.0) was reduced approximately 30% by the addition of 50 ppm Al as  $\text{Al}_2(\text{SO}_4)_3$  (Mackay et al., 1990), the lowest concentration tested. This lone study does not allow a high degree of confidence in the benchmark.

**Experiments conducted in solution.** Goransson and Eldhuset (1991) evaluated the effect of Al on root and shoot growth of seedlings of Norway spruce (*Picea abies* L.) and Scots pine (*Pinus sylvestris* L.) in a nutrient solution of pH 3.8. The spruce proved much more sensitive to Al with a 33% reduction in root growth weight after 21 days at 8.1 ppm Al in solution (5.4 ppm had no effect). Pine shoot growth rate was reduced 40% with 270 ppm, while 162 ppm Al had no effect.

Godbold and Kettner (1991) measured a 42% reduction in mean root length of 3-wk old Norway Spruce seedlings grown in a nutrient solution (pH 4) containing 5.4 ppm Al ( $\text{AlCl}_3$ ) for 8 days. Aluminum at 1.4 ppm had no effect on plant growth.

Nichol and Oliveira (1995) investigated the effect of aluminum in a hydroponic medium on root growth of a barley cultivar (*Hordeum vulgare*). For seeds germinated in the solution containing aluminum at 0.0027 ppm, root growth was inhibited by about 25% two days following germination and by about 60% four days following germination.

Pintro et al. (1996) studied the effect of aluminum (as chloride) in solution on the growth of two cultivars of corn, one aluminum-sensitive and one aluminum-tolerant (*Zea mays* L., HS7777 and C525-M). At 0.27 ppm, aluminum reduced root elongation and root weight of the aluminum-sensitive plants by about 50% and 20%, respectively. The NOEC was 0.13 ppm. A concentration of 0.405 ppm reduced the root elongation of the aluminum-tolerant cultivar by about 30%, with a NOEC of 0.13 ppm. The activities and ionic strengths of aluminum were calculated in the paper.

Zavas et al. (1996) exposed plants from two populations of perennial grass (*Piptatherum miliaceum*) to aluminum (as chloride) in solution for 16 days. The two populations were from a bauxite area and a pasture soil in Greece. Tests were conducted at pH 4.5 and 10. At the lower pH and at an

aluminum concentration of 12.9 ppm, the mean root length of plants from the bauxite area was reduced by 73% and that of plants from the pasture soil was reduced by 76%. The NOEC was 2.2 ppm. No toxicity was observed at a pH of 10.

Keltjens (1990) tested the responses of roots and shoots of 1-yr-old Douglas fir (*Pseudotsuga menziesii* L.) seedlings to Al (as chloride) in a pH 3.5 solution at 4, 6, 8, 16, and 32 ppm. Calcium and magnesium were added at 0.1, 0.5, or 2.5 mM. After a 9-month exposure to aluminum at 32 ppm with Ca and Mg present at 0.5 mM, root length was reduced 43%, and root weight was reduced about 30%. Shoot weight was reduced about 40% by exposure to 8 ppm Al with Ca and Mg added at 2.5 mM.

Lin and Myhre (1991) compared the tolerance of citrus rootstock seedlings to growth in solution (pH 4) containing Al (as  $\text{Al}_2(\text{SO}_4)_3$ ) by measuring root length, shoot height, and plant weight. After 60 days, three of the five rootstocks had reduced weight at 8.3 ppm Al. Percent reduction ranged from 22 to 45% at that concentration. The citrange rootstock root length was decreased 21% at 2.7 ppm Al. The Cleopatra mandarin rootstock had a 30% reduction in weight at 24.4 ppm.

Wheeler and Follet (1991) evaluated the effect of Al as  $\text{Al}_2(\text{SO}_4)_3$  in solution culture (pH 4.7) on root and shoot weights of onions (*Allium cepa* L.), asparagus (*Asparagus officinalis* L.), and squash (*Cucurbita maxima* L.). Root and shoot weights of onions were reduced 68 and 23% after 31 days of growth in solution containing 0.05 ppm (lowest concentration tested). Root and shoot weight of asparagus were reduced 49 and 70% in solution containing 0.13 ppm Al, while 0.05 ppm had no effect. Root weight of squash was reduced 25% after 26 days of growth in solution containing 0.27 ppm while 0.13 ppm had no effect.

McLean and Gilbert (1927) used nutrient solution culture to test the comparative resistance of different plants to Al toxicity. Carrot (*Daucus carota* L.) seedling weight was reduced approximately 75% after 126 days of growth in solution containing 3.6 ppm Al (lowest concentration tested) in two experiments. Radish (*Raphanus sativus* L.) seedling root and shoot weights were reduced 21% after 77 days of growth in solution containing 3.6 ppm Al, while 1.8 ppm had no effect. Turnip (*Brassica rapa* L.) seedling top weight was reduced 39% after 77 days of growth in solution containing 7.2 ppm Al, while 3.6 ppm had no effect. In two experiments with slightly different nutrient solutions, beet seedling weight was reduced approximately 25% after 126 days of growth in solution containing 1.8 ppm Al (lowest concentration tested). Seedling weight was diminished 74% by 1.8 ppm Al (lowest concentration tested) in a third experiment. In two 56-day experiments in slightly different nutrient solutions, lettuce weight was reduced 39% by 1.8 ppm (0.9 ppm had no effect) and 55% by 2.7 ppm Al (1.8 ppm had no effect) in solution.

In a third experiment lasting 42 days, lettuce top weight was reduced 25% by 1.1 ppm, while 0.5 ppm Al had no effect. Cabbage and oat seedling weights were reduced 43% and approximately 25% by 7.2 ppm Al (lowest concentration tested) after 98 and 63 days, respectively. After 77 days, barley seedling root and shoot weights were reduced 47 and 22% by 1.8 ppm Al (lowest concentration tested). After 63 days, rye seedling root weight was reduced 22% by 1.8 ppm Al (lowest concentration tested). In a second experiment, plants grown in an alternate nutrient solution suffered a root weight of 25% in the presence of 3.6 ppm Al, while 1.8 ppm had no effect.

Wallace and Romney (1977a) grew rice (*Oryza sativa* L.) and soybean (*Glycine max* L.) seedlings in solution culture containing Al as  $\text{Al}_2(\text{SO}_4)_3$  for 13 days. Root and shoot weights of rice were reduced 28 and 27% by 2.7 ppm Al, while 0.27 ppm had no effect. Leaf weight of soybeans was reduced 33% by the same concentration.

MacLeod and Jackson (1967) tested two varieties of barley (*Hordeum vulgare* L.) for tolerance to 4, 6, 8, 10, or 12 ppm Al (as  $\text{AlCl}_3$ ) in a pH 4.3 nutrient solution. After 30 days of growth, root and shoot weights of one variety were reduced approximately 50% by 10 ppm Al, while those of the other variety were reduced approximately 30% by 6 ppm Al.

Wong and Bradshaw (1982) evaluated the effect of Al on root and shoot length of ryegrass (*Lolium perenne* L.) grown in solution (pH 7) with Al added as  $\text{KAl}(\text{SO}_4)_2$ . After 14 days, they found a 29% reduction in the length of the longest root in response to 0.63 ppm (lowest concentration tested).

Sasaki et al. (1994) evaluated the effect of Al as  $\text{AlCl}_3$  in solution culture (pH 4.5) on root elongation of wheat cultivars (*Triticum sativum* L.). Root elongation was reduced 50% after 5 days growth in solution containing 0.27 ppm, while 0.14 ppm had no effect.

Lllugany et al. (1995) evaluated the effect of Al as  $\text{AlCl}_3$  in nutrient solution culture (pH 4.3) on root elongation of 4-d old seedlings of maize cultivars (*Zea mays* L.). For three of the four cultivars, root elongation was reduced 23 to 37% after 1 day of growth in the solution containing 0.54 ppm, the lowest concentration tested. Root elongation in the fourth cultivar during the same time period was reduced 36% in the presence of 1.35 ppm Al, while 0.54 ppm had no effect.

The authors have high confidence in the benchmark of 0.3 ppm Al. The low LOEC values are based on experiments with seedlings of field and horticultural crops. Trees, especially pines, appear to have the greatest tolerance to Al.

**Mechanism of phytotoxicity.** Aluminum interferes with cell division in roots; decreases root respiration; fixes P in unavailable forms in roots; interferes with uptake, transport, and use of Ca, Mg, P, K, and water; and interferes with enzyme activities (Foy et al. 1978). Symptoms of toxicity include stubby, brittle roots; stunting; late maturity; and collapse of growing points. Seedlings are more susceptible to damage from Al toxicity than are older plants.

### 3.1.2 Antimony

**Experiments conducted in soil.** No primary reference data exist that describe toxicity of Sb to plants grown in soil. The benchmark is based on a report of unspecified toxic effects on plants grown in a surface soil with the addition of 5 ppm Sb (Kabata-Pendias and Pendias 1984)). The authors have low confidence in the benchmark based on this study alone.

**Experiments conducted in solution.** No reference data exist that show toxicity of Sb to plants grown in solution.

**Mechanism of phytotoxicity.** Antimony is considered a nonessential metal and is easily taken up by plants if available in the soil in soluble forms (Kabata-Pendias and Pendias 1984). The only information found on phytotoxicity was a secondary reference noting undefined, qualitative phytotoxic effects on plants grown in a surface soil (Kabata-Pendias and Pendias 1984).

### 3.1.3 Arsenic

**Experiments conducted in soil.** The tolerance of spruce seedlings to As in soil was tested in field plots by Rosehart and Lee (1973). Three-year-old seedlings grown 335 days in soil to which 1000 ppm As was added as As(III) (lowest concentration tested) experienced a 50% reduction in height.

Deuel and Swoboda (1972) assessed the toxicity of As(III) added to two soils on the shoot weight of cotton (*Gossypium hirsutum* L.) and soybeans grown from seed for 6 weeks. In the fine sandy loam soil, shoot weight of both crops were reduced (cotton 22%; soybeans 45%) in the presence of 11 ppm As, the lowest concentration tested. Soybean growth in a black clay soil was reduced 28% by the addition of 22.4 ppm As, the lowest concentration tested. Cotton growth in this soil was reduced 29% by the addition of 89.6 ppm As.

Woolson et al. (1971) tested the toxicity of three sources of As(V) on corn grown from seed for 4 weeks in a loamy sand (pH 7.1). Corn fresh weight reductions rose from less than 10% with the addition of 10 ppm As in any form, to almost 100% for  $\text{NaH}_2\text{AsO}_4$ , over 75% for  $\text{Al}(\text{H}_2\text{AsO}_4)_3$ , and about 65% for  $\text{Ca}(\text{H}_2\text{AsO}_4)_2$  with the addition of 100 ppm As.

Jiang and Singh (1994) assessed the toxicity of As (III) and As (V) added to the two soils on the yield of barley and ryegrass grown from seed for 1 year in a greenhouse. The soils tested were a loam (pH 4.9, 3% organic carbon, and 19% clay) and a sand (pH 5.6, 0.4% organic carbon, and 3% clay). Sodium arsenite was more toxic to barley plants than sodium arsenate in both soils, with the greatest toxicity occurring in the sand (24% decrease at 2 ppm, the lowest concentration tested). Arsenic (V) at 250 ppm was associated with greater reduction in yield of ryegrass (63%) than the same concentration of As (III) in the loam soil (22%). In the sand, sodium arsenite reduced yield of ryegrass 34% at 50 ppm. A concentration of 250 ppm As (V) caused a 91% decrease in yield while 50 ppm had no effect. Confidence in a soil benchmark value of 10 ppm is moderate.

**Experiments conducted in solution.** Mhatre and Chaphekar (1982) tested several species at germination stage for their response to As. Seeds of sorghum, alfalfa (*Medicago sativa* L.), mung bean (*Phaseolus aureus* L.), cluster bean (*Cyamopsis tetragonoloba* L.), and radish were allowed to germinate in solutions containing 0.001, 0.01, 0.1 or 1 ppm As as  $\text{As}_2\text{O}_3$  (As III). Germination counts after 24 hours showed no effect of As. After 5 days, root length of cluster bean was reduced 29% by 0.001 ppm As. Root length of radish was reduced 21% by the addition of 0.01 ppm. Root and shoot lengths of alfalfa and mung bean were reduced (55 and 40%, 87 and 57%) by the addition of 1 ppm As.

The concentrations of As (V), from  $\text{Na}_2\text{HAsO}_4$ , required for a 50% reduction in seed germination and root length of mustard (*Sinapis alba*) after 3 days of exposure in solution (pH 7.3), was reported by Fargasova (1994).  $\text{LC}_{50}$  for germination was 30 ppm and  $\text{EC}_{50}$  for root length was 5.5 ppm As.

Bowen (1979) reported unspecified reductions in plant growth in a solution containing 0.02 ppm As.

Confidence in the solution benchmark of 0.001 ppm is low because there are less than 10 values and a limited variety of plant species tested.

**Mechanism of phytotoxicity.** Arsenic is not essential for plant growth. It is taken up actively by roots, with arsenate being more easily absorbed than arsenite. Arsenic and phosphate ions are likely taken up by the same carrier (Asher and Reay 1979). The phytotoxicity is strongly affected by the form in which it occurs in soils. Arsenite is more toxic than arsenate, and both are considerably more toxic than organic forms (Peterson et al. 1981). In experiments with toxic levels of As, rice and legumes appear to be more sensitive than other plants. Symptoms include wilting of new-cycle leaves, followed by retardation of root and top growth, and leaf necrosis (Aller et al. 1990). Because As is chemically similar to P, it is translocated in the plant in a similar manner and is able to replace P in many cell reactions. Arsenic (III) probably reacts with sulphhydryl enzymes leading to membrane degradation and

cell death. Arsenic (V) is known to uncouple phosphorylation and affect enzyme systems (Peterson et al. 1981). The mechanism of toxicity of organo-arsenicals is unclear.

### 3.1.4 Barium

**Experiments conducted in soil.** Chaudhry et al. (1977) investigated the effects of Ba added as  $\text{Ba}(\text{NO}_3)_2$  on shoot weight of barley and bush beans (*Phaseolus vulgaris* L.) grown from seed for 14 days in a loam soil. Shoot growth of barley was reduced 38% after 14 days by the addition of 500 ppm Ba, the lowest concentration tested. Shoot growth of bush beans was reduced 30% after 14 days by the addition of 2000 ppm Ba, but was not reduced at the next lowest level, 1000 ppm.

Confidence in a benchmark value of 500 ppm is low due to lack of supporting data.

**Experiments conducted in solution.** There were no reference data describing toxicity of Ba to plants grown in solution.

**Mechanism of phytotoxicity.** Barium is commonly present in plants but is not an essential component of plant tissues. It is taken up easily from acid soils (Kabata-Pendias and Pendias 1984). Mechanisms of toxicity may include competition with Ca for root uptake (Wallace and Romney 1971).

### 3.1.5 Beryllium

**Experiments conducted in soil.** Few tests of the toxicity of Be to plants grown in soil have been conducted. Confidence in the benchmark is low because the lowest of the three values is from a report of unspecified toxic effects on plants grown in a surface soil with the addition of 10 ppm Be (Kabata-Pendias and Pendias 1984).

**Experiments conducted in solution.** Romney et al. (1962) reported a 33% reduction in the weight of bush beans when grown for 48 days in a pH 5.3 nutrient solution containing 0.5 ppm Be (lowest concentration tested).

Sajwan et al. (1997) investigated the effect of beryllium on the mass of above-ground soybean plants in a sand and loamy sand soil. The addition of 25 ppm of beryllium to soil, the lowest concentration tested, led to a 49% reduction in soybean biomass in the sand and a 46% reduction in biomass in the loamy sand after 14 days. Additional tests undertaken with limed soil also showed toxicity at the 25 ppm concentration.

Romney and Childress (1965) investigated the effect of 2, 4, 8, and 16 ppm Be (as  $\text{BeCl}_2$ ; pH 5.3) on growth of barley, alfalfa, pea (*Pisum sativum* L.), and lettuce (*Lactuca sativa* L.). Barley (20 days), pea (24 days), and lettuce (28 days) weights were reduced 50, 21, and 37%, respectively, by 2 ppm Be. After 54 days, alfalfa weight was reduced 25% by 4 ppm Be.

The effects of Be, from  $\text{BeSO}_4$ , on germination and radicle length after 3 days of growth in solution of radish, cabbage (*Brassica oleracea* L.), turnip, lettuce, wheat, and millet (*Panicum miliaceum*) were determined by Carlson et al. (1991). There was no effect on seed germination up to 40 ppm Be. Treatment levels were 0, 0.5, 1, 2.5, 5, 7.5, 10, 20, 30, and 40 ppm Be. A concentration of 0.5 ppm reduced radicle length of lettuce and turnip by 62 and 63%. A concentration of 2.5 ppm reduced radicle length of cabbage by 35%. Five ppm Be reduced radicle length of radish by 32%, 20 ppm caused a 30% decrease in wheat, and 40 ppm reduced radicle length of millet by 35%.

Confidence in the benchmark for Be in solution (0.5 ppm) is low. There were 11 values to consider but a greater than 30% reduction occurred in the measure approximating the 10th percentile.

**Mechanism of phytotoxicity.** Soluble forms of Be are easily taken up by plants, probably in a manner similar to Ca and Mg, but it is not readily translocated from roots to shoots (Peterson and Girling 1981). Be has been reported to inhibit seed germination, enzyme activation, and uptake of Ca and Mg by roots. Common symptoms are brown, retarded roots and stunted foliage (Romney and Childress 1965).

### 3.1.6 Bismuth

**Experiments conducted in soil.** No reference data were available that describe toxicity of Bi to plants grown in soil.

**Experiments conducted in solution.** There were no primary reference data showing toxicity of Bi to plants grown in solution. The benchmark is based on a report of unspecified toxic effects on plants grown in a solution with the addition of 27 ppm Bi (Bowen, 1979). The authors have low confidence in the benchmark of 20 ppm Bi based on this work alone.

**Mechanism of phytotoxicity.** Although Bi has been shown to reduce the weight of some plants in solution culture (Bowen, 1979), no information on specific mechanisms of toxicity was found.

### 3.1.7 Boron

**Experiments conducted in soil.** John et al. (1977) investigated the effects of B added as  $H_3BO_3$  on shoot weight of corn seedlings grown 7 weeks in muck and two silt loam soils (growth chamber). Addition of 50 ppm B to the muck soil (pH 4.5; % organic matter 56; CEC 117 meq/100g soil) resulted in a 56% reduction in plant growth, while the next lowest concentration tested, 10 ppm B, did not cause a 20% decrease. Growth was reduced 37% by the addition of the lowest concentration tested (0.5 ppm) in the Marble Hill silt loam soil (pH 5.7; % organic matter 6; CEC 23 meq/100g soil). Growth was reduced 83% by the addition of 50 ppm B in the Monroe silt loam soil (pH 5.7; % organic matter 3; CEC 16 meq/100g soil), but not reduced by 10 ppm added B .

Confidence in a benchmark value of 0.5 ppm is low because it is based on fewer than 10 values.

**Experiments conducted in solution.** Wallace et al. (1977b) evaluated the effect of B (as  $H_3BO_3$ ) on leaf, stem, and root weights of bush bean seedlings in solution. After 16 days, root and leaf weights were reduced 35 and 45% by 5.4 ppm B, while 1.1 ppm had no effect.

Bowen (1979) reported unspecified toxic effects on plants grown in a solution with the addition of 1 ppm B.

Confidence in the benchmark of 1 ppm for B is low because it is based on only two values.

**Mechanism of phytotoxicity.** Boron is a plant micronutrient involved in transport of sugars across membranes, synthesis of nucleic acids, and protein utilization. It is rapidly taken up, mainly as the neutral  $B(OH)_3$  molecule and equally distributed between roots and shoots (Wallace and Romney 1977b).

Toxicity symptoms include needle tip necrosis and discoloration in pines (Neary et al. 1975) and burning of leaf edges in other plants. Grasses and legumes appear to have greater than average tolerance



to high B concentrations (Gupta 1984), and pines appear to be particularly sensitive (Stone and Baird, 1956).

### 3.1.8 Bromine

**Experiments conducted in soil.** There were no primary reference data showing toxicity of Br to plants grown in soil. Kabata-Pendias and Pendias (1984) reports unspecified toxic effects on plants grown in a surface soil with the addition of 10 ppm Br. Confidence in this benchmark is low. Newton and Toth (1952) found no toxicity symptoms or reduction in weight of tomato (*Lycopersicon esculentum* L.) at concentrations up to 20 ppm Br in soil.

**Experiments conducted in solution.** There were no primary reference data describing toxicity of Br to plants grown in solution. The benchmark is based on a report of unspecified toxic effects on plants grown with the addition of 15 ppm Br (Martin 1966a). Confidence in the benchmark of 10 ppm based on this work is low.

**Mechanism of phytotoxicity.** Bromine can substitute for part of the Cl<sup>-</sup> requirement of plants. Symptoms of excess Br are similar to those of excess salt (leaf edge necrosis and poor seed germination) (Martin et al. 1956).

### 3.1.9 Cadmium

**Experiments conducted in soil.** Miles and Parker (1979a) investigated the effects of Cd added as CdCl<sub>2</sub> on seed germination and root and shoot weights of a variety of native plants grown from seed for 6 weeks in a sandy soil (pH 4.8, % organic matter 1.9, CEC 6.3 meq/100g soil). Seed germination of Black-eyed Susan (*Rudbeckia hirta*), and root and/or shoot growth of Black-eyed Susan, rough blazing star (*Listris spicata*), long-fruited thimbleweed (*Anemone cylindrica*), and wild bergamot (*Monarda fistulosa*) were reduced by more than 20% with the addition of 10 ppm Cd, the lowest concentration tested. Growth of Kentucky bluegrass (*Poa pretensis*) roots and shoots, little bluestem (*Andropogon scoparius*) roots, and poison-ivy (*Rhus radicans*) roots and shoots was reduced by approximately 90%, 60%, and 70%, respectively, with the addition of 30 ppm Cd where 10 ppm Cd did not have an effect.

Miles and Parker (1979b) found approximately 45% reductions in root and shoot weights of little bluestem grown from seed for 12 weeks in a sandy soil (pH 7.8, % organic matter 2.5, CEC 12 meq/100g soil), when 10 ppm Cd as CdCl<sub>2</sub> was added. This was the only concentration tested.

In a pot culture starting with 2-year-old beech (*Fagus sylvatica* L.), trees growing in an organic-rich forest soil (pH 4.8), Hagemeyer et al. (1993) measured an approximately 25% reduction in annual ring growth in the presence of 5.6 ppm 1M ammonium acetate-extractable Cd when trees were grown for two seasons. Cadmium at 1.1 ppm did not affect growth. The results of this study are not directly comparable to others that report the amount of Cd added to the soil; however, the information is presented for reference to increase the number of plant types covered.

Dixon (1988) measured the response of red oak (*Quercus rubra* L.) seedlings grown for 16 weeks in a sandy loam soil (pH 6, % organic matter 1.5) with addition of Cd (CdCl<sub>2</sub>). Cadmium at 20 ppm reduced tree weight by 28%, while 10 ppm had no effect.

Carlson and Bazzaz (1977) measured root, woody stem, green stem and foliage weights, and main stem diameter of 2 to 3-year old American sycamore (*Plantanus occidentalis* L.) saplings associated

with 90 days of exposure to Cd as CdCl<sub>2</sub> added to a silty clay loam soil. The lowest concentration tested (5 ppm Cd) was responsible for a 30% reduction in leaf weight.

Burton et al. (1984) grew Sitka-spruce (*Picea sitchensis*) seedlings from 4 weeks of age in a mixture of acidic peaty gley soil and sand with Cd added (0.1, 0.4, 1, 2, 4, 8, and 16 ppm as CdCl<sub>2</sub>). Two ppm Cd lead to a reduction of about 45% in root and shoot weight of the 18-week old seedlings.

John et al. (1972b) reported the effects of Cd on radish growth (from seed for 3 weeks in a growth chamber) averaged more than 30 surface soils with the following characteristics: pH of 5.6 (±0.8), % organic matter 12.9 (±15.7), % clay 19.3 ± 14.4, CEC 32.8 meq/100g soil (± 25.2). Root weight was reduced by an average of 67%, and shoot weight by an average of 47%, by the addition of 100 ppm Cd as CdCl<sub>2</sub>, the lowest concentration evaluated.

Reber (1989) found a 21% reduction in wheat growth from seed for 4 weeks in a Phaeosem soil (pH 6.9, % organic matter 2.2) with the addition of 113 ppm Cd as Cd acetate (C<sub>4</sub>H<sub>6</sub>CdO<sub>4</sub>). Only 14 ppm Cd were required to get this same reduction in an acid cambisol soil (pH 5.6, % organic matter 1.7).

In a mixture (1:1) of sandy and clay loam soils (pH 8.4, % organic matter 0.5, CEC 15 meq/100g soil), Singh et al. (1991) measured a 44% reduction in the grain and straw yield of wheat grown from seed to maturity with the addition of 20 ppm Cd as CdCl<sub>2</sub>.

Carlson and Rolfe (1979) found that 100 ppm Cd added as CdCl<sub>2</sub> to a soil was necessary to give a 33% reduction in clipping weight of ryegrass grown in a silt loam soil (pH 5.9, CEC 21 meq/100g soil) from seed.

Number of soybean seeds produced per plant was decreased by 67% when plants were grown in an average garden soil to which 10 ppm Cd was added as CdCl<sub>2</sub> (Aery and Sakar 1991). Cadmium at 5 ppm had no effect. Plants were grown from seed to maturity.

Strickland et al. (1979) evaluated the effects of Cd (1.25, 2.5, 5, 10, and 20 ppm Cd as CdCl<sub>2</sub>) on soybeans grown from seed for 6 weeks in varying ratios of sand and peat. While increasing the amount of organic matter in the mixture from 0 to 2%, the concentration of Cd required to reduce plant growth by 20% was increased from 1.25 ppm (lowest concentration tested) to 20 ppm.

Hassett et al. (1976) measured a 43% reduction in corn root length after 7 days of growth from seed in a loamy sand soil (pH 6.5, % organic matter 2, CEC 2 meq/100g soil) to which 25 ppm Cd (as CdCl<sub>2</sub>) was added. Cadmium at 15 ppm did not affect growth.

Traynor and Knezek (1973) measured a 24% reduction in corn plant weight with the addition of 28 ppm Cd (as CdCl<sub>2</sub>; lowest concentration tested) to a sandy soil (pH 5, % organic matter 2, CEC 6 meq/100g soil) in which the plants had been grown for 5 weeks from seed.

Muramoto et al. (1990) measured the effects on wheat and rice grown from seed to maturity of addition of Cd as CdO to an alluvial soil (pH 6). Root and shoot weights of rice were reduced 32 and 21% by 100 ppm Cd, while 30 ppm had no effect. Wheat grain yield was reduced 34% by 30 ppm Cd, while 10 ppm had no effect.

Sadana and Singh (1987a and b) investigated the effects of Cd added to a loamy sand soil (pH 8.4, % organic matter 1) on lettuce and grain yield of wheat grown from seed to maturity. Lettuce growth was

reduced 23% by the addition of 4 ppm Cd and wheat grain yield was reduced by 28% by 10 ppm Cd (lowest concentration tested).

Miller et al. (1976) investigated the effects of 1, 10, and 100 ppm Cd (as CdCl<sub>2</sub>) on vegetative growth of soybeans from seed for 28 days in soils with a range of pH and CEC values. There was an average of 50% (range 33-77%) reduction in shoot weight of crops grown in three silt loam and one loamy sand soil after the addition of 10 ppm Cd. These soils had pH values ranging from 4.5 to 7.0, and CEC values from 2 to 9 meq/100g soil. Soybeans in one silt loam soil (pH 5.5, CEC 8 meq/100g soil) experienced a 30% reduction in shoot weight after addition of 1 ppm Cd. In another silt loam (pH 6.5, CEC 16 meq/100g soil), a 47% reduction in shoot weight was seen when 100 ppm Cd was added. Corn (*Zea mays* L.) grown from seed for 31 days in a loamy sand used in the 1976 work (pH 6, CEC 2 meq/100g soil) experienced a 28% decrease in plant weight after addition of 2.5 ppm Cd (lowest concentration tested) (Miller et al. 1977).

Bingham et al. (1975) evaluated the effects of a range of Cd concentrations added as CdSO<sub>4</sub> on a variety of horticultural crops grown from seed to maturity in a silt loam soil (pH 7.5, CEC 14 meq/100g soil). Additions of Cd (in ppm) causing a 25% reduction in shoot or reproductive portion weights were as follows: spinach (*Spinacia oleracea* L.) 4, radishes 96, lettuce 13, carrots 20, soybean 5, curlycress (*Lepidium sativum* L.) 8, corn 18, turnip 28, field bean (*Phaseolus vulgaris* L.) 40, wheat 50, tomato and zucchini (*Curcubita pepo* L.) 160, and cabbage 170.

John (1973) evaluated the effects of 40 and 200 ppm Cd added as CdCl<sub>2</sub> on a variety of horticultural crops grown to maturity in a silt loam soil (pH 5.1, % organic matter 12, CEC 38 meq/100g soil). Addition of 40 ppm Cd caused a reduction in plant weights of spinach, peas, and radishes, and grain yield of oats (*Avena sativa* L.) by 96%, 32%, 27%, and 37%, respectively. Addition of 200 ppm Cd caused a reduction in plant weights of lettuce, broccoli (*Brassica oleracea* L.), cauliflower (*Brassica oleracea* L.), and carrots by 90%, 63%, 97%, and 95%, respectively.

Haghiri (1973) determined the effects of additions of 2, 5, 10, and up to 100 ppm Cd as CdCl<sub>2</sub> to a silty clay loam soil (pH 6.7, % organic matter 4, CEC 31 meq/100g soil) on dry matter yield of several crops. Growth of lettuce and radish were reduced 40 and 36% by the lowest treatment level after 37 days and 26 days, respectively. Weights of wheat were reduced by 29% by 5 ppm Cd, and those of soybeans about 50% by addition of 10 ppm Cd. Both crops were grown from seed for 5 weeks.

In two studies using Brown earth soils, Khan and Frankland investigated the effects of Cd added as CdCl<sub>2</sub>, the less soluble CdO, or a combination, on growth of radish (1983, 1984), wheat and oats (1984). Radish root and shoot growth were reduced 22% by the addition of 10 ppm Cd as CdCl<sub>2</sub>, or 100 ppm as CdO (29%), to a soil having a pH of 5.4 (1983). Addition of 50 ppm as CdCl<sub>2</sub>+CdO (1:1) reduced radish root growth 43% in soil having a pH of 4.6 (1984). The plants were grown from seed for 42 days. Wheat growth was reduced 61% by the addition of 50 ppm Cd as CdCl<sub>2</sub> and 47% by the addition of 100 ppm CdO. Oat growth was reduced 25% by the addition of 10 ppm Cd as CdCl<sub>2</sub>. All concentrations were the lowest tested. Wheat and oats were grown from seedlings for 42 days.

Adema and Henzen (1989) calculated EC<sub>50</sub> concentrations for effects of Cd added as CdCl<sub>2</sub> on lettuce, tomato, and oats grown in a growth chamber from seed for 14 days. The EC<sub>50</sub> for lettuce in a humic sand soil (pH 5.1, % organic matter 3.7) was 136 ppm, while in a loam soil (pH 7.4, % organic matter 1.4) it was 33 ppm Cd. The EC<sub>50</sub> for tomato in the humic sand soil was 16 ppm, while in the loam soil it was 171 ppm Cd. The EC<sub>50</sub> for oats in the humic sand soil was 97 ppm, while in the loam soil it was 159 ppm Cd.

Two cultivars of cotton were tested for tolerance to Cd in soil (Rehab and Wallace, 1978). Two-week-old seedlings grown for 35 days in soil (pH 6.8) to which 300 ppm Cd was added (lowest concentration tested) experienced reduced leaf and stem weights—75 and 83% for the first cultivar and 40 and 78% for the second.

Confidence in a benchmark value of 4 ppm Cd is high because of the high number (74) of values available for its derivation. Approximately 40% of the concentrations responsible for greater than 20% reductions in plant growth parameters fall between 1 and 10 ppm Cd added to soil. This range includes wild and cultivated plants such as legumes, trees, grasses, leafy vegetables and other dicotyledonous plants in soils with a relatively wide range of physical and chemical characteristics.

**Experiments conducted in solution.** The effect of Cd, as CdSO<sub>4</sub>, on root elongation of 3-week-old Norway spruce seedlings grown for 7 days in nutrient solution (pH 4) was examined by Lamersdorf et al. (1991). The only concentration tested, 0.11 ppm Cd, reduced root elongation by 23%.

Al-attar et al. (1988) investigated the effect of cadmium acetate on the length and weight of the roots and shoots of perennial ryegrass (*Lolium perenne*) seedlings in solution. A concentration of 0.001 ppm, the lowest concentration tested, resulted in a 20% reduction in root dry weight, a 10% decrease shoot dry weight, and no decrease in shoot or root length.

Kummerova and Brandejsova (1994) studied the toxicity of cadmium (as nitrate) to young maize plants in solution. A concentration of 1.12 ppm, the lowest concentration tested, was associated with a 40% decrease in stalk weight.

Ouzounidou et al. (1997) studied the effect of cadmium (as Cd(NO<sub>3</sub>)<sub>2</sub>) in nutrient solution on root and shoot-leaf length of wheat (*Triticum aestivum*). The lowest concentration tested, 29.8 ppm, resulted in a 53% decrease in root length, a 40% decrease in shoot-leaf length, a 42% reduction in root mass, and a 17% decrease in shoot mass compared to control plants.

Greger et al. (1991) exposed sugar beet (*Beta vulgaris*) seedlings to a series of concentrations of cadmium in a nutrient solution. A reduction of 46% dry shoot weight and 42% dry plant weight occurred at 2.25 ppm. The NOEC was 0.56 ppm. Root weight and growth rate were reduced by less than 20%.

Gussarsson (1994) investigated the effect of cadmium chloride on the growth rate of birch (*Betula pendula*) seedlings in a nutrient solution. The percentage dry weight increase after eight days of cadmium exposure was 20% to 25% lower for shoots exposed to 0.056 ppm than for control shoots. The NOEC was 0.022 ppm. No toxic effect on root growth was observed at concentrations up to 2 ppm, the highest concentration tested.

Godbold and Huttermann (1985) measured a 30% reduction in root elongation rate of 4-wk old Norway spruce seedlings grown in a nutrient solution (pH 4.3) containing 0.56 ppm Cd (CdSO<sub>4</sub>; lowest concentration tested) for 7 d.

Misra et al. (1994) evaluated the effects of several metals on seed germination and root growth of broad bean (*Vicia faba*). Seed germination was not affected by Cd (CdCl<sub>2</sub>) at concentrations up to 10 ppm Cd in nutrient solution. Root elongation of plants exposed for 3 days to 6 ppm Cd was reduced 25%, while 4 ppm had no effect.

Wallace (1979) found 98 and 94% decreases in root and shoot weights of bush bean when grown for 15 days in nutrient solution (pH 5) with 11 ppm Cd (as CdSO<sub>4</sub>), while 0.11 ppm had no effect on growth.

Stiborova et al. (1986) measured a 31% decrease in seedling weight of corn when germinated and grown for 10 days in nutrient solution with 0.11 ppm Cd (as CdSO<sub>4</sub>; lowest concentration tested).

The effect of Cd on weight and grain yield of 2-week-old corn seedlings grown in nutrient solutions containing CdCl<sub>2</sub> was examined in three experiments by Iwai et al. (1975). Seedlings grown for 58 days in pH 5.5 solution experienced 21 and 32% reductions weight and grain yield with 0.1 ppm Cd, while 0.01 ppm had no effect. In an experiment conducted with the same solution but lasting only 19 days, 1 ppm Cd was required to reduce plant weight 23%, while 0.1 ppm had no effect. In a third experiment looking at the effect of pH and toxicity of Cd by using nutrient solutions of pH 4, 5, and 6, plants were grown for 12 days. Plant weight was reduced in all pH treatments by 2 ppm Cd (37, 41, and 45% reductions, respectively) while 0.2 ppm had no effect.

El-Enany (1995) measured 25, 32, and 39% decreases in seed germination, radicle length, and fresh plant weight of corn when germinated and grown for 5 d in nutrient solution with 45 ppm Cd (as CdCl<sub>2</sub>; lowest concentration tested).

Rascio et al. (1993) found reductions of approximately 45 and 35% in root and shoot length of corn seedlings grown 18 days in nutrient solution containing 28.1 ppm Cd (as Cd(NO<sub>3</sub>)<sub>2</sub>). Cadmium concentration of 11.2 ppm had no effect.

Wong and Bradshaw (1982) measured 37 and 27% decreases in lengths of longest root and shoot of ryegrass when germinated and grown for 14 days in nutrient solution (pH 7) with 1.25 ppm Cd (as CdSO<sub>4</sub>; lowest concentration tested).

Patel et al. (1976) found 55 and 24% decreases in root and stem weights of chrysanthemum seedlings when grown for 21 days in nutrient solution with 0.11 ppm Cd (as CdSO<sub>4</sub>; lowest concentration tested).

Cunningham et al. (1975) examined the effect of Cd on leaf, stem, and root weight of 4-day-old soybean seedlings grown for 21 days in nutrient solution (pH 5.2). A concentration of 0.05 ppm (the lowest concentration tested) reduced leaf, stem and root weights by 73, 62, and 38%, respectively.

In 1977, Cunningham reported 56, 47, and 53% reductions in leaf, stem, and root weights of soybean seedlings when grown for 21 days in nutrient solution (pH 6.2) with 0.05 ppm Cd (as Cd(NO<sub>3</sub>)<sub>2</sub>; lowest concentration tested).

Adema and Henzen (1989) evaluated the effect of Cd (as CdCl<sub>2</sub>) on germination and growth of lettuce, tomato, and oat seedlings in nutrient solution. They report 50% reductions in top growth weight at 0.84 ppm for lettuce, 3 ppm for tomato, and 6 ppm for oats.

Turner (1973) grew seedlings of various vegetables in nutrient solution (pH 6.3) containing Cd at 0.01, 0.1, and 1 ppm Cd as CdCl<sub>2</sub>. Carrots were the least tolerant with a 25% reduction in top weight at 0.01 ppm Cd after 35 days. Tomato seedlings grown for 14 days showed a 45% reduction in top weight with 0.1 ppm. Beets (*Beta vulgaris* L.) and Swiss chard (*Beta vulgaris* L.), both grown for 35 days, had reductions of 54% in top weight at 1 ppm Cd.

Inouhe et al. (1994) grew 7-d old seedlings of various crops in solution containing Cd as CdSO<sub>4</sub>. Sesame (*Sesamum indicum*), pea, radish, cucumber, tomato, and Azuki bean (*Vigna angularis*) experienced reduced root growth (40 to 85%) at 1 ppm Cd, the lowest concentration tested. Lettuce and barley were more tolerant with 63 and 35% reductions at 3.4 ppm Cd after 7 d. Root growth of oat seedlings was reduced 50% when grown in solution containing 6.8 ppm Cd.

Jalil et al. (1994) measured a 24% reduction in shoot dry weight of 2-d old wheat seedlings when grown for 13 d in nutrient solution (pH 5.5) with 0.1 ppm Cd (as CdCl<sub>2</sub>; lowest concentration tested).

Wang (1994) evaluated the effects of several metals and organic compounds on radicle weight of rice seed germinated and grown for 6 d in solution. The calculated EC<sub>50</sub> for Cd (CdCl<sub>2</sub>) was 1.4 ppm.

Page et al. (1972) grew corn, field bean, beet, and turnip seedlings for 21 days in nutrient solution containing Cd as CdSO<sub>4</sub> at 0.1, 0.25, 0.5, and 1 ppm. Weights of bean, beet, and turnip were reduced 36, 45, and 22% by 0.1 ppm Cd. Weight of corn was reduced 33% by 0.5 ppm. They also grew lettuce, tomato, pepper, cabbage, and barley for 21 days in solutions containing 1, 2.5, and 5 ppm Cd. Weights of lettuce, tomato, pepper, and barley were reduced 53, 25, 38, and 30% by 1 ppm. Cabbage had a 24% reduction in weight with 2.5 ppm Cd.

Garate et al. (1993) evaluated the effect of Cd (as CdSO<sub>4</sub>) in nutrient solution on root and leaf growth of lettuce and endive (*Lactuca serriola* L.). They found a 28% reduction in root weight of giant endive after 35 days of growth with 0.1 ppm Cd (lowest concentration tested).

The concentrations of Cd, from CdCl<sub>2</sub>, required for a 50% reduction in seed germination and root length of mustard after a 3-day exposure in solution (pH 6.6) was reported by Fargasova (1994). LC50 for germination was 692 ppm and EC<sub>50</sub> for root length was 48 ppm Cd.

The effect of Cd, as CdCl<sub>2</sub>, on plant weight of cotton grown in nutrient solution (pH 5.5) was evaluated by Rehab and Wallace (1978). Plant weight was reduced 47% by 1.1 ppm Cd, the lowest concentration tested.

Confidence in the 0.1 ppm benchmark is high. It is based on 52 values from experiments using a variety of plant species.

**Mechanism of phytotoxicity.** Cadmium is not essential for plant growth. If present in available form, it is readily taken up by the roots and translocated through the plant and accumulated. Cadmium is chemically similar to Zn, an essential element. Competition between the two for organic ligands may explain some of the toxic effects of Cd and the ameliorative effects of Zn on Cd toxicity. Cadmium depresses uptake of Fe, Mn, and probably Ca, Mg, and N (Wallace et al., 1977e; Iwai, et al. 1975). Cadmium is toxic at low concentrations. Symptoms resemble Fe chlorosis and include necrosis, wilting, reduced Zn levels, and reduction in growth. The mechanisms of toxicity include reduced photosynthetic rate, poor root system development, reduced conductivity of stems, and ion interactions in the plant. Agronomic crops are more sensitive to Cd toxicity than trees (Adriano 1986).

### 3.1.10 Chromium

**Experiments conducted in soil.** Turner and Rust (1971) investigated the effect of Cr added as Cr(VI) on soybean seedlings grown 3 days in a loam soil. Fresh shoot weight was reduced 30% by 30 ppm Cr, while 10 ppm had no effect.

Adema and Henzen (1989) calculated  $EC_{50}$  concentrations for effects of Cr added as Cr(VI) on lettuce, tomato and oats grown in a growth chamber from seed for 14 days. The  $EC_{50}$  for lettuce in a humic sand soil (pH 5.1, % organic matter 3.7) was greater than 11 ppm, while in a loam soil (pH 7.4, % organic matter 1.4) it was 1.8 ppm Cr. The  $EC_{50}$  for tomato in the humic sand soil was 21 ppm, while in the loam soil it was 6.8 ppm Cr. The  $EC_{50}$  for oats in the humic sand soil was 31 ppm, while in the loam soil it was 7.4 ppm Cr.

Confidence in the benchmark of 1 ppm Cr is low because of the small number of studies on which it is based.

**Experiments conducted in solution.** Adema and Henzen (1989) calculated  $EC_{50}$  concentrations for effects of Cr added as  $K_2Cr_2O_7$  (Cr VI) on lettuce, tomato and oats grown in a growth chamber from seed for 14 days. The  $EC_{50}$  values for lettuce, tomato and oats were 0.16, 0.29, and 1.4 ppm Cr.

Wang (1994) evaluated the effects of several metals and organic compounds on radicle weight of rice seed germinated and grown for 6 d in solution. The calculated  $EC_{50}$  for Cr (VI) ( $K_2CrO_4$ ) was 4.8 ppm.

Moral et al. (1995) measured 24 and 32% reductions in root length and fresh root weight of tomato seedlings grown in nutrient solution containing 100 ppm Cr (III) as  $CrCl_3$ . Chromium at 50 ppm had no effect.

Top weight of soybean seedlings grown for 5 d in nutrient solution containing Cr(VI) was reduced 21% by 1 ppm Cr, while 0.5 ppm had no effect (Turner and Rust, 1971).

Wallace et al. (1977a) measured a 30% reduction in leaf weight of bush beans grown 11 d in nutrient solution containing 0.54 ppm Cr as Cr(VI) ( $K_2Cr_2O_7$ ).

Length of the longest root of rye grass was reduced 69% by exposure to 2.5 ppm Cr(VI) ( $K_2Cr_2O_7$ ; lowest concentration tested) in nutrient solution (pH 7) for 14 d (Wong and Bradshaw, 1982). Length of the longest shoot was not affected at this concentration.

Breeze (1973) found little difference in the toxicity of Cr(III) [ $Cr_2(SO_4)_3$ ] and Cr(VI) ( $K_2Cr_2O_7$ ) to rye grass seed germination. Seed exposed to solutions containing 50 ppm Cr (III) or (VI) reduced germination 37 and 38% after 2.5 days.

Nutrient solution containing 0.05 ppm Cr(III) [ $Cr_2(SO_4)_3$ ] reduced leaf and stem weights of chrysanthemum seedlings exposed for 21 days by 31 and 36% (Patel et al., 1976). This was the lowest concentration tested and root weight was not affected.

Using a 1:1 combination of Cr(III) ( $CrCl_3$ ) and Cr(VI) ( $K_2CrO_7$ ) in nutrient solution (pH 5), Hara et al. (1976) measured a 68% reduction in weight of cabbage with 10 ppm Cr. Chromium at 2 ppm had no effect.

The concentrations of Cr(VI), from  $(NH_4)_2CrO_4$ , required for a 50% reduction in seed germination and root length of mustard after a 3-day exposure in solution (pH 7.3), was reported by Fargasova (1994).  $LC_{50}$  for germination was 100 ppm and  $EC_{50}$  for root length was 46 ppm Cr.

Confidence in the solution Cr benchmark of 0.05 ppm is moderate, however the concentration approximating the 10th percentile was the lowest concentration tested and caused a greater than 30% reduction in the growth parameter.

**Mechanism of phytotoxicity.** Chromium is not an essential element in plants. The (VI) form is more soluble and available to plants than the (III) form and is considered the more toxic form (Smith et al. 1989). In soils within a normal Eh and pH range, Cr(VI), a strong oxidant, is likely to be reduced to the less available Cr(III) form although the (III) form may be oxidized to the (VI) form in the presence of oxidized Mn (Bartlett and James 1979). In nutrient solution, however, both forms are about equally taken up by plants and toxic to plants (McGrath 1982). Cr(VI), as  $\text{CrO}_4^{2-}$ , may share a root membrane carrier with  $\text{SO}_4^{2-}$ . Cr(VI) is more mobile in plants than Cr(III) but translocation varies with plant type. After plant uptake it generally remains in the roots because of the many binding sites in the cell wall capable of binding especially the Cr(III) ions (Smith et al. 1989). Within the plant Cr(VI) may be reduced to the Cr(III) form and complexed as an anion with organic molecules. Symptoms of toxicity include stunted growth, poorly developed roots, and leaf curling. Chromium may interfere with C, N, P, Fe, and Mo metabolism, and enzyme reactions (Kabata-Pendias and Pendias 1984).

### 3.1.11 Cobalt

**Experiments conducted in soil.** There was no primary reference data showing toxicity of Co to plants grown in soil. Kabata-Pendias and Pendias (1984) reported unspecified toxic effects on plants grown in a surface soil with the addition of 20 ppm Co. We have low confidence in the benchmark based on this study alone.

**Experiments Conducted in Solution.** Wallace et al. (1977a) evaluated the effect of Co as  $\text{CoSO}_4$  on bush beans grown for 21 d in nutrient solution. Leaf dry weight was reduced 22% by the addition of 0.06 ppm Co, the lowest concentration tested. Root and stem weights were not affected at this concentration. Chrysanthemum seedling root weight was reduced 55% after 21 days of growth in nutrient solution containing the same concentration of Co as  $\text{CoSO}_4$  (Patel et al. 1976). Leaf and stem weight were not affected at this concentration.

Misra et al. (1994) evaluated the effects of several metals on seed germination and root growth of broad bean (*Vicia faba*). Seed germination was not affected by Co ( $\text{CoCl}_2$ ) at concentrations up to 10 ppm Co in nutrient solution. Root elongation of plants exposed for 3 days to 10 ppm Co was reduced 30%, while 8 ppm had no effect.

Patterson and Olson (1983) evaluated the effect of several metals in solution (pH 5 to 6) on white spruce (*Picea glauca*), black spruce (*Picea mariana*), paper birch (*Betula papyrifera*), jack pine (*Pinus banksiana*), white pine (*Pinus strobus*), red pine (*Pinus resinosa*), and honeysuckle (*Lonicera tatarica*) grown from seed for 5 to 21 d. There was no clear relationship between general tree group (i.e., spruce or pine) and effect of Co on radicle elongation. Toxic concentrations ranged from 5 ppm (35 and 47% reductions in honeysuckle and paper birch) to 100 ppm (53% reductions in white pine). The other species were intermediate in their response to Cu in solution.

Confidence in the solution benchmark of 0.06 ppm Co is low because it is based on a limited number of types of plants.

**Mechanism of phytotoxicity.** Cobalt is not known to be essential to plants except legumes in symbiosis with  $\text{N}_2$ -fixing microorganisms. When translocated from roots it travels in the xylem as the Co(II) ion (Tiffin 1967). Toxicity symptoms due to excess Co are typical of Fe deficiency induced



chlorosis and necrosis, and root tip damage (Wallace et al. 1977a). There appears to be inhibition of mitosis and chromosome damage (Aller et al. 1990).

### 3.1.12 Copper

**Experiments conducted in soil.** Miles and Parker (1979b) found approximately 68% reductions in root and shoot weights of little bluestem grown from seed for 12 weeks in a sandy soil (pH 7.8, % organic matter 2.5, CEC 12 meq/100g soil), when 100 ppm Cu as  $\text{CuSO}_4$  was added. This was the only concentration tested. Growth was reduced in a second sandy soil (pH 4.8, % organic matter 1.9, CEC 6 meq/100g soil) by 86% with the addition of 100 ppm Cu (only concentration tested).

Wallace et al. (1977b) evaluated the effects of Cu, added as  $\text{CuSO}_4$  to a loam soil, on leaf and stem weights of bush beans grown from seed for 17 days. Leaf weight was reduced 26% by 200 ppm Cu, while 100 ppm had no effect.

Confidence is low in the benchmark of 100 ppm Cu in soil because it is derived from fewer than 10 values.

**Experiments in solution.** The effect of Cu on stem diameter increase and plant weight of red pine, maple (*Acer rubrum*), dogwood (*Cornus stolonifera*), and honeysuckle was examined by Heale and Ormrod (1982). All seedlings (90-d old) grown for 110 d in nutrient solution containing 4 ppm Cu from  $\text{CuSO}_4$  (lowest concentration tested) were affected. Reductions in rate of stem diameter increase and in plant weight were 41 and 50%, 79 and 67%, and 97 and 74% for maple, dogwood, and honeysuckle, respectively. Red pine experienced a 28% decrease in plant weight at 4 ppm Cu but the stem diameter increase was unaffected up to 20 ppm Cu (highest concentration tested).

Mocquot et al. (1996) investigated the toxicity of copper (as  $\text{CuSO}_4$ ) to maize (*Zea mays*) seedlings in solution. At 0.64 ppm of copper, shoot length and root mass were decreased by about 23% and 42%, respectively. At 0.45 ppm of Cu, no toxicity was observed.

Patterson and Olson (1983) evaluated the effect of several metals in solution (pH 5 to 6) on white spruce, black spruce, paper birch, jack pine, white pine, red pine, and honeysuckle grown from seed for 5 to 21 d. Paper birch, which was least tolerant of Co, was also most affected by Cu, with a 39% reduction in radicle elongation at 1 ppm Cu. As in the case of Co, white pine was the most tolerant plant tested with a 42% reduction in radicle elongation at 100 ppm Cu. The other species were intermediate in their response to Cu in solution.

Wang (1994) evaluated the effects of several metals and organic compounds on radicle weight of rice seed germinated and grown for 6 d in solution. The calculated  $\text{EC}_{50}$  for Cu ( $\text{CuCl}_2$ ) was 0.22 ppm.

Wong and Bradshaw (1982) measured reductions in lengths of longest roots and shoots of rye grass grown for 14 d in nutrient solution (pH 7) to which Cu as  $\text{CuSO}_4$  was added. The length of the longest root was reduced 71% by 0.031 ppm Cu, the lowest concentration tested.

Maize seedlings germinated and grown for 10 d in solution containing  $\text{CuSO}_4$  had a 40% reduction in total fresh weight in the 0.06 ppm Cu treatment (lowest concentration tested) (Stiborova et al. 1986). This same concentration caused a 45% reduction in root weight of chrysanthemums grown for 21 d in nutrient solution with  $\text{CuSO}_4$  added (Patel et al. 1976). Leaf and stem weights were not affected.

Gupta and Mukherji (1977) evaluated the effect of Cu as a  $\text{CuSO}_4$  solution on rice seedling shoot and root lengths. After 4 days, root length was reduced 64% by 64 ppm Cu, while 6.4 ppm had no effect.

Confidence in the solution benchmark for Cu, 0.06 ppm, is moderate; however it is based on a limited number of types of plants.

**Mechanism of Phytotoxicity.** Copper is a micronutrient essential for plant nutrition. It is required as a co-factor for many enzymes and is an essential part of a copper protein involved in photosynthesis. Copper occurs as part of enzymes and enzyme systems. Root absorption appears to be passive, perhaps in organo-copper complexes (Jarvis and Whitehead 1983), and active through a specific carrier (Fernandes and Henriques 1991). Copper may be deficient in low-copper soils because the metal is adsorbed to cells in the root system. The form in which it is taken into the root affects its binding there (Wallace and Romney 1977b). Copper can be transported in the xylem and phloem of plants complexed with amino acids.

The most common toxicity symptoms include reduced growth, poorly developed root system, and leaf chlorosis (Wong and Bradshaw 1982). The basic deleterious effect of Cu is related to the root system where it interferes with enzyme functioning (Mukherji and Das Gupta 1972). It also strongly interferes with photosynthesis and fatty acid synthesis (Smith et al. 1985).

### 3.1.13 Fluorine

**Experiments conducted in soil.** The benchmark is based on a report of unspecified reductions in plant growth in a surface soil with the addition of 200 ppm F (Kabata-Pendias and Pendias 1984). Confidence in the benchmark for F is low because it based on this reference alone.

**Experiments conducted in solution.** Bowen (1979) reported unspecified reductions in plant growth in a solution culture with the addition of 5 ppm F. We have low confidence in the benchmark based on this study alone.

**Mechanism of Phytotoxicity.** Fluorine is not an essential plant element. Toxicity symptoms are the same as seen in plants exposed to HF gas; marginal leaf chlorosis and interveinal chlorosis (Brewer 1966).

### 3.1.14 Iodine

**Experiments conducted in soil.** Newton and Toth (1952) measured the effects of I, added to soils (pH 6.8) as KI at 0.4 and 4 ppm, on top weight of tomatoes grown from seed for 97 days. They found a 47% reduction in top weight in a sandy soil, 25% in one loam soil and 52% in another, and 30% reduction in top weight in a silt loam soil at 4 ppm I.

The benchmark of 4 ppm is taken from this study. Confidence in this benchmark is low.

**Experiments conducted in solution.** Top weight of corn seedlings grown for 60 days in nutrient solution (pH 5.8) was reduced 31% by the addition of 0.5 ppm I added as KI (Lewis and Powers 1941). Iodine at 0.1 ppm had no effect on plant growth.

Newton and Toth (1952) measured the effects of I, added to nutrient solution as KI at 0.5 and 5 ppm, on top weight of tomato seedlings grown for 60 days. Iodine at 5 ppm reduced top weight 46%, while 0.5 ppm had no effect.

Confidence in the solution benchmark of 0.5 ppm I is low because of the limited amount of data on which it is based.

**Mechanism of phytotoxicity.** Iodine is not known to be essential for plant growth. It is present in available form in very small amounts in soil. Toxicity symptoms are similar to salt burn, that is, burning of leaf edges and subsequent leaf necrosis (Martin 1966b).

### 3.1.15 Iron

**Experiments conducted in soil.** No information was found on which to base a toxicity benchmark for plants growing in soil.

**Experiments conducted in solution.** Wallihan (1966) reported unspecified reductions in plant growth in a solution culture with the addition of 10 ppm Fe. Wallace et al. (1977b) evaluated the effects of Fe (as  $\text{FeSO}_4$ ) on leaf, stem, and root weights of bush bean seedlings grown for 15 days in nutrient solution. Iron at 28 ppm reduced all three measures 67, 52, and 67%, respectively, while 11.2 ppm had no effect.

After 55 days cabbage seedling plant weight was reduced 45% by 50 ppm Fe added as  $\text{FeSO}_4$  to nutrient solution (pH 5), while 10 ppm had no effect on growth (Hara et al. 1976).

Confidence in the benchmark for Fe in solution (10 ppm) is low because it is based on less than 10 values.

**Mechanism of phytotoxicity.** Iron is the key metal required for energy transformations needed for cellular functioning. It occurs in heme and nonheme proteins and is concentrated in chloroplasts. Organic Fe complexes are involved in photosynthetic electron transfer. Plant symptoms of toxicity are not specific and differ among plant species and growth stages (Foy et al. 1978).

### 3.1.16 Lead

**Experiments conducted in soil.** Rolfe and Bazzaz (1975) measured the effects of Pb, added to a 1:1:1 mixture of soil, sand and peat moss as  $\text{PbCl}_2$ , on 1-year-old seedlings of autumn olive (*Elaeagnus umbellata*) grown for 49 days. They found a reduction in transpiration of approximately 25% with the addition of 160 ppm Pb, while 80 ppm had no effect.

Dixon (1988) measured the response of red oak seedlings grown for 16 weeks in a sandy loam soil (pH 6, % organic matter 1.5) with addition of Pb ( $\text{PbCl}_2$ ). Lead at 50 ppm reduced tree weight by 26%, while 20 ppm had no effect.

Carlson and Bazzaz (1977) measured foliage biomass, trunk diameter, and new stem and root growth reductions in 2- to 3-year-old American sycamore saplings associated with a 90-day exposure to Pb as  $\text{PbCl}_2$  added to a silty clay loam soil. The lowest concentration tested (50 ppm Cd) was responsible for a 30% reduction in leaf weight.

Burton et al. (1984) grew Sitka-spruce seedlings from 4 weeks of age in a mixture of acidic peaty gley soil and sand with Pb added as  $\text{PbCl}_2$ . Lead added at 100 ppm resulted in a reduction of about 25% in root and shoot weight of the 18-week old seedlings.

Miles and Parker (1979b) found approximately 52% reductions in root and shoot weights of little bluestem grown from seed for 12 weeks in a sandy soil (pH 7.8, % organic matter 2.5, CEC 12 meq/100g soil), when 450 ppm Pb as  $\text{PbCl}_2$  was added. This was the only concentration tested. Root growth was reduced in a second sandy soil (pH 4.8, % organic matter 1.9, CEC 6 meq/100g soil) by 22% with the addition of 450 ppm Pb (only concentration tested).

Carlson and Rolfe (1979) found that 5000 ppm Pb added as  $\text{PbCl}_2$  to a soil was necessary to give 46 and 31% reductions in clipping weight of ryegrass and fescue (*Festuca rubra*) grown in a silt loam soil (pH 5.9, CEC 21 meq/100g soil) from seed.

Muramoto et al. (1990) measured the effects of addition of Pb as PbO to an alluvial soil (pH 6) on growth and yield of wheat grown from seed to maturity. Root weight was reduced 22% by 1000 ppm Pb, while 300 ppm had no effect.

In a study using Brown earth soil, Khan and Frankland (1984) investigated the effects of Pb added as  $\text{PbCl}_2$ , the less soluble PbO, or a combination, on root weight of wheat and oats. Wheat root weight was reduced 34% by the addition of 1000 ppm Pb as  $\text{PbCl}_2$ , while 500 ppm had no effect. Oat growth was reduced 37% by the addition of 500 ppm Pb as  $\text{PbCl}_2$ , while 100 ppm had no effect. Wheat and oats were grown from seedlings for 42 days.

Hassett et al. (1976) measured a 48% reduction in corn root length after 7 days of growth from seed in a loamy sand soil (pH 6.5, % organic matter 2, CEC 2 meq/100g soil) to which 500 ppm (as  $\text{PbCl}_2$ ) was added. Lead at 250 ppm did not affect growth.

Corn (*Zea mays* L.) grown from seed for 31 days in a loamy sand used in the 1976 work (pH 6, CEC 2 meq/100g soil) experienced a 42% decrease in plant weight after addition of 250 ppm Pb (Miller et al., 1977). Lead at 125 ppm did not affect growth.

In a study using Brown earth soil, Khan and Frankland (1983) investigated the effects of Pb added as  $\text{PbCl}_2$ , the less soluble PbO, or a combination, on radish root and top weights. Radish root growth was reduced 24% by the addition of 500 ppm Pb as  $\text{PbCl}_2$ , or 1000 ppm (lowest concentration tested) as PbO (27% reduction), to a soil having a pH of 5.4. Plants were grown from seed for 42 days.

John and Van Laerhoven (1972) investigated the effects of lead, added in various forms, to a silty clay loam soil (pH 3.8, % organic matter 17, CEC 45 meq/100g soil). Lettuce was grown from seed for 30 days before tops were harvested. Lead added at a rate 1000 ppm (lowest concentration tested) as  $\text{PbCl}_2$  and  $\text{Pb}(\text{NO}_3)_2$  reduced plant weight by 35 and 25%.

Moderate confidence is assumed for the 50 ppm benchmark for Pb because it is based on 17 values from experiments conducted with a range of different plant species.

**Experiments conducted in solution.** The effect of Pb, as  $\text{PbCl}_2$ , on root elongation of 3-week-old Norway spruce seedlings grown for 7 days in nutrient solution (pH 4) was examined by Lamersdorf et al. (1991). The only concentration tested, 0.02 ppm Pb, reduced root elongation by 26%.

Godbold and Kettner (1991) measured a 24% reduction in rate of root elongation of 3-wk old Norway spruce seedlings grown in a nutrient solution (pH 4) containing 0.2 ppm Pb ( $\text{PbCl}_2$ ; lowest concentration tested) for 7 d.

The rate of root growth of onions grown for 4 d in a solution containing lead as  $\text{Pb}(\text{NO}_3)_2$  was reduced 33% by 0.2 ppm Pb, while 0.02 ppm had no effect (Liu et al. 1994).

Wang (1994) evaluated the effects of several metals and organic compounds on radicle weight of rice seed germinated and grown for 6 d in solution. The calculated  $\text{EC}_{50}$  for Pb [ $\text{Pb}(\text{C}_2\text{H}_3\text{O}_2)_2$ ] was 9.7 ppm.

Mhatre and Chaphekar (1982) tested several species at germination stage for their response to Pb. Seeds of sorghum, alfalfa, mung bean, cluster bean, and radish were allowed to germinate in solutions containing 0.001, 0.01, 0.1, or 1 ppm Pb as  $\text{Pb}(\text{NO}_3)_2$ . Germination counts after 24 hours showed no effect of Pb. After 5 days, root length of cluster bean was reduced 34% by 0.001 ppm Pb. Root length of alfalfa was reduced 25% by the addition of 0.1 ppm. Root and shoot lengths of radish were reduced 27 and 32% by the addition of 1 ppm Pb. Root length of mung bean was reduced 23% by the addition of 1 ppm.

The effect of Pb on root length of barley and maize seedlings after a 7-day exposure in nutrient solution was examined by Wierbicka and Antosiewicz (1993). Root length of maize was reduced 25% by 1 ppm Pb (lowest concentration tested) and that of barley reduced 27% by 2 ppm Pb, while 1 ppm had no effect.

Wong and Bradshaw (1982) evaluated the effect of Pb on root and shoot elongation of rye grass grown in solution (pH 7) with Pb added as  $\text{Pb}(\text{NO}_3)_2$ . After 14 days they found 77 and 36% reductions in lengths of the longest roots and shoots in response to 2.5 ppm Pb (lowest concentration tested).

Wong and Lau (1985) evaluated the effect of Pb on root length of several cultivars of Bermuda grass and wire grass (*Eleusine indica* L.) grown in solution with Pb added as  $\text{Pb}(\text{NO}_3)_2$ . After 14 days they found root length of all Bermuda grass cultivars reduced an average of 64% in response to 10 ppm Pb (lowest concentration tested). The response of wire grass was more variable with 75 and 27% reductions in root length at 10 ppm for two cultivars, and a 87% reduction at 20 ppm for the third (10 ppm had no effect).

Hooper (1937) ran a series of experiments to evaluate the effect of Pb as  $\text{PbSO}_4$  on fresh weight of french beans grown in nutrient solution. In three of the runs she found an average 32% reduction in response to 10 ppm Pb, while 5 ppm had no effect. In two other runs, fresh weight was reduced approximately 25% by 30 ppm, while 20 ppm Pb had no effect.

The fresh weight of maize seedlings grown for 10 days in a Pb-containing solution ( $\text{Pb}(\text{NO}_3)_2$ ) was reduced 45% by 207 ppm Pb, while 20.7 ppm had no effect (Stiborova et al. 1986).

The concentrations of Pb, from  $\text{Pb}(\text{CH}_3\text{COO})_2$ , required for a 50% reduction in seed germination and root length of mustard after a 3-day exposure in solution (pH 5.5), was reported by Fargasova (1994).  $\text{LC}_{50}$  for germination was 1148 ppm and  $\text{EC}_{50}$  for root length was 263 ppm Pb.

Confidence in the 0.02 ppm toxicity benchmark for plants growing in solution is moderate.

**Mechanism of phytotoxicity.** Lead is taken up passively by roots and translocation to shoots is limited (Wallace and Romney 1977b). It is bound to the outside of roots, in the apoplast, and in cell walls and organelles of absorbing roots (Koepppe 1981). In the plant, lead may exist in naturally chelated form, or in pyro- or orthophosphate forms. The phytotoxicity of lead is relatively low compared with

other trace elements. It affects mitochondrial respiration and photosynthesis by disturbing electron transfer reactions (Miles et al. 1972).

### 3.1.17 Lithium

**Experiments conducted in soil.** Wallace (1979) investigated the effects of Li ( $\text{Li}_2\text{C}_2\text{O}_4$ ) added to a loam soil (pH 6) on barley seedlings grown for 10 days. Lithium at 500 ppm (lowest concentration tested) resulted in a 66% reduction in shoot weight.

Wallace et al. (1977c) measured the reduction in leaf and stem weights of cotton and bush bean seedlings resulting from additions of Li, as LiCl or  $\text{LiNO}_3$ , to a loam soil (pH 6). Cotton leaf and stem weights were reduced 33 and 56% by the addition of 50 ppm Li as  $\text{LiNO}_3$ , while 25 ppm had no effect. Bush bean leaf weight was reduced 32% by the addition of 25 ppm Li as LiCl, while 10 ppm had no effect. Cotton was exposed for 16 days and bush beans for 10 days.

Aldrich et al. (1951) recorded an undefined phytotoxic effect on sweet orange seedlings grown in a surface soil for 6 months with 2 ppm Li as  $\text{LiSO}_4$  (lowest concentration tested).

Confidence in the benchmark of 2 ppm is low.

**Experiments conducted in solution.** Wallace et al. (1977c) measured the reduction in leaf, stem, and root weights of bush bean seedlings resulting from additions of Li, as  $\text{LiNO}_3$ , to nutrient solution. Stem weight was reduced 30% by 3.5 ppm Li, the lowest concentration tested.

Confidence in the 3 ppm toxicity benchmark for plants growing in solution is low.

**Mechanism of phytotoxicity.** The soluble Li in soils is easily taken up by plants. It appears to share the  $\text{K}^+$  transport carrier and is mainly found in leaf tissues. Toxicity symptoms include damage to root tips and necrosis of interveinal leaf tissue (Aldrich et al. 1951).

### 3.1.18 Manganese

**Experiments conducted in soil.** Wallace et al. (1977b) evaluated the effects of Mn, added as  $\text{MnSO}_4$  to a loam soil, on leaf and stem weights of bush beans grown from seed for 17 days. Stem weight was reduced 29% by 500 ppm Mn (lowest concentration tested).

Because the 500 ppm benchmark for Mn is based on this one study, confidence in it is low.

**Experiments conducted in solution.** Langheinrich et al. (1992) evaluated effects of solution pH, N supply and Mn ( $\text{MnSO}_4$ ) on growth parameters of Norway spruce seedlings. In an experiment run at pH 6 for 32 d, root growth was measured (length and weight). Manganese added at 44 ppm (lowest concentration tested) reduced root growth 50% when N was added as  $\text{NH}_4$ , and reduced root length by 37% when N was added as  $\text{NO}_3$  (11 ppm had no effect). In experiments run at pH 4 for 77 days, epicotyl height, length of the primary root, and percent plants with terminal buds were determined. Manganese added at 44 ppm (only concentration tested) reduced all measures approximately 40% when N was added as  $\text{NO}_3$  and reduced height of epicotyl and percent plant with terminal buds by approximately 55% when N was added as  $\text{NH}_4$ .

Wong and Bradshaw (1982) evaluated the effect of Mn on root and shoot elongation of rye grass grown in solution (pH 7) with Mn added as  $\text{MnSO}_4$ . After 14 days, they found a 71% reduction in the length of the longest root in response to 0.75 ppm (lowest concentration tested).

Wallace et al. (1977b) evaluated the effect of Mn (as  $\text{MnSO}_4$ ) on leaf, stem, and root weights of bush bean seedlings in grown in nutrient solution. After 16 days, in one experiment, the three weights were reduced approximately 25% by 5.5 ppm Mn, the lowest concentration tested. In a second, 21-day experiment, the three weights were reduced approximately 40% by 55 ppm, while 5.5 ppm Mn had no effect.

LeBot et al. (1990) evaluated the effect of Mn, as  $\text{MnSO}_4$ , on weight of tomato plants growing in nutrient solution (pH 5.5) for 17 days. Manganese at 5.5 ppm reduced plant weight by 27%, while 2.8 ppm had no effect.

Foy et al. (1995) evaluated the effect of pH and plant genotype on the toxicity of Mn ( $\text{MnSO}_4$ ) on 9-d old seedlings of cotton grown in nutrient solution for 18 d. Leaf and root dry weights of one genotype were reduced approximately 25% at 8 ppm in the pH 5 solution, while in the pH 4.6 solution, Mn was not toxic until a concentration of 16 ppm was reached (35 to 77% reductions). For the second genotype, the lowest concentration of Mn tested (4 ppm) reduced plant weight approximately 35% at both pH levels.

Wang (1994) evaluated the effects of several metals and organic compounds on radicle weight of rice seed germinated and grown for 6 d in solution. The calculated  $\text{EC}_{50}$  for Mn ( $\text{MnCl}_2$ ) was 100 ppm.

Burke et al. (1990) compared the effects of 30 days of hourly root submersion in an Mn in solution ( $\text{MnSO}_4$ , pH 4.8) on root and shoot weights of five wheat cultivars. For three of the five cultivars, root weight was reduced an average of 43% (25 to 60%) by the addition of 30 ppm Mn, the lowest concentration. This concentration reduced both root and shoot weight of a fourth cultivar by 25%. The fifth cultivar experienced 60 and 35% reductions in root and shoot weight with the addition of 90 ppm Mn, while 30 ppm had no effect.

The effect of Mn on weight of potato (*Solanum tuberosum* L.) shoots grown for 32 days in nutrient solution was examined by Marsh and Peterson (1990). A concentration of 33.5 ppm (lowest concentration tested) caused a 23% reduction in shoot weight.

Confidence in the solution Mn benchmark of 4 ppm is moderate.

**Mechanism of phytotoxicity.** Manganese is essential for plant growth. It is involved in N assimilation, as a catalyst in plant metabolism and functions with Fe in the synthesis of chlorophyll (Labanauskas 1966). Toxicity symptoms include marginal chlorosis and necrosis of leaves and root browning. Excess Mn interferes with enzymes, decreases respiration, and is involved in the destruction of auxin (Foy et al. 1978). It is fairly uniformly distributed between roots and shoots (Wallace and Romney 1977b).

### 3.1.19 Mercury

**Experiments conducted in soil.** There were no primary reference data describing toxicity of Hg to plants grown in soil. Kabata-Pendias and Pendias (1984) report unspecified toxic effects on plants grown in a surface soil with the addition of 0.3 ppm Hg.

Panda et al. (1992) evaluated the phytotoxicity of mercury from the solid waste deposits of a chloralkali plant. After exposure of barley to mercury waste for 7 days, seedling height was reduced by 19% at 64 ppm mercury in soil. Germination of barley was reduced by 20% at 103 ppm. The NOEC was 34.9 ppm. The authors did not apparently test the waste for contaminants other than mercury. However, mercury was the major contaminant in the waste, at 2550 mg/kg.

Confidence in the inorganic mercury benchmark of 0.3 ppm is low because it is based on a secondary reference, and the toxicity threshold in the chloralkali study was more than two orders of magnitude higher.

**Experiments conducted in solution.** The effect of Hg, as  $\text{HgCl}_2$ , on root elongation of 3-week-old Norway spruce seedlings grown for 7 days in nutrient solution (pH 4) was examined by Lamersdorf et al. (1991). The only concentration tested, 0.002 ppm Hg, reduced root elongation by 31%. Methyl mercury ( $\text{CH}_3\text{HgCl}$ ) completely stopped root elongation at a concentration of 0.0002 ppm, the only concentration tested.

Al-attar et al. (1988) investigated the effect of mercury (II) acetate on the length and weight of the roots and shoots of perennial ryegrass (*Lolium perenne*) seedlings in solution. This study was not used in the derivation of the benchmark for inorganic mercury or methyl mercury, but it is included here as an example of a toxicity test using mercury. A concentration of 0.0005 ppm of mercury (II) resulted in a 29% reduction in root dry weight, a 24% decrease shoot dry weight, and a 2% decrease in root length. The NOEC was 0.0001 ppm.

Schlegel et al. (1987) investigated the effects of inorganic ( $\text{HgCl}_2$ ) and organic ( $\text{CH}_3\text{HgCl}$ ) Hg on needle chlorophyll content, transpiration rate, and  $\text{CO}_2$  uptake of 2-week-old spruce seedlings in nutrient solution (pH 4.3) for 35 days. Methyl Hg at 0.002 ppm Hg (lowest concentration tested) reduced transpiration rate and  $\text{CO}_2$  uptake by 49, and 73%. At 0.02 ppm Hg (lowest concentration tested), both forms reduced needle chlorophyll content approximately 28%.

Godbold and Huttermann (1985) measured a 64% reduction in root elongation rate of 4-wk old Norway spruce seedlings grown in a nutrient solution (pH 4.3) containing 0.02 ppm Hg ( $\text{HgCl}_2$ ; lowest concentration tested) for 7 d.

Suszcynsky and Shann (1995) measured 50% reductions in root and shoot dry weights of 5-wk old tobacco (*Nicotiana glauca*) seedlings grown in nutrient solution (pH 6) containing 1 ppm Hg as  $\text{HgCl}_2$ . Mercury at 0.1 ppm had no effect.

Mhatre and Chaphekar (1982) tested several species at germination stage for their response to Hg. Seeds of sorghum, alfalfa, mung bean, cluster bean, and radish were allowed to germinate and grow for 5 days in solutions containing 0.001, 0.01, 0.1, or 1 ppm Hg as  $\text{HgCl}_2$ . At 0.01 ppm Hg, root length reductions ranged from 22 for radish to 52% for alfalfa, with Pennisetum, mustard, sorghum, and cluster bean having intermediate reductions. Shoot length of Pennisetum, alfalfa, and cluster bean were also reduced at this concentration 25, 37, and 26%, respectively. Root length of pea was reduced 40% by the addition of 0.1 ppm. Root and shoot lengths of mung bean were reduced 28 and 50% at this concentration.

Mukhiya et al. (1983) compared the toxicity of different Hg compounds to barley root and shoot length, and plant weight in solution at concentrations of 1, 5, 10, and 50 ppm and found organic forms to be more toxic than inorganic forms. After 7 days, mercury as  $\text{C}_8\text{H}_8\text{HgO}_2$  (phenyl mercuric acetate) at 5 ppm reduced shoot length and plant weight 27 and 25%. Mercuric acetate ( $\text{C}_4\text{H}_6\text{HgO}_4$ ) at 10 ppm Hg



reduced root length and plant weight 23%. Mercurous chloride ( $\text{Hg}_2\text{Cl}_2$ ) at 50 ppm reduced root length and plant weight 22 and 25%, and 50 ppm mercuric chloride ( $\text{HgCl}_2$ ) reduced plant weight 25%, root length 28%, and shoot length 35%.

The concentrations of Hg, from  $\text{HgCl}_2$ , required for a 50% reduction in seed germination and root length of mustard after 3 days of exposure in solution (pH 7.4), was reported by Fargasova (1994). LC50 for germination was 129 ppm and EC<sub>50</sub> for root length was 9.3 ppm Hg.

After 14 days, lengths of longest root and shoot of germinating rye grass seedlings were reduced 40 and 23% by 5 ppm Hg (lowest concentration tested) added to nutrient solution (pH 7) as  $\text{HgCl}_2$  (Wong and Bradshaw, 1982).

Confidence in the solution phytotoxicity benchmark for inorganic mercury (0.005 ppm) is moderate because it is based on 17 values and a range of plant species.

Confidence in the solution phytotoxicity benchmark for organic mercury (0.002 ppm Hg) is low because it is based on less than 10 values. Furthermore, the concentration approximating the 10th percentile was the lowest concentration tested and caused a 100% reduction in the growth parameter.

**Mechanism of Phytotoxicity.** Mercury and its compounds taken up by roots are translocated to only a limited extent in plants. Organic forms of Hg may be translocated to a greater degree than inorganic forms in some plants (Huckabee and Blaylock 1973). Gay (1975) reports that pea plants (*Pisum sativum*) form methyl mercury as an intermediate product from Hg added to the soil in organic and inorganic forms.

### 3.1.20 Molybdenum

**Experiments conducted in soil.** Kabata-Pendias and Pendias (1984) reported unspecified toxic effects on plants with the addition of 2 ppm Mo. Confidence in the benchmark of 2 ppm, based on this study alone, is low. Because the bioavailability of molybdenum increases with pH, toxicity would also likely increase with pH (unlike many metals). Neuman et al. (1987) assert that phytotoxicity of molybdenum in the field has never been recorded. Also, molybdenum is required by the nitrogenase of nitrogen-fixing microorganisms; thus, legumes are sometimes fertilized with the element. Although studies of molybdenum fertilization have been undertaken, none of these studies use molybdenum added to soil at a concentration of 2 ppm or above. Aghatise and Tayo (1994) observed slight increases in many growth parameters of soybean (*Glycine max*) with fertilization with molybdenum up to 0.8 kg/ha (about 0.5 mg/kg if 15 cm incorporation depth is assumed).

**Experiments conducted in solution.** Wallace et al. (1977b) evaluated the effect of Mo (as  $\text{H}_2\text{MoO}_4$ ) on root, leaf, and stem weights of bush bean seedlings in nutrient solution. After 14 days, leaf weight was reduced 36% by 9.6 ppm Mo, the lowest concentration tested.

Wallace (1979) measured a 35% decrease in leaf weight of bush bean when grown for 14 days in nutrient solution (pH 5) with 5.7 ppm Mo (as  $\text{H}_2\text{MoO}_4$ ), the lowest concentration tested. Root weight was not affected at this concentration.

Johnson (1966) reported unspecified toxic effects on plants grown in a solution with the addition of 0.5 ppm Mo.

Saco et al. (1995) observed in a fertilization experiment that 1 ppm of molybdenum did not affect nitrate and nitrite reductase activity, increased ammonium and nitrite content of the leaves and protein content in the root, and produced spots on the leaves. This quantity is not included in the derivation of the benchmark because other growth-related studies have been obtained.

Confidence in the 0.5 ppm benchmark for toxicity to plants growing in solution culture is low because it is based on fewer than 10 values.

**Mechanism of Phytotoxicity.** Molybdenum is required for symbiotic N<sub>2</sub> fixation by legumes and for growth of nonleguminous plants. The most important functions of Mo in plants are related to enzymes active in N metabolism (activation of nitrogenase and nitrate reductase). The majority of Mo taken up by the root system tends to remain in the roots although significant amounts may be translocated to the shoots in some cases (Wallace and Romney 1977b). Toxicity symptoms include chlorosis, apparently due to interference with Fe metabolism (Warrington 1954).

### 3.1.21 Nickel

**Experiments conducted in soil.** Dixon (1988) measured the response of red oak seedlings grown for 16 weeks in a sandy loam soil (pH 6, % organic matter 1.5) with addition of Ni (NiCl<sub>2</sub>). Nickel at 50 ppm reduced tree weight by 30%, while 20 ppm had no effect.

Khalid and Tinsley (1980) measured a 66% reduction in ryegrass shoot weight with the addition of 180 ppm Ni (as NiSO<sub>4</sub>) to a loam soil (pH 4.7). Addition of 90 ppm Ni had no effect. Plants were grown 4 weeks from seed.

Oats grown from seed for 110 days in the presence of 50 ppm Ni (as NiCl<sub>2</sub>) in soil (pH 6.1, CEC 6 meq/100 g, and % organic matter 1.4) had reductions of 38 and 63% in grain and straw weight (Halstead et al., 1969). In a second soil (pH 5.7, CEC 11.7 meq/100 g, % organic matter 4.1) only straw weight was reduced (45%) by addition of 100 ppm Ni (50 ppm had no effect).

Two cultivars of cotton were tested for tolerance to Ni in soil (Rehab and Wallace, 1978). Two-week-old seedlings grown for 35 days in soil (pH 6.8) to which 100 ppm Ni was added (lowest concentration tested) experienced reduced leaf and stem weights; 46 and 28% for the first cultivar, and 44 and 59% for the second.

Wallace et al. (1977d) report the results of experiments on the effects of Ni (as NiSO<sub>4</sub>) on seedlings of a variety of plants grown in a loam soil at several pHs. Corn grown in soil at pH 4.2, 5.6, and 7.5 experienced 74, 80, and 50% reductions in shoot weight after 14 days of growth with the addition of 250 ppm Ni. Ni at 100 ppm had no effect. At pH 5.8, bush beans grown for 16 days had a 64% reduction in shoot weight with the addition of 100 ppm (lowest concentration tested). At pH 7.5, a 36% reduction in plant weight occurred with 250 ppm Ni, while 100 ppm had no effect. After 28 days of growth in a loam soil at pH 5.8, bush bean leaf weight was reduced 45% by the addition of 100 ppm Ni, while 25 ppm had no effect. For barley under these same growth conditions, 25 ppm Ni (lowest concentration tested) reduced shoot weight 88%.

Traynor and Knezek (1973) measured a 21% reduction in corn plant weight with the addition of 294 ppm Ni (as NiCl<sub>2</sub>) to a sandy soil (pH 5, % organic matter 2, CEC 6 meq/100g soil) in which the plants had been grown for 5 weeks from seed. Addition of 220 ppm had no effect.

Confidence in the 30 ppm benchmark for Ni is low. Although there were 14 values, the concentration closest to the 10th percentile was the lowest concentration tested and caused an 88% reduction in the measured growth parameter. The next closest concentration was also responsible for a greater than 30% reduction in plant growth.

**Experiments conducted in solution.** The effect of Ni on stem diameter increase and plant weight of red pine, maple, dogwood, and honeysuckle was examined by Heale and Ormrod (1982). Seedlings (90-d from cutting) of red pine and honeysuckle grown for 110 days in nutrient solution containing 2 ppm Cu from NiSO<sub>4</sub> (lowest concentration tested) had reductions in stem diameter increase and plant weight of 100, and 25%, and 84 and 65%, respectively. Reductions in stem diameter increase in plant weight were 70% dogwood grown in solution containing 10 ppm Ni, while 2 ppm had no effect. Maple experienced a 48% decrease in plant weight only at 10 ppm Ni with the stem diameter increase remaining unaffected up to 20 ppm Ni (highest concentration tested).

Patterson and Olson (1983) evaluated the effect of several metals in solution (pH 5 to 6) on white spruce, black spruce, paper birch, jack pine, white pine, red pine, and honeysuckle grown from seed for 5 to 21 d. Paper birch, which was least tolerant of Co and Cu, was also most affected by Ni, with a 21% reduction in radicle elongation at 1 ppm Ni. As in the case of Co and Cu, white pine was the most tolerant plant tested with a 24% reduction in radicle elongation at 50 ppm Ni. The other species were intermediate in their response to Ni in solution.

Misra et al. (1994) evaluated the effects of several metals on seed germination and root growth of broad bean (*Vicia faba*). Seed germination was not affected by Ni (NiCl<sub>2</sub>) at concentrations up to 10 ppm Ni in nutrient solution. Root elongation in plants exposed for 3 days to 8 ppm Ni was reduced 30%, while 6 ppm had no effect.

Wang (1994) evaluated the effects of several metals and organic compounds on radicle weight of rice seed germinated and grown for 6 d in solution. The calculated EC<sub>50</sub> for Ni (NiCl<sub>2</sub>) was 0.85 ppm.

Wong and Bradshaw (1982) measured a 29% decrease in length of longest root of rye grass when germinated and grown for 14 days in nutrient solution (pH 7) with 0.13 ppm Ni [Ni(NH<sub>4</sub>)<sub>2</sub>(SO<sub>4</sub>)<sub>2</sub>], the lowest concentration tested. Length of the longest shoot was unaffected at this concentration.

The effects of Ni, from NiSO<sub>4</sub>, on germination and radicle length of radish, cabbage, turnip, lettuce, wheat, and millet after 3 days of growth in solution were determined by Carlson et al. (1991). There was no effect on seed germination up to 20 ppm Ni. Treatment levels were 0, 0.25, 0.5, 1, 2, 4, 8, 12, 16, and 20 ppm Ni. A concentration of 1 ppm reduced radicle length of lettuce and turnip by 24 and 25%. A concentration of 2 ppm reduced radicle length of cabbage by 30%. Eight ppm Ni reduced radicle length of radish by 32% and wheat by 45%, and 12 ppm caused a reduction in radicle length of millet by 40%.

Patel et al. (1976) found 26 and 27% decreases in leaf and stem weights of chrysanthemum seedlings when grown for 14 days in nutrient solution with 0.59 ppm Ni (NiSO<sub>4</sub>), while 0.006 ppm had no effect. Root weight was not affected at 0.59 ppm Ni.

Wallace (1979) measured 92 and 68% decreases in root and leaf weights of bush bean seedlings when grown for 21 days in nutrient solution (pH 5) with 1.2 ppm Ni, the only concentration tested.

The effect of Ni, as NiCl<sub>2</sub>, on plant weight of cotton grown in nutrient solution (pH 6) was evaluated by Rehab and Wallace (1978). Plant weight was reduced 92% by 5.9 ppm Ni, while 0.59 ppm had no effect.

Confidence in the 0.5 ppm phytotoxicity benchmark for Ni is high.

**Mechanism of Phytotoxicity.** Nickel is not generally considered to be an essential element for plants. However, it may be required by nodulated legumes for internal N transport as part of the urease enzyme (Aller et al. 1990). It is generally adsorbed as the Ni(II) ion and translocated in xylem and phloem with an organic chelate (Hutchinson 1981). Nickel is fairly uniformly distributed between roots and shoots (Wallace and Romney 1977b). Symptoms of Ni toxicity are generally Fe-deficiency induced chlorosis and foliar necrosis (Khalid and Tinsley 1980). Excess nickel affects nutrient absorption by roots, root development, and metabolism, and it inhibits photosynthesis and transpiration. Nickel can replace Co and other heavy metals located at active sites in metallo-enzymes and disrupt their functioning.

### 3.1.22 Selenium

**Experiments conducted in soil.** Wan et al. (1988) investigated the effects of Se(VI), as  $\text{Na}_2\text{SeO}_4$ , on alfalfa grown in three soils. In the sandy loam soil (pH 6.7, % organic matter 13) and in the two clay loam soils (pH 5.6, % organic matter 15; pH 6.9, % organic matter 13), shoot weight was reduced 83, 33, and 56% by the addition of 1.5 ppm Se(VI), while 0.5 ppm had no effect. Alfalfa was grown from seed to 0.25 bloom stage.

Singh and Singh (1978) investigated the toxicity of selenium (as  $\text{Na}_2\text{SeO}_3$ ) to wheat (*Triticum aestivum*) in a sandy soil from India. The authors observed a 23% decrease in biomass as 50 days and a 22% decrease in biomass and a 28% decrease in grain yield at maturity (135 days) associated with the addition of the lowest concentration of selenium tested, 2.5 ppm.

The effect of Se(VI) ( $\text{Na}_2\text{SeO}_4$ ) on alfalfa grown from seed to bloom was examined in five silty clay loam soils, ranging in pH from 6.9 to 7.8, by Soltanpour and Workman (1980). Shoot weight was reduced by 2 ppm in 4 of the 5 soils (91, 74, 23, and 27% reductions), with the greatest reductions in soils with the lowest organic matter content (% organic matter 3.1, 3.7, 5, and 6.5, respectively). Shoot weight was diminished 94% in the fifth soil (pH 7.0, % organic matter 6.3) with 4 ppm Se, 2 ppm having no effect.

Carlson et al. (1991) investigated the effects of Se(VI) (as  $\text{Na}_2\text{SeO}_4$ ) and Se (IV) (as  $\text{Na}_2\text{SeO}_3$ ) on sorgrass (*Sorghum vulgare*) grown from seed for 42 days in several soils. In a loamy sand soil (% organic matter 19, CEC 4 meq/100g soil) at pH 5.5 and 6.0, there were 59 and 53% reductions in shoot weight with the addition of 1 ppm Se(VI), (lowest concentration tested). No reductions were observed with additions of up to 4 ppm Se(IV). In a sandy soil (% organic matter 11, CEC 3 meq/100g soil) at pH 4.9, 1 ppm Se(VI) and 2 ppm Se(IV) caused 64 and 61% reductions in shoot weight. In this same sandy soil limed to pH 6.5, Se(IV) had no effect up to 4 ppm and Se(VI) reduced shoot weight 66% at 1 ppm.

Confidence in the 1 ppm benchmark for Se is low. Although there were 14 values, the concentration closest to the 10th percentile was the lowest concentration tested and consistently caused severe decreases in the measured growth parameter.

**Experiments conducted in solution.** Martin (1937a) evaluated the effect of Se (IV) from  $\text{Na}_2\text{SeO}_3$  on root and shoot weight, and plant height of wheat and buckwheat (*Fagopyrum esculentum* L.) seedlings growing in nutrient solution for 42 days. Selenium at 1 ppm (lowest concentration tested) reduced wheat root and shoot weight, and plant height 41, 40, and 23%. This concentration also reduced buckwheat root and shoot weight, and plant height 59, 75, and 44%.

Wu et al. (1988) investigated the toxicity of selenium as sodium selenate to five grass species in nutrient solution. After 3 weeks of exposure, reduction in root length and shoot height was observed for: bermudagrass (*Cynodon dactylon*) at 0.5 ppm, the lowest concentration tested; crested wheatgrass (*Agropyron desertorum*) at 1 ppm; seaside bentgrass (*Agrostis stolonifera*) at 1 ppm; and buffalograss (*Buchloe dactyloides*) at 2 ppm. The reductions in root and shoot growth at the LOEC were about 40% and 50% for bermudagrass, 30% and 30% for crested wheatgrass, 20% and 25% for seaside bentgrass, and 70% and 50% for buffalograss. Toxicity to tall fescue (*Festuca arundinaceae*) at concentrations of selenium up to 2 ppm was not observed.

Banuelos et al. (1997) studied the effect of selenium (added as  $\text{NaSeO}_4$ ) on nine “land races” of *Brassica juncea* (Indian mustard) and one race of *Brassica carinata* grown in solution culture. At 2 ppm of selenium, the lowest concentration tested, decreases in shoot yield of Indian mustard ranged from 12 to 23% and of root yield ranged from 11 to 34%. A 32% decrease in root yield of *Brassica carinata* was observed at the same 2 ppm concentration.

In experiments with plants found only in Se-rich soil, Trelease and Trelease (1938) found a 37% reduction in weight of milk-vetch (*Astragalus racemosus* L.) when grown in solution containing 27 ppm Se(IV) ( $\text{Na}_2\text{SeO}_3$ ), while 9 ppm had no effect on growth.

Wallace et al. (1980) examined the toxicity of selenate-Se ( $\text{Na}_2\text{SeO}_4$ ) on root and shoot weights of bush bean seedlings grown in nutrient solution (pH 4.4). Root weight was reduced 21% by 0.79 ppm Se, the lowest concentration tested, while shoot weight was unaffected.

Confidence in the 0.7 ppm phytotoxicity benchmark for Se is low because it is based on less than 10 values.

**Mechanism of phytotoxicity.** Selenium is not proven to be essential for plant growth. It is absorbed by plants as selenite, selenate or in organic form and the selenate may be the more toxic. It is believed that selenate is taken up actively while selenite uptake is largely passive (Peterson et al. 1981). Selenium is translocated to all parts of the plant, including the seed, in low molecular weight compounds (Broyer et al. 1972). Toxicity symptoms include chlorosis, stunting, and yellowing of the leaves. The mechanism of toxicity is thought to be indiscriminate replacement of S by Se in proteins and nucleic acids with disruptions in metabolism (Trelease et al. 1960). For example, selenomethionine is a less effective substrate than methionine for peptide bond formation, which could reduce protein synthesis (Eustice et al. 1981). Selenium-accumulating plant species incorporate less selenium into proteins than other species (Brown and Shrift 1981).

### 3.1.23 Silver

**Experiments conducted in soil.** There were no primary reference data showing toxicity of Ag to plants grown in soil. Confidence is low in the benchmark because it is based on a report of unspecified toxic effects on plants grown in a surface soil with the addition of 2 ppm Ag (Kabata-Pendias and Pendias 1984).

**Experiments conducted in solution.** Wallace (1979) examined the effect of Ag from  $\text{AgNO}_3$  on shoot weight of bush bean seedlings grown in nutrient solution (pH 5) for 13 days. Silver at 0.16 ppm reduced shoot weight 58% while 0.016 ppm had no effect.

Wang (1994) evaluated the effects of several metals and organic compounds on radicle weight of rice seed germinated and grown for 6 d in solution. The calculated  $\text{EC}_{50}$  for Ag ( $\text{AgNO}_3$ ) was 0.55 ppm.

Confidence in the 0.1 ppm benchmark for toxicity to plants growing in solution is low due to lack of data.

**Mechanism of Phytotoxicity.** Silver taken up by plants remains in the root system precipitated with phosphate or chloride (Ward et al., 1979). The toxicity of Ag is related to the binding potential of  $\text{Ag}^+$  ions to enzymes and other active molecules at cell surfaces (Cooper and Jolly 1970).

### 3.1.24 Technetium

**Experiments conducted in soil.** Wildung et al. (1977) investigated the effect of Tc on wheat and soybean grown in a silt loam soil (pH 6.8, % organic matter 1.4) from seed for 30 days. Addition of 1 ppm Tc as  $\text{TcO}_4^-$ , reduced shoot weight of wheat 100% and soybeans 99%, while 0.1 ppm had no effect.

Confidence in the benchmark of 0.2 ppm Tc is low because it is based on this study alone. The authors' chose to divide the LOEC by 5 because, although it was not expressed as such in the study, the severity of the effects seemed to border on mortality of the plants.

**Experiments conducted in solution.** Berlyn et al. (1980) conducted several experiments to examine the effect of Tc on fresh weight of soybean seedlings. When seedlings were germinated and allowed to grow for 20 days in nutrient solution containing 0.2 ppm Tc ( $\text{TcO}_4^-$ ), plant weight was reduced 31%. Technetium at 0.04 ppm had no effect. However, when seedlings were germinated and allowed to grow for 5 days before Tc was supplied, weight was reduced 36% at 20 ppm Tc, while 5 ppm had no effect.

Gast et al. (1978) examined the effect of Tc as pertechnetate ( $\text{NH}_4\text{TcO}_4$ ) on shoot and root weight of several plants grown from seed for 10 days in nutrient solution containing Tc. Technetium at 0.3 ppm reduced shoot weights of wheat and barley by 22 and 24%, while 0.03 ppm had no effect. A concentration of 1.2 ppm Tc caused decreases of 53% in root and shoot weights of oats, and a 24% reduction in shoot weight of radish, while 0.3 ppm had no effect. Corn shoot weight was reduced 31% by 5.8 ppm Tc, while 3 ppm had no effect. Soybean shoot weight was diminished 50% by 7.8 ppm Tc, while 5.8 ppm had no effect.

Confidence in the 0.2 ppm benchmark for toxicity to plants growing in solution is low because it is based on less than 10 values.

**Mechanism of phytotoxicity.** There are very little data on phytotoxicity of Tc. It is taken up and transported in plants as the pertechnetate ion ( $\text{TcO}_4^-$ ). The active uptake and toxicity of Tc may be due to its functioning as a nutrient analog, possibly P, S, or Mo (Wildung et al. 1979). The minimal amount of radiation measured in the experimental plants lead researchers to the conclusion that the effects were the result of the element rather than radiation (Wildung et al. 1977).

### 3.1.25 Tellurium

**Experiments conducted in soil.** No information was found on which to base a toxicity benchmark for plants growing in soil.

**Experiments conducted in solution.** Martin (1937b) evaluated the effect of Te (as  $\text{K}_2\text{TeO}_3$ ) on root and shoot weight, and plant height of wheat seedlings grown in nutrient solution containing Te for 42 days. Tellurium at 2 ppm (lowest concentration tested) reduced root and shoot weights 32 and 35%.

Confidence in the 2 ppm benchmark for toxicity to plants growing in solution is low due to lack of data.

**Mechanism of phytotoxicity.** Very little information on phytotoxicity of Te was found. The biological cycling of the element resembles that of Se although it is not accumulated in plant tissues in concentrations as high as Se (Kabata-Pendias and Pendias 1984). Although plant growth reductions have been measured in plants grown in solution culture to which Te has been added, no information on specific mechanisms of toxicity was found.

### 3.1.26 Thallium

**Experiments conducted in soil.** There are no primary reference data showing toxicity of Tl to plants grown in soil. Confidence in the benchmark is low because it based on a report of unspecified toxic effects on plants grown in a surface soil with the addition of 1 ppm Tl (Kabata-Pendias and Pendias 1984).

Al-attar et al. (1988) investigated the effect of thallium (as nitrate) on the length and weight of the roots and shoots of perennial ryegrass (*Lolium perenne*) seedlings in solution. A concentration of 0.5 ppm of thallium resulted in a 57% reduction in root dry weight, a 59% decrease shoot dry weight, a 26% decrease in root length, and a 54% decrease in shoot length. The NOEC was 0.1 ppm.

**Experiments conducted in solution.** The effect of Tl, as  $TlCl_3$ , on root elongation of 3-week old Norway spruce seedlings grown for 7 days in nutrient solution (pH 4) was examined by Lamersdorf et al. (1991). The only concentration tested, 0.02 ppm Tl, reduced root elongation by 27%.

The effects of Tl, from  $Tl_2SO_4$ , on germination and radicle length of radish, cabbage, turnip, lettuce, wheat, and millet after 3 days of growth in solution were determined by Carlson et al. (1991). There was no effect on seed germination up to 40 ppm Tl. Treatment levels were 0, 0.5, 1, 2.5, 5, 7.5, 10, 20, 30, and 40 ppm Tl. A concentration of 0.5 ppm reduced radicle length of lettuce by 65%. A concentration of 1 ppm reduced radicle length of turnip by 63%. Five ppm Tl reduced radicle length of radish by 22%, wheat by 30%, and millet by 35%. Radicle length of millet was reduced 23% by 7.5 ppm Tl.

Carlson et al. (1975) measured 40 and 55% reductions in photosynthesis when corn and sunflower (*Helianthus annuus* L.) seedlings were grown in nutrient solution containing 1 ppm Tl ( $TlCl_2$ ) (lowest concentration tested). Bowen (1979) reports undefined toxic effects on plant growth at this concentration also.

Confidence in the 0.02 ppm benchmark for toxicity to plants growing in solution is moderate.

**Mechanism of phytotoxicity.** Thallium is not essential for plant growth. When soluble forms are available, Tl is readily taken up by plants and translocated to aerial parts, probably because of its similarity to K. Toxic effects on plants include impairment of chlorophyll synthesis and seed germination, reduced transpiration due to interference in stomatal processes, growth reduction, stunting of roots, and leaf chlorosis (Adriano 1986).

### 3.1.27 Tin

**Experiments conducted in soil.** Romney et al. (1975) studied the effect of Sn (as  $SnCl_2$ ) on shoot weight of bush beans grown for 17 days in soil (pH 6). Shoot weight was reduced 22% by 500 ppm Sn, while 50 ppm had no effect.

Kabata-Pendias and Pendias (1984) reported unspecified toxic effects on plants grown in a surface soil with the addition of 50 ppm Sn. Confidence in the benchmark of 50 ppm for Sn is low.

**Experiments conducted in solution.** Romney et al. (1975) studied the effect of Sn (as  $\text{SnCl}_2$ ) on shoot weight of bush beans grown for 26 days in nutrient solution. A concentration of 119 ppm reduced shoot weight 81%, while 12 ppm had no effect. Confidence in the benchmark of 100 ppm for Sn in solution is low.

**Mechanism of phytotoxicity.** Tin is not essential to plants although it is readily taken up from nutrient solution. Most remains in the root system (Wallace and Romney 1977b). Tin is an element that is considered relatively innocuous but may be biomethylated to a more toxic form. Although plant growth reductions have been measured in plants grown in solution culture to which Sn has been added, no information on specific mechanisms of toxicity was found.

### 3.1.28 Titanium

**Experiments conducted in soil.** No information was found on which to base a toxicity benchmark for plants growing in soil.

**Experiments conducted in solution.** Wallace et al. (1977a) evaluated the effect of Ti ( $\text{TiCl}_3$ ) on root, stem, and leaf weight of bush beans grown in nutrient solution for 21 days. They measured a 23% decrease in leaf weight at 0.069 ppm Ti, the lowest concentration tested.

Hara et al. (1976) measured a 24% reduction in cabbage seedling weight after 55 days of growth in nutrient solution (pH 5) containing 4 ppm Ti ( $\text{TiCl}_3$ ). Titanium in solution at 0.4 ppm had no effect.

Confidence in the 0.06 ppm Ti in solution benchmark is low because of lack of data.

**Mechanism of phytotoxicity.** Titanium is not essential for plant growth and when taken up, it remains in the root system (Wallace and Romney 1977b). Toxicity symptoms include chlorosis, necrosis, and stunted growth. No information on specific mechanisms of toxicity was found.

### 3.1.29 Uranium

**Experiments conducted in soil.** Sheppard et al. (1983) grew Swiss chard in a sandy (pH 6.4, CEC 1.2 meq/100 g) and a peaty (pH 3, CEC 65 meq/100 g, % organic matter 92) soil to test the effects of  $^{238}\text{U}$  added as uranyl nitrate [ $\text{UO}_2(\text{NO}_3)_2$ ]. In the sandy soil, root weight was reduced 23% by 5 ppm U (lowest concentration tested), while shoot weight was not affected. In the peaty soil, root weight was reduced 44% by 10 ppm U (lowest concentration tested), while shoot weight was not affected. Confidence in the benchmark of 5 ppm U in soil is low because it is based on this study alone.

**Experiments conducted in solution.** Murthy et al. (1984) examined the effect of U, as  $\text{UO}_2$ , on germination and seedling length of soybean in nutrient solution for 6 days. A concentration of 42 ppm reduced seedling length 33%, while 0.42 ppm had no effect. Seed germination remained unaffected. Confidence in the benchmark of 40 ppm U in solution is low.

**Mechanism of phytotoxicity.** Uranium exists in the water-soluble fraction of plant tissue, probably as the uranyl ion and bound to cell wall proteins (Whitehead et al. 1971). The mechanisms of U phytotoxicity involve inhibition of enzyme systems and possibly binding to nucleic acids (Feldman et



al. 1967). The minimal amount of radiation measured in the experimental plants has led researchers to the conclusion that toxic effects are the result of the element rather than radiation (Sheppard et al. 1983).

### 3.1.30 Vanadium

**Experiments conducted in soil.** There are no primary reference data describing toxicity of V to plants grown in soil. Kabata-Pendias and Pendias (1984) report unspecified toxic effects on plants grown in a surface soil with the addition of 50 ppm V. Vanadium added at a concentration of 2.5 ppm was toxic to plants in a study reported by EPA (1980). Confidence in the 2 ppm benchmark for V is low.

**Experiments conducted in solution.** Wallace (1979) examined the effect of V from  $\text{NH}_4\text{VO}_3$  on root and shoot weight of bush bean seedlings grown in nutrient solution (pH 5) for 14 days. Vanadium at 0.51 ppm (lowest concentration tested) reduced root weight 46%. After 55 days, cabbage seedling plant weight was reduced 34% by 4 ppm V added as  $\text{VCl}_3$  to nutrient solution (pH 5), while 0.4 ppm had no effect on growth (Hara et al. 1976). Plant weight of soybean seedlings grown for 33 days in nutrient solution containing 6 ppm V (as  $\text{VOSO}_4$ ) was reduced 36%, while 3 ppm had no effect (Kaplan et al. 1990) on growth.

Nowakowski (1992) determined the effects of V ( $\text{NH}_4\text{VO}_3$ ) on root and shoot weights of three cultivars of peas when allowed to germinate and grow 14 days in solution containing V. Vanadium at 20 ppm reduced root and shoot weights of the cultivars approximately 40 and 25%.

The effects of V, from  $\text{VOSO}_4$ , on germination and radicle length after 3 days of growth in solution of radish, cabbage, turnip, lettuce, wheat, and millet were determined by Carlson et al. (1991). There was no effect on seed germination up to 40 ppm. Treatment levels were 0, 0.5, 1, 2.5, 5, 7.5, 10, 20, 30, and 40 ppm V for all but millet which was exposed additionally to 50, 60, 70, 80, and 100 ppm V. A concentration of 2.5 ppm reduced radicle length of lettuce by 30%, turnip by 50%, and cabbage by 42%. 10 ppm reduced radicle length of radish by 23%. Wheat was unaffected up to 40 ppm V. Radicle length of millet was reduced 50% by 60 ppm.

Gil et al. (1995) measured 26 and 28% reductions in root fresh weight and shoot dry weight of 2-wk old lettuce seedlings grown in nutrient solution (pH 4.7) containing 0.2 ppm V as  $\text{NH}_4\text{VO}_3$ . Vanadium at 0.1 ppm had no effect.

Confidence in the 0.2 ppm V in solution benchmark is low.

**Mechanism of phytotoxicity.** Vanadium is not known to be essential for plant growth although it may be involved in  $\text{N}_2$  fixation in nodules of legume roots. Toxicity symptoms include chlorosis, dwarfing, and inhibited root growth (Pratt 1966). Vanadium inhibits various enzyme systems while stimulating others, the overall effect on plant growth being negative (Peterson and Girling 1981). After uptake, most vanadium remains in the root system in insoluble form with Ca (Wallace and Romney 1977b).

### 3.1.31 Zinc

**Experiments conducted in soil.** In a pot culture starting with 2-year-old beech trees growing in an organic-rich forest soil (pH 4.8), Hagemeyer et al. (1993) measured a reduction of approximately 40% in annual ring growth in the presence of 3.3 ppm 1M ammonium acetate-extractable Zn when trees were grown for two seasons (lowest concentration tested). Zinc was added as  $\text{ZnSO}_4$ . The results of this

study are not directly comparable to others that report the amount of Zn added to the soil; however, the information is presented for reference in order to increase the number of plant types covered.

Muramoto et al. (1990) measured the effects of addition of Zn as ZnO to an alluvial soil (pH 6) on root and stem weights, stem length, and grain yield of wheat and rice grown from seed to maturity. Root weight of rice was reduced about 29% by 1000 ppm (lowest concentration tested). Wheat grain yield and plant weight were reduced 66 and 28% by 1000 ppm (lowest concentration tested).

The number of soybean seeds produced per plant was decreased by 28% when plants were grown in an average garden soil to which 25 ppm Zn was added as ZnSO<sub>4</sub> (Aery and Sakar 1991). Zn at 10 ppm had no effect. Nodule weight and number and seed weight were not affected by 25 ppm Zn. Plants were grown from seed to maturity.

White et al. (1979) evaluated the effect of Zn, as ZnSO<sub>4</sub>, on leaf and root weights of soybeans grown in a sandy loam soil at two pH levels. Leaf weight was reduced 30% by 131 ppm Zn at pH 5.5, while 115 ppm had no effect. At pH 6.5, leaf weight was reduced 33% by 393 ppm Zn.

Lata and Veer (1990) measured reductions in root and shoot lengths and weights of spinach and coriander (*Coriandrum sativum* L.) after 60 days in soil with added Zn form. Total soil Zn concentrations of 87 ppm reduced plant weight of spinach about 45%, and coriander about 22%.

Gall and Barnette (1940) investigated the effect of Zn, as ZnSO<sub>4</sub>, on corn and cowpeas (*Vigna sinensis* L.) grown in three soils for 30 days from seed. Results of this study are not directly comparable to most others because the authors report effective concentrations as "exchangeable", that is, Zn associated with the colloidal portion of the soil. Corn shoot weight was reduced 68% in a sandy soil at 404 ppm exchangeable Zn, while 202 ppm had no effect. In a sandy loam soil, the reduction was 38% at 334 ppm, while 222 ppm had no effect. In a clay loam soil, the reduction was 33% at 632 ppm, while 474 ppm had no effect. Cowpea shoot weight was reduced 29% in a sandy soil at 141 ppm exchangeable Zn, while 81 ppm had no effect. In a sandy loam soil, the reduction was 46% at 222 ppm, while 112 ppm had no effect. In a clay loam soil, the reduction was 28% at 316 ppm, while 158 ppm had no effect.

Confidence in the 50 ppm benchmark is moderate.

**Experiments conducted in solution.** Carroll and Loneragan (1968) measured effects of Zn on weight of 1-week old seedlings of barrel medic (*Medicago truncatula* L.), subterranean clover (*Trifolium subterraneum* L.), and lucerne (*Medicago sativa* L.) grown for 46 days in nutrient solution (pH 6). Zinc at 0.41 ppm reduced weight 80, 40, and 37%, respectively, while 0.08 ppm had no effect.

Wong and Bradshaw (1982) evaluated the effect of Zn on root and shoot length of rye grass grown in solution (pH 7) with Zn added as ZnSO<sub>4</sub>. After 14 days, they found a 63% reduction in the length of the longest root in response to 1.85 ppm (lowest concentration tested).

Patel et al. (1976) found a 30% decrease in root and stem weights of chrysanthemum seedlings when grown for 21 days in nutrient solution with 6.5 ppm Zn (as ZnSO<sub>4</sub>), while 0.65 ppm had no effect.

Wallace et al. (1977b) evaluated the effect of Zn (as ZnSO<sub>4</sub>) on leaf, stem, and root weights of bush bean seedlings in solution. After 16 days, weights were reduced 34, 41, and 44%, respectively, by 6.6 ppm Zn, while 0.66 ppm had no effect.

Misra et al. (1994) evaluated the effects of several metals on seed germination and root growth of broad bean (*Vicia faba*). Seed germination was not affected by Zn ( $\text{ZnCl}_2$ ) at concentrations up to 10 ppm Zn in nutrient solution. Root elongation of plants exposed for 3 days to 10 ppm Zn was reduced 30%, while 8 ppm had no effect.

Wang (1994) evaluated the effects of several metals and organic compounds on radicle weight of rice seed germinated and grown for 6 d in solution. The calculated  $\text{EC}_{50}$  for  $\text{ZnCl}_2$  was 26 ppm.

The benchmark of 0.4 ppm Zn is based on the work of Carroll and Loneragan (1986). Confidence in the benchmark is low because it is based on less than 10 values from experiments conducted with a limited range of plant species.

**Mechanism of phytotoxicity.** Zinc is an essential element for plant growth. It has a part in many enzymes and is involved in disease protection and metabolism of carbohydrates and proteins. Zinc is actively taken up by roots in ionic form and, to a lesser extent, in organically chelated form (Collins 1981). It is fairly uniformly distributed between roots and shoots being transported in the xylem in ionic form (Wallace and Romney 1977b). Transport in the phloem appears to be as an anionic complex (van Goor and Wiersma 1976). Toxicity symptoms include chlorosis and depressed plant growth (Chapman 1966). It acts to inhibit  $\text{CO}_2$  fixation, phloem transport of carbohydrates, and alter membrane permeability (Collins 1981).

## 3.2 ORGANIC COMPOUNDS

Numerous organic chemicals exist for which phytotoxicity has not been measured. Feng et al. (1997) and van Gestel et al. (1997) provide Quantitative Structure Activity Relationships (QSAR) for plant growth parameters as affected by substituted anilines and phenols, including chlorinated compounds.

### 3.2.1 Acenaphthene

Hulzebos et al. (1993) evaluated the effects of 75 organic compounds on growth of lettuce from seed for 14 d in two loam soils, and of 1-wk old lettuce seedlings in nutrient solution for 16 to 21 d. The difference in the loams was the clay content (12 and 24%). The calculated  $\text{EC}_{50}$  value for acenaphthene was 25 ppm in the soil containing 24% clay and  $>0.1$ ,  $<0.32$  ppm in solution.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.2 Aniline

Feng et al. (1996) calculated the  $\text{EC}_{50}$  for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The  $\text{EC}_{50}$  reported for aniline was 203.5 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.3 Biphenyl

Hulzebos et al. (1993) evaluated the effects of biphenyl on the growth of lettuce in solution and in a loam soil containing 12% clay, as described for acenaphthene. The calculated EC<sub>50</sub> value was 68 ppm for the soil and 2.1 ppm in solution.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.4 4-Bromoaniline

Feng et al. (1996) calculated the EC<sub>50</sub> for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC<sub>50</sub> reported for 4-bromoaniline was 37.8 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.5 3-Chloroaniline

Van Gestel et al. (1996) reported the toxicity of four chlorophenols and three chloroanilines on the growth of lettuce (*Lactuca sativa*) in a loam soil. The EC<sub>50</sub> for a 14-day test using 3-chloroaniline was 23 ppm. The NOEC was approximately a factor of 3.2 lower.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.6 4-Chloroaniline

Feng et al. (1996) calculated the EC<sub>50</sub> for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC<sub>50</sub> reported for 4-chloroaniline was 39.4 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.7 2-Chlorophenol

Feng et al. (1996) calculated the EC<sub>50</sub> for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC<sub>50</sub> reported for 2-chlorophenol was 58.3 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.8 3-Chlorophenol

Van Gestel et al. (1996) reported the toxicity of four chlorophenols and three chloroanilines on the growth of lettuce (*Lactuca sativa*) in a loam soil. The EC<sub>50</sub> for a 14-day test using 3-chlorophenol was 7 ppm. The NOEC was approximately a factor of 3.2 lower.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.9 4-Chlorophenol

Feng et al. (1996) calculated the EC50 for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC50 reported for 4-chlorophenol was 47.4 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.10 2-Cresol

Feng et al. (1996) calculated the EC50 for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC50 reported for 2-cresol was 54.9 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.11 Di-n-butyl phthalate

Overcash et al. (1982) evaluated the phytotoxicity of di-n-butyl phthalate and toluene on plant growth in two soils. Fescue, corn, and soybeans were grown from seed for 21 days in a clay soil (pH 5, % organic matter 1.6, CEC 8.7 meq/100g soil) and a sandy loam soil (% organic matter approximately 1, CEC approximately 3 meq/100g soil). Both soils were tested at pH 4 and 6. The treatment levels for di-n-butyl phthalate were 200, 2000, or 20000 ppm. In the clay soil, no effect was seen on seed germination at the highest concentration. Corn fresh weight was reduced 23% by 200 ppm. Fescue fresh weight was reduced 73% by 2000 ppm. In the sandy loam soil at pH 4, soybean seed germination was reduced 56% by 200 ppm. Corn fresh weight was reduced 34% by 200 ppm. In the sandy loam soil at pH 6, no effect on seed germination was noted. Fresh weights of corn and soybean were reduced 44 and 29% by 200 ppm. Fescue fresh weight was reduced 56% by 2000 ppm. Confidence in the benchmark of 200 ppm is low.

**Mechanism of toxicity.** Di-n-butyl phthalate has a low vapor pressure and is nonionic. It is biologically and chemically decomposed in soil. Di-n-butyl phthalate may be produced in plants (some phthalate esters are known to be), and it is metabolically degraded by plants and animals (Overcash et al. 1982).

### 3.2.12 3,4-Dichloroaniline

Feng et al. (1996) calculated the EC50 for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC50 reported for 3,4-dichloroaniline was 14.1 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.13 2,4-Dichlorophenol

Feng et al. (1996) calculated the EC50 for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC50 reported for 2,4-dichlorophenol was 17.1 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.14 3,4-Dichlorophenol

Van Gestel et al. (1996) reported the toxicity of four chlorophenols and three chloroanilines on the growth of lettuce (*Lactuca sativa*) in a loam soil. The EC<sub>50</sub> for a 7-day test using 3,4-dichlorophenol was 25 ppm. The NOEC was approximately a factor of 3.2 lower.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.15 Diethylphthalate

Hulzebos et al. (1993) evaluated the effects of diethylphthalate on growth of lettuce in solution in a loam soil containing 12% clay, as described for acenaphthene. The calculated EC<sub>50</sub> value was 134 ppm for the soil and 25 ppm in solution.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.16 2,4-Dinitrophenol

Overcash et al. (1982) evaluated the phytotoxicity of 2,4-dinitrophenol on plant growth in two soils as described for di-n-butyl phthalate. Treatment levels were 10, 20, 40, 80, and 100 ppm. In the clay soil, no effect on seed germination was noted. Soybean fresh weight was reduced 63% by 20 ppm. Corn and fescue fresh weights were reduced 35 and 80% by 40 ppm. In the sandy loam soil at pH 4, soybean seed germination was reduced 30%, and fresh weight 65%, by 40 ppm. Corn seed germination was reduced 42% by 80 ppm, while fresh weight was reduced 25% by 20 ppm. Fescue fresh weights were reduced 29% by 40 ppm. In the sandy loam soil at pH 6, no effect on seed germination was noted. Fresh weight of soybean was reduced 23% by 20 ppm, of corn 25% by 40 ppm, and of fescue 24% by 80 ppm. Confidence in the benchmark of 20 ppm is moderate.

**Mechanisms of toxicity.** 2,4-dinitrophenol is more toxic to plants at low pH, where the weak acid is largely in the molecular, undissociated form which is more easily taken up by, and active in, plants than the dissociated anion. Primary modes of action on plants are increasing respiration, uncoupling of oxidative phosphorylation, and activation of ATP-ase. It is relatively persistent in soils, especially at low pH. The pH range of 4 to 6 included in the studies of Overcash et al. (1982) was not great enough to show differences in toxicity due to soil adsorption and differential ionic activity.

### 3.2.17 Furan

Hulzebos et al. (1993) evaluated the effects of furan on growth of lettuce in two soils and solution, as described for acenaphthene. The calculated EC<sub>50</sub> values for the soil containing 24% clay was >1,000 ppm, and for the soil containing 12% clay it was 617 ppm. The EC<sub>50</sub> values in solution were 130 and 135 ppm.

### 3.2.18 Heptane

Hulzebos et al. (1993) evaluated the effects of heptane on growth of lettuce in two soils and solution, as described for acenaphthene. The calculated EC<sub>50</sub> values for both soils was >1000 ppm. The EC<sub>50</sub> values in solution were 1.7 and 47 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.19 Hexachlorocyclopentadiene

Hulzebos et al. (1993) evaluated the effects of hexachlorocyclopentadiene on growth of lettuce in solution and in a loam soil containing 24% clay, as described for acenaphthene. The calculated EC<sub>50</sub> value was 10 ppm for the soil and 0.1 ppm in solution.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.20 Naphthalene

Hulzebos et al. (1993) evaluated the effects of naphthalene on growth of lettuce in solution and in loam soil containing 12% clay, as described for acenaphthene. The calculated EC<sub>50</sub> value was >100 ppm for the soil and 13 ppm in solution.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.21 3-Nitroaniline

Feng et al. (1996) calculated the EC<sub>50</sub> for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC<sub>50</sub> reported for 3-nitroaniline was 69.2 ppm.

### 3.2.22 4-Nitroaniline

Feng et al. (1996) calculated the EC<sub>50</sub> for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC<sub>50</sub> reported for 4-nitroaniline was 43.6 ppm.

### 3.2.23 Nitrobenzene

McFarlane et al. (1990) examined the effect of nitrobenzene on soybean, barley, lettuce, Russian olive (*Elaeagnus angustifolia* L.), autumn olive, green ash (*Fraxinus pennsylvanica*), hybrid poplar (*Populus x robusta*), and honeysuckle grown in nutrient solution. One-year-old autumn olive seedlings exposed for 2 days to 8 ppm nitrobenzene (only concentration tested) experienced reductions of 95 and 90% in photosynthesis and transpiration. Confidence in the solution benchmark is low because it is based on this study alone.

**Mechanism of toxicity.** No information was found on phytotoxicity of nitrobenzene except for the studies showing reduced photosynthesis and transpiration of autumn olive discussed above (McFarlane et al. 1990).

### 3.2.24 4-Nitrophenol

Feng et al. (1996) calculated the EC<sub>50</sub> for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the

log of chemical concentration in solution was plotted. The EC<sub>50</sub> reported for 4-nitrophenol was 12.3 ppm.

### 3.2.25 Pentachlorophenol

Hulzebos et al. (1993) evaluated the effects of pentachlorophenol on growth of lettuce in two soils and solution, as described for acenaphthene. Soils with a higher clay content had a higher EC<sub>50</sub> value for phenol (8 and 3.2 ppm); the EC<sub>50</sub> value of 0.03 ppm in the solution was lower than in either soil.

Gunther and Pestemer (1990) reported the toxic levels of pentachlorophenol causing reduced fresh weight of shoots of oats (*Avena sativa*) after 14 days of exposure and turnips (*Brassica rapa*) after 10 days of exposure in a sandy loam soil. The EC<sub>50</sub>s for oats and turnips were 20 and 10 ppm, respectively.

Confidence in the benchmarks for soil and solution, 3 and 0.3 ppm, is low.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.26 Phenol

Hulzebos et al. (1993) evaluated the effects of phenol on growth of lettuce in solution and in two soils, as described for acenaphthene. Soils with a higher clay content had a lower EC<sub>50</sub> value for phenol (79 and 168 ppm); the EC<sub>50</sub> values in solution were lower (14 and 20 ppm) than in either soil.

Feng et al. (1996) calculated the EC<sub>50</sub> for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC<sub>50</sub> reported for phenol was 125.6 ppm.

Confidence in the benchmarks for soil and solution, 70 and 10 ppm, is low.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.27 Polychlorinated biphenyls (PCBs)

**Aroclor 1254.** Strek and Weber (1980) investigated the effects of the PCB Aroclor 1254 on fescue, sorghum (*Sorghum bicolor* L.), corn, soybean, and beets grown in a sandy soil (pH 4.7, % organic matter 1, CEC 1.5 meq/100g soil) from seed for 16 days. Height, water use, and top fresh weight of corn, sorghum, and fescue were unaffected by the 1000 ppm test concentration. Fresh top weight of three soybean varieties was reduced an average of 28% and water use 43%. Beet height and fresh top weight were reduced 100% and water use 94%. Fresh foliage weight of pigweed (*Amaranthus retroflexus* L.) was assessed in soil containing up to 100 ppm Aroclor 1254. The more sensitive variety had a 22% reduction in weight at 40 ppm, while 20 ppm had no effect.

Strek and Weber (1982b) also evaluated the effect of Aroclor 1254 on pigweed grown in the sandy soil used by Strek and Weber in the 1980 work. They found a 23% reduction in the height of plants grown from seed for 28 days in soil containing 100 ppm. A treatment level of 50 ppm had no effect.

Weber and Mrozek (1979) evaluated the effect of Aroclor 1254 on soybean grown in the sandy soil used by Strek and Weber in the 1980 work. They found a 27% reduction in the fresh shoot weight of plants grown from seed for 26 days in soil containing 100 ppm. A treatment level of 10 ppm had no effect. There was also a 45% reduction in water use at the 100 ppm level.



Confidence in the benchmark of 40 ppm for PCBs is low because it is based on fewer than 10 values.

**Mechanism of toxicity.** Commercial formulations of PCBs are various, usually unquantified, mixes of polychlorinated biphenyls. Although plant growth reductions resulting from PCB addition to soil have been measured, no mechanism of toxicity was suggested. Because cumulative water use seems to be more sensitive to PCBs than plant growth (Weber and Mrozek 1979), it has been suggested that effects on plants may be indirect, following an effect on transpiration (Strek and Weber 1982a). *In vitro* cultures of plant cells are capable of metabolizing and detoxifying PCBs (Fletcher et al. 1987).

### 3.2.28 Styrene

Hulzebos et al. (1993) evaluated the effects of styrene on growth of lettuce in solution and in a loam soil containing 24% clay, as described for acenaphthene. The calculated EC<sub>50</sub> value was 320 ppm for the soil and 18 ppm in solution.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.29 2,3,5,6-Tetrachloroaniline

Van Gestel et al. (1996) reported the toxicity of four chlorophenols and three chloroanilines on the growth of lettuce (*Latuca sativa*) in a loam soil. The EC<sub>50</sub> for a 14-day test using 2,3,5,6-tetrachloroaniline was 17 ppm. The NOEC was approximately a factor of 3.2 lower.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.30 Tetrachloroethene

Hulzebos et al (1993) evaluated the effects of tetrachloroethene on growth of lettuce in solution and in a loam soil containing 24% clay, as described for acenaphthene. The calculated EC<sub>50</sub> value was >1,000 ppm for the soil and 12 ppm in solution.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.31 Toluene

Overcash et al. (1982) evaluated the phytotoxicity of toluene on plant growth in two soils as described for di-n-butyl phthalate. In the clay soil, no effect on seed germination was noted. Corn fresh weight was reduced 30% by 200 ppm. Soybean fresh weight was reduced 32% by 20,000 ppm. In the sandy loam soil at pH 4, soybean seed germination was reduced 50% by 2000 ppm. Corn seed germination was reduced 86% by 20,000 ppm. In the sandy loam soil at pH 6, no effect on seed germination was noted. Fresh weight of soybean was reduced 40% by 200 ppm, and of corn and fescue, 68 and 22% by 20,000 ppm. Confidence in the benchmark of 200 ppm toluene is low.

Hulzebos et al. (1993) evaluated the effects of toluene on growth of lettuce in solution and in a loam soil containing 12% clay, as described for phenol. The calculated EC<sub>50</sub> value was >1,000 ppm for the soil and 16 ppm in solution.

**Mechanism of toxicity.** Toluene is a lipophilic compound that is more toxic in vapor form because of its ability to dissolve lipids of cuticle and plasma membranes. It is not actively taken up by plants

from soils but may adsorb to root surfaces and enter by dissolving membrane components. Toluene is known to be oxidatively detoxified by plants.

Toluene has been found to negatively affect seed germination and plant weight. Toxic effects appear to be acute because toluene is not accumulated in plants. In the case of seeds, it is thought that high levels of toluene may kill the embryo (Overcash et al. 1982).

### 3.2.32 4-Toluidine

Feng et al. (1996) calculated the EC<sub>50</sub> for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC<sub>50</sub> reported for 4-toluidine was 102.2 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.33 2,4,5-Trichloroaniline

Van Gestel et al. (1996) reported the toxicity of four chlorophenols and three chloroanilines on the growth of lettuce (*Latuca sativa*) in a loam soil. The EC<sub>50</sub> for a 14-day test using 2,4,5-trichloroaniline was 23 ppm. The NOEC was approximately a factor of 3.2 lower.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.34 Trichloroethane

Hulzebos et al. (1993) evaluated the effects of trichloroethane on growth of lettuce in solution and in a loam soil containing 24% clay, as described for acenaphthene. The calculated EC<sub>50</sub> value was >1000 ppm for the soil and 104 ppm in solution.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.35 2,4,5-Trichlorophenol

Van Gestel et al. (1996) reported the toxicity of four chlorophenols and three chloroanilines on the growth of lettuce (*Latuca sativa*) in a loam soil. The EC<sub>50</sub> for a 14-day test using 2,4,5-trichlorophenol was 4.3 ppm. The NOEC was approximately a factor of 3.2 lower.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.36 2,4,6-Trichlorophenol

Feng et al. (1996) calculated the EC<sub>50</sub> for the effect of several anilines and phenols on the root length of 5-day-old Chinese cabbage plants in solution. The inhibition rate in root elongation versus the log of chemical concentration in solution was plotted. The EC<sub>50</sub> reported for 2,4,6-trichlorophenol was 12.7 ppm.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.37 Ortho-xylene

Hulzebos et al. (1993) evaluated the effects of ortho-xylene on growth of lettuce in solution and in loam soil containing 24% clay as described for phenol. The calculated EC<sub>50</sub> value was >1,000 ppm for the soil and >1, <3.2 ppm in solution.

**Mechanism of toxicity.** No information was available on the mechanism of toxicity.

### 3.2.38 Xylene

Allen et al. (1961) evaluated the effect of xylene in insecticides on emergence of sugar beet seedlings exposed in solution (pH 6) for 2 days. Root length was reduced 32% by 100 ppm xylene, the lowest concentration tested. Confidence in the solution benchmark is low because it is based on this study alone.

**Mechanism of toxicity.** There was no information found on phytotoxicity of xylene except for the study showing reduced beet root growth (Allen et al.1961).

## **4. RELATIONSHIP BETWEEN SOIL PHYTOTOXICITY BENCHMARKS AND OTHER ECOTOXICOLOGICAL CRITERIA**

### **4.1 COMPARISON OF PHYTOTOXICITY BENCHMARKS FOR CONTAMINANTS IN SOIL TO CANADIAN ENVIRONMENTAL QUALITY CRITERIA FOR CONTAMINATED SITES**

The Canadian Council of Ministers of the Environment has developed Environmental Quality Criteria for contaminated sites. These are “numerical limits for contaminants in soil and water intended to maintain, improve, or protect environmental quality and human health at contaminated sites in general” (CCME 1991). Remediation criteria are presented for comparison to the phytotoxicity benchmarks because they represent levels considered generally protective of human health and the environment for specified uses of soil (in this case the most conservative use, agriculture, has been chosen) without taking into account site-specific conditions. If contaminant concentrations exceed the remediation criteria for a current or future land use, further investigation or remediation is needed. These criteria have an interim status and their derivation is in the process of refinement. They have been adopted from several Canadian jurisdictions and many lack supporting rationale (CCME, 1991). The remediation criteria are not strictly comparable to our phytotoxicity benchmarks because they also take into account human health and, presumably, soil organisms and the entire food chain dependent upon the soil. New CCME Soil Quality Guidelines are being developed and will be made available in late 1995. The CCME remediation criteria and the soil and solution benchmarks are listed in Table 2.

Contaminant phytotoxicity benchmarks derived by our method are more conservative than those of the CCME except for Be, Cd, F, Sn, Tl, 2,4-dinitrophenol, PCP, and styrene. These differences may be due to the Canadian consideration of a larger number of endpoints or a different level of protection. There is no indication in the source publication as to the level of protection being afforded by the CCME Remediation Criteria; however, if human health is considered in the conservative agriculture land use scenario, one would expect it to be high. This is seen in the case of 2,4-dinitrophenol which has a high mammalian toxicity.

### **4.2 COMPARISON OF PHYTOTOXICITY BENCHMARKS FOR CONTAMINANTS IN SOIL TO RIVM (NETHERLANDS) ECOTOXICOLOGICAL INTERVENTION VALUES FOR CONTAMINANTS IN SOILS**

The National Institute of Public Health and Environmental Protection developed Ecotoxicological Intervention Values which represent concentrations of contaminants in soil causing 50% of the species potentially present in an ecosystem to experience adverse effects (van den Berg et al. 1993). They take into account plants, soil fauna, and microorganisms. The method for deriving the values (the RAB method) is described by Denneman and van Gestel in several RIVM publications in Dutch. In order to take the influence of soil characteristics on the bioavailability of compounds, data were corrected for organic matter and clay content as described by van den Berg et al. (1993). Risks resulting from biomagnification were included. The RIVM values and the soil and solution benchmarks are listed in Table 2.

**Table 2. Comparison of screening benchmark concentrations for the phytotoxicity of chemicals in soil to CCME remediation criteria (RC), RIVM ecotoxicological intervention values (EIVs), arithmetic means of elements in uncontaminated soils of the Oak Ridge Reservation (ORR), and geometric means of elements in soils and surficial material of the eastern U.S.**

Chemical	Screening benchmark (mg/kg)	CCME RC <sup>a</sup> (mg/kg)	RIVM EIVs (mg/kg)	ORR (mg/kg)	USGS eastern U.S. (mg/kg)
Aluminum	50	—	—	15700	33000
Antimony	5	20	—	0.46	0.52
Arsenic	10	20	40	9.7	4.8
Barium	500	750	625	87.9	290
Beryllium	10	4	—	0.77	0.55
Boron	0.5	2 <sup>b</sup>	—	10.4	—
Bromine	10	—	—	—	0.62
Cadmium	4	3	12	0.22	—
Chromium (total)	1	750	230	24	33
Chromium (VI)	—	8	—	—	—
Cobalt	20	40	240	15.6	5.9
Copper	100	150	190	11.2	13
Fluorine	200	200	—	—	130
Iodine	4	—	—	—	0.68
Lead	50	375	290	26.8	14
Lithium	2	—	—	9.4	17
Manganese	500	—	—	1318	260
Mercury	0.3 <sup>c</sup>	0.8 <sup>d</sup>	10 <sup>d</sup>	0.20 <sup>d</sup>	0.08 <sup>d</sup>
Molybdenum	2	5	<480	3.9	0.32

Table 2 (continued)

Chemical	Screening benchmark (mg/kg)	CCME RC <sup>a</sup> (mg/kg)	RIVM EIVs (mg/kg)	ORR (mg/kg)	USGS eastern U.S. (mg/kg)
Nickel	30	150	210	15.1	11
Selenium	1	2	—	0.73	0.3
Silver	2	20	—	1.22	—
Technetium	0.2	—	—	—	—
Thallium	1	1	—	0.50	—
Tin	50	2	—	—	0.86
Uranium	5	—	—	—	2.1
Vanadium	2	200	—	32.3	43
Zinc	50	600	720	46.2	40
3-Chloroaniline	20				
2-Chlorophenol	—	0.05	10		
3-Chlorophenol	7	0.05	10		
4-Chlorophenol	—	0.05	10		
2-Cresol	—	0.1	50		
2,4-Dichlorophenol	—	0.05	10		
3,4-Dichlorophenol	20	0.05	10		
2,4 Dinitrophenol	20	0.1 <sup>e</sup>	—	—	—
Di-n-butyl phthalate	200	—	—	—	—
4-nitrophenol	—	0.1			

**Table 2 (continued)**

<b>Chemical</b>	<b>Screening benchmark (mg/kg)</b>	<b>CCME RC<sup>a</sup> (mg/kg)</b>	<b>RIVM EIVs (mg/kg)</b>	<b>ORR (mg/kg)</b>	<b>USGS eastern U.S. (mg/kg)</b>
Pentachlorophenol	3	0.05 <sup>f</sup>	5	—	—
PCBs	40	0.5	70	—	—
Styrene	300	0.10	—	—	—
2,3,5,6-tetrachloroaniline	20				
Toluene	200	0.1	130	—	—
2,4,5-Trichloroaniline	20				
2,4,5-Trichlorophenol	4	0.05	10		
2,4,6-Trichlorophenol	—	0.05	10		

<sup>a</sup>Agricultural land-use context.

<sup>b</sup>Hot water soluble B.

<sup>c</sup>Inorganic Hg.

<sup>d</sup>Does not indicate form (organic or inorganic).

<sup>e</sup>Each nonspecified non-chlorinated phenolic compound is not to exceed 0.1 ppm.

<sup>f</sup>Each nonspecific chlorinated phenolic compound is not to exceed 0.05 ppm.

## **5. COMPARISON OF PHYTOTOXICITY BENCHMARKS FOR CONTAMINANTS IN SOIL TO CONCENTRATIONS OF CHEMICALS IN UNPOLLUTED SOILS**

### **5.1 COMPARISON TO USGS ELEMENT CONCENTRATIONS IN SOILS AND OTHER SURFICIAL MATERIALS OF THE EASTERN UNITED STATES**

To place the three sets of critical values into a broader perspective, soil chemical concentrations are presented as reported by the U.S. Geological Survey (USGS) in a survey of soils of the eastern United States (Shacklette and Boerngen 1984) (Table 2). These samples were collected and analyzed by the USGS to represent, as far as possible, soils that were very little altered from their natural condition and that supported native plants. The values are presented as "total" concentrations.

It is interesting to compare the levels of elements cited in the literature as toxic against concentrations of those same elements found in natural (i.e., not directly contaminated) soils. This comparison is reasonable in most cases because benchmarks were generally based on nominal soil concentrations (i.e., those added to the soil by the experimenter) as opposed to a measure of either total concentration or of the plant-available quantity of the element in the soil. Seldom was the background level of the "contaminant" element in the soil measured, the assumption being that there is very little of the element existing naturally in the soil compared to treatment levels added. This is often, but not always, a reasonable assumption. The USGS compilation contains concentrations of elements mainly derived from strong acid extractions, although, in the case of uranium, neutron activation was used to measure a true total concentration. Soils of the eastern United States were chosen for comparison because most of the experimental results used to develop the benchmarks were derived from agricultural soils of the eastern United States. Surficial deposits of the western United States, especially arid and mountainous regions, may contain unusually high concentrations of naturally-occurring trace elements.

For several of the metals, the phytotoxicity benchmark was below the geometric mean for the element in soils and surficial deposits in the eastern United States. Comparing the benchmarks to the acid-extractable element data, a large discrepancy is realized between the USGS soil Al value and the low soil benchmark based on a quantity of Al added to soil. Al is present in most soils in exchangeable and amorphous forms that are not readily available to plants. The acid extraction removes for measurement all exchangeable and some portion of the amorphous Al. In the case of Cr, Li, and V, the form of the element added or some other aspect of the experimental design may account for the low benchmark concentration as compared to mean levels in soils.

### **5.2 COMPARISON TO DOE OAK RIDGE RESERVATION BACKGROUND SOIL CHARACTERIZATION ELEMENT CONCENTRATIONS IN SOILS**

The Background Soil Characterization Project at the Oak Ridge Reservation was established to determine the background concentrations of organics, metals, and radionuclides in natural soils that are important to environmental restoration projects (Watkins et al. 1993). Soils were sampled, field classified, and analyzed for chemicals using several methods. The data presented in Table 2 are arithmetic means of 46 sampling sites of elements extracted using nitric acid and hydrogen peroxide (EPA 1986). This standard EPA acid digestion for sediments, sludges, and soils is not explicitly meant to extract total elements from a sample. A comparison with total soil concentrations of elements measured by neutron activation analysis shows that for many elements (Sb, As, Cr, Co, Mn, Si, V, Zn)



the acids used do extract most of the element in question (Watkins et al. 1993). Unfortunately, not all elements are amenable to measurement by neutron activation analysis.

As with the USGS data, there is a large discrepancy between the Background Soil Characterization Project soil Al value and the soil phytotoxicity benchmark based on a quantity of Al added to soil. The high manganese levels of geologic origin at the Oak Ridge Reservation emphasize the need for local reference soils for comparison to waste site soils. In the case of Cr, Li, and V, the form of the element added or some other aspect of the experimental design may account for the low benchmark concentrations as compared to levels found in Oak Ridge Reservation soils.

## 6. RECOMMENDATIONS AND CONCLUSIONS

The values presented in Table 1 are intended for contaminant screening in the hazard identification (problem formulation) phase of ecological risk assessments. Chemicals with soil concentrations that exceed both the phytotoxicity benchmark for soil and the background soil concentration for the soil type, and which may be derived from waste disposal, are contaminants of potential concern. Background soil concentrations have been derived for the Oak Ridge Reservation and should be generated for other Comprehensive Environmental Response, Compensation and Liability Act (CERCLA) sites as well. Similarly, soil solution or shallow groundwater concentrations that exceed both the phytotoxicity benchmark for solutions and the background water concentration for the aquifer, which may be derived from waste disposal, and to which plant roots may be exposed are contaminants of potential concern.

For baseline ecological risk assessments, and other assessments that may lead to regulatory actions, assessors should consult the primary sources of toxicity data and then determine the applicability of the data to their specific site. In addition, assessments should not blindly rely on laboratory toxicity data. Where phytotoxicity is suspected, phytotoxicity tests should be performed with the contaminated soil. In addition, the site should be surveyed for signs of phytotoxicity such as inexplicable bare areas, low plant diversity, low plant vigor, or symptoms of toxic injury.

## 7. REFERENCES

- Adema D. M. M. and L. Henzen. 1989. A comparison of plant toxicities of some industrial chemicals in soil culture and soilless culture. *Ecotoxicol. Environ. Saf.* 18:219-29.
- Adriano, D. C. 1986. *Trace Elements in the Terrestrial Environment*. Springer-Verlag, New York.
- Aery, N. C. and S. Sakar. 1991. Studies on the effect of heavy metal stress on growth parameters of soybean. *J. Environ. Biol.* 12(1):15-24.
- Aghatise, V. O., and T. O. Tayo. 1994. Response of soybean (*Glycine max*) to molybdenum application in Nigeria. *Indian J. of Agric. Sci.* 64:597-603.
- Al-Attar, A. F., M. H. Martin, and G. Nickless. 1988. Uptake and toxicity of cadmium, mercury and thallium to *Lolium perenne* seedlings. *Chemosphere* 17:1219-1225.
- Aldrich, D. G., A. P. Vanselow, and G. R. Bradford. 1951. Lithium toxicity in citrus. *Soil Sci.* 71:291-95.
- Allen, W. R., W. L. Askew, and K. Schreiber. 1961. Effect of insecticide fertilizer mixtures and seed treatments on emergence of sugar beet seedlings. *J. Econ. Entom.* 54(1):181-187.
- Aller, A. J., J. L. Bernal, M. J. del Nozal, and L. Deban. 1990. Effects of selected trace elements on plant growth. *J. Sci. Food Agric.* 51:447-479.
- Asher, C. J., and P. F. Reay. 1979. Arsenic uptake by barley seedlings. *Aust. J. Plant Physiol.* 6:459-466.
- Banuelos, G. S., H. A. Ajwa, L. Wu, X. Guo, S. Akohoue, and S. Zambrzuski. 1997. Selenium-induced growth reduction in *Brassica* land races considered for phytoremediation. *Ecotoxicology and Environ. Safety* 36:282-287.
- Bartlett, R. J., and B. James. 1979. Behavior of chromium in soils: III. Oxidation. *J. Environ. Qual.* 8:31-35.
- Berlyn, G. P., S. S. Dhillon, and E. E. Koslow. 1980. Technetium: A toxic waste product of the nuclear fuel cycle: Effects on soybean growth and development. *Environ. Management* 4(2):149-156.
- Bingham, F. T., A. L. Page, R. J. Mahler, and T. J. Ganje. 1975. Growth and cadmium accumulation of plants grown on a soil with a cadmium-enriched sewage sludge. *J. Environ. Qual.* 4(2):207-211.
- Bowen, H. J. M. 1979. *Environmental Chemistry of the Elements*. Academic Press, NY.
- Breeze, V. G. 1973. Land reclamation and river pollution problems in the Croal Valley caused by waste from chromate manufacture. *J. Appl. Ecol.* 10:513-525.
- Brewer, R. 1966. Fluorine. In *Diagnostic Criteria for Plants and Soils*. H.D. Chapman (ed). Univ. of California, Div. Agric. Sci., Riverside. pp. 180-196.

- Brown, T. A., and A. Shrift. 1981. Exclusion of selenium from proteins of selenium-tolerant *Astragalus* species. *Plant Physiol.* 67:1051-1053.
- Broyer, T. C., C. M. Johnson, and R. P. Huston. 1972. Selenium and nutrition of *Astragalus*. I. Effects of selenite or selenate supply on growth and selenium content. *Plant Soil* 36:635-649.
- Burke, D. G., K. Watkins, and B. J. Scott. 1990. Manganese toxicity effects on visible symptoms, yield, manganese levels, and organic acid levels in tolerant and sensitive wheat cultivars. *Crop Sci.* 30:275-80.
- Burton, K. W., E. Morgan, and A. Roig. 1984. The influence of heavy metals upon the growth of sitka-spruce in South Wales forests. II. Greenhouse experiments. *Plant Soil* 78:271-82.
- Canadian Council of Ministers of the Environment (CCME). 1991. *Interim Canadian Environmental Quality Criteria for Contaminated Sites*. CCME EPC-CS34. Winnipeg, Manitoba.
- Carlson, C. L., D. C. Adriano, and P. M. Dixon. 1991. Effects of soil-applied selenium on the growth and selenium content of forage species. *J. Environ. Qual.* 20:363-68.
- Carlson, R. W. and F. A. Bazzaz. 1977. Growth reduction in American Sycamore (*Plantanus occidentalis* L.) caused by Pb-Cd interaction. *Environ. Pollut.* 12:243-53.
- Carlson, R. W. and G. L. Rolfe. 1979. Growth of rye grass and fescue as affected by lead-cadmium-fertilizer interaction. *J. Environ. Qual.* 8(3):348-352.
- Carlson, R. W., F. A. Bazzaz, and G. L. Rolfe. 1975. The effects of heavy metals on plants. II. Net photosynthesis and transpiration of whole corn and sunflower plants treated with Pb, Cd, Ni, and Tl. *Environ. Res.* 10:113-120.
- Carlson, C. L., D. C. Adriano, K. S. Sajwan, S. L. Abels, D. P. Thoma, and J. T. Driver. 1991. Effects of selected trace metals on germinating seeds of six plant species. *Water Air Soil Pollut.* 59:231-240.
- Carroll, M. D., and J. F. Loneragan. 1968. Response of plant species to concentrations of zinc in solution. *Aust. J. Agric. Res.* 19:859-68.
- Chapman, H. D. 1966. Zinc. In *Diagnostic Criteria for Plants and Soils*. H. D. Chapman (ed). Univ. of California, Div. Agric. Sci. pp. 484-499.
- Chaudhry, F. M., A. Wallace, and R. T. Mueller. 1977. Barium toxicity in plants. *Commun. Soil Sci. Plant Anal.* 8(9):795-97.
- Chou, S. F. J., and R. A. Griffin. 1986. Solubility and soil mobility of polychlorinated biphenyls. In J. S. Waid (ed) *PCBs and the Environment, Vol. 1*. pp. 101-119.
- Collins, J. C. 1981. Zinc. In *Effects of Heavy Metal Pollution on Plants, Vol. 1. Effects of Trace Metals on Plant Function*. N. W. Lepp (ed) Applied Science Publishers, New Jersey. pp. 145-170.
- Cooper. C. F., and W. C. Jolly. 1970. Ecological effects of silver iodide and other weather modification agents: A review. *Water Resour. Res.* 6:88-98.

- Cunningham, L. M. 1977. Physiological and biochemical aspects of cadmium in soybean: The effects of induced Cd toxicity on the uptake and translocation of Zn, Fe, Mg, Ca and K. *Proc. Annual Conf. on Trace Substances in the Environment*, pp. 133-45.
- Cunningham, L. M., F. W. Collins, and T. C. Hutchinson. 1975. Physiological and biochemical aspects of cadmium toxicity in soybean. I. Toxicity symptoms and autoradiographic distribution of Cd in roots, stems and leaves. In *Proceedings - International Conference on Heavy Metals in the Environment*. pp. 97-120.
- Deuel, L. E. and A. R. Swoboda. 1972. Arsenic toxicity to cotton and soybeans. *J. Environ. Qual.* 1:317-20.
- Dixon, R. K. 1988. Response of ectomycorrhizal *Quercus rubra* to soil cadmium, nickel and lead. *Soil Biol. Biochem.* 20(4):555-59.
- El-Enany, A. E. 1995. Alleviation of cadmium toxicity on maize seedlings by calcium. *Biologia Plantarum* 37(1): 93-99.
- EPA (U.S. Environmental Protection Agency). 1980. *A Screening Procedure for the Impacts of Air Pollution Sources on Plants, Soils, and Animals*. EPA 450/2-81-078. Washington, D.C.
- EPA. 1986. *Test Methods for Evaluating Solid Waste. Vol. 1: Laboratory Manual Physical/Chemical Methods*. SW846. Office of Solid Waste and Emergency Response. Washington, D.C.
- Eustice, D. C., F. J. Kull, and A. Shrift. 1981. Selenium toxicity: aminoacylation and peptide bond formation with selenomethionine. *Plant Physiol.* 67:1054-1058.
- Fargasova, A. 1994. Effect of Pb, Cd, Hg, As, and Cr on germination and root growth of *Sinapis alba* seeds. *Bull. Environ. Contam. Toxicol.* 52:452-456.
- Feldman, I., J. Jones, and R. Cross. 1967. Chelation of uranyl ions adenine nucleotides. *J. Am. Chem. Soc.* 89:49-55.
- Feng, L., L. Wang, Y. Zhao, and B. Song. 1996. Effects of substituted anilines and phenols on root elongation of cabbage seed. *Chemosphere* 32:1575-1583.
- Fernandes, J. C., and F. S. Henriques. 1991. Biochemical, physiological, and structural effects of excess copper in plants. *Bot. Rev.* 57(3):246-272.
- Fletcher J., A. Groeger, J. McCrady, and J. McFarlane. 1987. Polychlorobiphenyl (PCB) metabolism by plant cells. *Biotech. Letters* 9(11):817-820.
- Foy, C. D., R. L. Chaney, and M. C. White. 1978. The physiology of metal toxicity in plants. *Ann. Review Plant Physiol.* 29:511-566.
- Foy, C. D., R. R. Weil, and C. A. Coradetti. 1995. Differential manganese tolerances of cotton genotypes in nutrient solution. *J. Plant Nutr.* 18(4): 685-706.
- Gall, O. E. and R. M. Barnette. 1940. Toxic limits of replaceable zinc to corn and cowpeas grown of three Florida soils. *J. Am. Soc. Agron.* 32:23-32.

- Garate, A., I Ramos, M. Manzanares, and J. J. Lucena. 1993. Cadmium uptake and distribution in three cultivars of *Latuca* sp. *Bull. Environ. Contam. Toxicol.* 50:709-716.
- Gast, R. G., L. H. Thorvig, E. R. Landa, and K. J. Gallagher. 1978. Technetium-99 toxicity to plants and sorption by soils. In D.C. Adriano and I.L. Brisbin, Jr. (eds) *Environmental Chemistry and Cycling Processes*. Tech Info Ctr, Us DOE., pp. 550-567.
- Gay, D. D. 1975. Biotransformation and chemical form of mercury in plants. In *International Conference on Heavy Metals in the Environment Symposium Proceedings, Vol. II*, pt. 1. pp. 87-95.
- Gil, J., C. E. Alvarez, M. C. Martinez, and N. Perez. 1995. Effect of vanadium on lettuce growth, cationic nutrition, and yield. *J. Environ. Sci. Health A30* (1):73-87.
- Godbold, D. L., and A. Huttermann. 1985. Effect of zinc, cadmium, and mercury on root elongation of *Picea abies* (Karst.) Seedlings and the significance of these metals to forest die-back. *Environ. Pollut.* A38:375-381.
- Godbold, D. L., and C. Kettner. 1991. Use of root elongation studies to determine aluminum and lead toxicity in *Picea abies* seedlings. *J. Plant Physiol.* 138:231-235.
- Görransson, A. and T. D. Eldhuset. 1991. Effects of aluminum on growth and nutrient uptake of small *Picea abies* and *Pinus sylvestris* plants. *Trees* 5:136-42.
- Greger, M., E. Brammer, S. Lindberg, G. Larsson, and J. Idestam-Almquist. 1991. Uptake and physiological effects of cadmium in sugar beet (*Beta vulgaris*) related to mineral provision. *J. Exp. Botany* 42:729-737.
- Gunther, P., and W. Pestemer. 1990. Risk assessment for selected xenobiotics by bioassay method with higher plants. *Environ. Manage.* 14:381-388.
- Gupta, U. C. 1984. Boron nutrition of alfalfa, red clover, and timothy grown on podzol soils of eastern Canada. *Soil Sci.* 137(1):16-22.
- Gupta, D. B. and S. Mukherji. 1977. Effects of toxic concentrations of copper on growth and metabolism of rice seedlings. *Z. Pflanzenphysiol. Bd.* 82:95-106.
- Gussarsson, M. 1994. Cadmium-induced alterations in nutrient composition and growth of *Betula pendula* seedlings: the significance of fine roots as a primary target for cadmium toxicity. *J. Plant Nutr.* 17:2151-2163.
- Hagemeyer, J., D. Lohrmann, and S.-W. Breckle. 1993. Development of annual xylem rings and shoot growth of young beech (*Fagus sylvatica* L.) grown in soil with various Cd and Zn levels. *Water, Air Soil Pollut.* 69:351-361.
- Haghiri, F. 1973. Cadmium uptake by plants. *J. Environ. Qual.* 2(1):93-95.
- Halstead, R. L., B. J. Finn, and A. J. MacLean. 1969. Extractability of nickel added to soils and its concentration in plants. *Can. J. Soil Sci.* 49:335-342.

- Hara, T., Y. Sonoda, and I. Iwai. 1976. Growth response of cabbage plants to transition elements under water culture conditions. *Soil Sci. Plant Nutr.* 22(3):307-315.
- Hassett, J. J., J. E. Miller, and D. E. Koeppe. 1976. Interaction of lead and cadmium on maize root growth and uptake of lead and cadmium by roots. *Environ. Pollut.* 11:297-302.
- Heale, E. L., and D. P. Ormrod. 1982. Effects of nickel and copper on *Acer rubrum*, *Cornus stolonifera*, *Lonicera tatarica*, and *Pinus resinosa*. *Can. J. Bot.* 60:2674-2681.
- Hooper, M. C. 1937. An investigation of the effect of lead on plants. *Ann. Appl. Biol.* 24:690-695.
- Huckabee, J. W., and B. G. Blaylock. 1973. Transfer of mercury and cadmium from terrestrial to aquatic ecosystems. *Adv. Exp. Med. Biol.* 40:125-160.
- Hulzebos, E. M., D. M. M. Adema, E. M. Dirven-van Breemen, L. Henzen, W. A. van Dis, H. A. Herbold, J. A. Hoekstra, R. Baerselman, and C. A. M. van Gestel. 1993. Phytotoxicity studies with *Latuca sativa* in soil and nutrient solution. *Environ. Toxicol. Chem.* 12:1079-1094.
- Hutchinson, T. C. 1981. Nickel. In: *Effects of Heavy Metal Pollution on Plants, Vol. 1. Effects of Trace Metals on Plant Function*. N.W. Lepp (ed) Applied Science Publishers, New Jersey. pp. 171-212.
- Inouhe, M., S. Ninomiya, H. Tohyama, M. Joho, and T. Murayama. 1994. Different characteristics of roots in the cadmium-tolerance and Cd-binding complex formation between mono- and dicotyledonous plants. *J. Plant Res.* 107:201-207.
- Iwai, I., T. Hara, and Y. Sonoda. 1975. Factors affecting cadmium uptake by the corn plant. *Soil Sci. Plant Nutr.* 21:37-46.
- Jalil, A., F. Selles, and J. M. Clarke. 1994. Growth and cadmium accumulation in two durum wheat cultivars. *Comm. Soil Sci. Plant Anal.* 25 (15&16):2597-2611.
- Jarvis, S. C., and D. C. Whitehead. 1983. The absorption, distribution and concentration of copper in white clover grown on a range of soils. *Plant Soil* 75:427-434.
- Jenne, E. A., and S. N. Luoma. 1977. Forms of trace elements in soils, sediments, and associated waters: An overview of their determination and biological activity. In H. Drucker and R.E. Wildung (eds) *Biological Implications of Metals in the Environment*. CONF-750929. Tech. Info. Ctr., ERDA. Washington. pp. 110-143.
- Jiang, Q. Q., and B. R. Singh. 1994. Effect of different sources and forms of arsenic on crop yield and arsenic concentration. *Water Air Soil Pollut.* 74:321-343.
- John, M. K. 1973. Cadmium uptake by eight food crops as influenced by various soil levels of cadmium. *Environ. Pollut.* 4:7-15.
- John, M. K. and C. Van Laerhoven. 1972. Lead uptake by lettuce and oats as affected by lime, nitrogen, and sources of lead. *J. Environ. Qual.* 1(2):169-171.
- John, M. K., H. H. Chuah, and C. Van Laerhoven. 1972a. Cadmium contamination of soil and its uptake by oats. *J. Environ. Qual.* 6(6):555-557.

- John, M. K. , C. Van Laerhoven, and H.H. Chuah. 1972b. Factors affecting plant uptake and phytotoxicity of cadmium added to soils. *Environ. Sci. Technol.* 6(12):1005-1009.
- John, M. K., H. H. Chuah, and C. J. Van Laerhoven. 1977. Boron response and toxicity as affected by soil properties and rates of boron. *Soil Sci.* 124:34-39.
- Johnson, C. M. 1966. Molybdenum. In *Diagnostic Criteria for Plants and Soils*. Chapman, H. D. (ed.). Univ. of California, Div. Agric. Sci., Riverside.
- Kabata-Pendias, A., and H. Pendias. 1984. *Trace Elements in Soils and Plants*. CRC Press, Inc. Boca Raton, Florida.
- Kaplan, D. I., D. C. Adriano, C.L. Carlson, and K.S. Sajwan. 1990. Vanadium: Toxicity and accumulation by beans. *Water Air Soil Pollut.* 49:81-91.
- Keltjens, W. G. 1990. Effects of aluminum on growth and nutrient status of Douglas-fir seedlings grown in culture solution. *Tree Physiol.* 6:165-75.
- Khalid, B. Y. and J. Tinsley. 1980. Some effects of nickel toxicity on rye grass. *Plant Soil.* 55:139-44.
- Khan, D. H. and B. Frankland. 1983. Effects of cadmium and lead on radish plants with particular reference to movement of metals through soil profile and plant. *Plant Soil.* 70:335-345.
- Khan, D. H. and B. Frankland. 1984. Cellulolytic activity and root biomass production in some metal-contaminated soils. *Environ. Pollut.* 33:63-74.
- Koeppel, D. E. 1981. Lead: Understanding the minimal toxicity of lead in plants. In *Effects of Heavy Metal Pollution on Plants, Vol. 1. Effects of Trace Metals on Plant Function*. N. W. Lepp (ed) Applied Science Publishers, New Jersey pp. 55-76.
- Kummerova, M., and R. Brandejsova. 1994. Project TOCOEN. The fate of selected pollutants in the environment. Part XIX. The phytotoxicity of organic and inorganic pollutants--cadmium. The effect of cadmium on the growth of germinating maize plants. *Toxicological and Environ. Chem.* 42:115-132.
- Labanauskas, C. K. 1966. Manganese. In *Diagnostic Criteria for Plants and Soils*. H. D. Chapman (ed). Univ. of California, Div. Agric. Sci., Riverside. pp. 264-285.
- Lamersdorf, N. P., D. L. Godbold, and D. Knoche. 1991. Risk assessment of some heavy metals for the growth of Norway spruce. *Water Air Soil Pollut.* 57-58:535-543.
- Langheinrich, U., R. Tischner, and D. L. Godbold. 1992. Influence of a high Mn supply on Norway spruce [*Picea abies* (L.) Karst.] seedlings in relation to the nitrogen source. *Tree Physiol.* 10:259-71.
- Lata, K. and B. Veer. 1990. Phytotoxicity of Zn amended soil to *Spinacia* and *Coriandrum*. *Acta Bot. Indica.* 18:194-198.
- Le Bot, J., E. A. Kirkby, and M. L. van Beusichem. 1990. Manganese toxicity in tomato plants: Effects on cation uptake and distribution. *J. Plant Nutr.* 13(5):513-25.



- Lepp, N. W. 1981. Copper. In *Effects of Heavy Metal Pollution on Plants, Vol. 1. Effects of Trace Metals on Plant Function*. N. W. Lepp (ed) Applied Science Publishers, New Jersey. pp. 111-144.
- Lewis, J. C. and W. L. Powers. 1941. Antagonistic action of chlorides on the toxicity of iodides to corn. *Plant Physiol.* 393-98.
- Lin, Z. and D. L. Myhre. 1991. Differential response of citrus rootstocks to aluminum levels in nutrient solutions: I. Plant growth. *J. Plant Nutr.* 14(11):1223-38.
- Lindsay, W. L. 1979. *Chemical Equilibria in Soils*. John Wiley and Sons, New York.
- Liu, D., W. Jiang, W. Wang, F. Zhao, and C. Lu. 1994. Effects of lead on root growth, cell division, and nucleolus of *Allium cepa*. *Environ. Pollut.* 86:1-4.
- Llugany, M., C. Poschenrieder, and J. Barcelo. 1995. Monitoring of aluminum-induced inhibition of root elongation in four maize cultivars differing in tolerance to aluminum and proton toxicity. *Physiologia Plantarum* 93:265-271.
- Long, R. E. and L. G. Morgan. 1990. *The potential for biological effects of sediment-sorbed contaminants tested in the national status and trends program*. NOAA Technical Memorandum NOS OMA 52.
- Mackay, A. D., J. R. Caradus, and M. W. Pritchard. 1990. Variation for aluminum tolerance in white clover. *Plant Soil* 123:101-105.
- Macleod, L. B. and L. P. Jackson. 1967. Aluminum tolerance of two barley varieties in nutrient solution, peat, and soil culture. *Agronomy J.* 59:359-63.
- Marsh, K. B., and L. A. Peterson. 1990. Gradients in Mn accumulation and changes in plant form for potato plants affected by Mn toxicity. *Plant Soil* 121:157-163.
- Martin, A. L. 1937a. Toxicity of selenium to plants and animals. *Am. J. Bot.* 23:471-483.
- Martin, A. L. 1937b. A comparison of the effects of tellurium and selenium on plants and animals. *Am. J. Bot.* 24:198-203.
- Martin, J. P. 1966a. Bromine. In *Diagnostic Criteria for Plants and Soils*. H.D. Chapman (ed). Univ. of California, Div. Agric. Sci., Riverside. pp. 62-64.
- Martin, J. P. 1966b. Iodine. In *Diagnostic Criteria for Plants and Soils*. Chapman, H.D. (ed.). Univ. of California, Div. Agric. Sci., Riverside. pp. 200-202.
- Martin, J. P., G. K. Helmkamp, and J. O. Ervin. 1956. Effect of bromide from a soil fumigant and from  $\text{CaBr}_2$  on the growth and chemical composition of citrus plants. *Soil Sci. Am. Proc.* 20:209-212.
- McFarlane, C. M., T. Pfleeger, and J. Fletcher. 1990. Effect, uptake and disposition of nitrobenzene in several terrestrial plants. *Environ. Toxicol. Chem.* 9:513-520.

- McGrath, S. P. 1982. The uptake and translocation of tri- and hexa-valent chromium and effects on the growth of oat in flowing nutrient solution. *New Phytol.* 92:381-390.
- McLean, F. T. and B. E. Gilbert. 1927. The relative aluminum tolerance of crop plants. *Soil Sci.* 24:163-74.
- Mhatre, G. N., and S. B. Chaphekar. 1982. Effect of heavy metals on seed germination and early growth. *J. Environ. Biol.* 3(2):53-63
- Miles, L. J. and G. R. Parker. 1979a. The effect of soil-added cadmium on several plant species. *J. Environ. Qual.* 8(2):229-232.
- Miles, L. J. and G. R. Parker. 1979b. Heavy metal interaction for *Andropogon scoparius* and *Rudbeckia hirta* grown on soil from urban and rural sites with heavy metals additions. *J. Environ. Qual.* 8(4):443-49.
- Miles, C. D., J. R. Brandle, D. J. Daniel, O. Chu-Der, P. D. Schnore, and D.J. Uhlik. 1972. Inhibition of photosystem II in isolated chloroplasts by lead. *Plant Physiol.* 49:820-825.
- Miller, J.E., J.J. Hassett, and D.E. Koeppe. 1976. Uptake of cadmium by soybeans as innced by soil cation exchange capacity, pH, and available phosphorus. *J. Environ. Qual.* 5(2):157-160.
- Miller, J. E., J. J. Hassett, and D. E. Koeppe. 1977. Interactions of lead and cadmium on metal uptake and growth of corn plants. *J. Environ. Qual.* 6(1):18-20.
- Misra, J., V. Pandey, and N. Singh. 1994. Effects of some heavy metals on root growth of germinating seeds of *Vicia faba*. *J. Environ. Sci. Health A29* (10):2229-2234.
- Mocquot, B., J. Vangronsveld, H. Clijsters, and M. Mench. 1996. Copper toxicity in young maize (*Zea mays* L.) plants: effects on growth, mineral and chlorophyll contents, and enzyme activities. *Plant and Soil* 182:287-300.
- Moral, R., J. N. Pedreno, I. Gomez, and J. Mataix. Effects of chromium on the nutrient element content and morphology of tomato. *J. Plant Nutr.* 18(4):815-822.
- Mukherji, S., and B. Das Gupta. 1972. Characterization of copper toxicity in lettuce seedlings. *Physiol. Plant.* 27:126-129.
- Mukhiya, Y. K., K. C. Gupta, N. Shrotriya, J. K. Joshi, and V. P. Singh. 1983. Comparative responses of the action of different mercury compounds on barley. *Intern. J. Environ. Studies* 20:323-327.
- Muramoto, S., H. Nishizaki, and I. Aoyama. 1990. The critical levels and the maximum metal uptake for wheat and rice plants when applying metal oxides to soil. *J. Environ. Sci. Health, Part B* 25(2):273-80.
- Murthy, T. C. S., P. Weinberger, and M. P. Measures. 1984. Uranium effects on the growth of soybean (*Glycine max* (L.) Merr.). *Bull. Environ. Contam. Toxicol.* 32:580-586.
- Neary, D. G., G. Schneider, and D. P. White. 1975. Boron toxicity in red pine following municipal waste water irrigation. *Soil Sci. Soc. Am. Proc.* 39:981-982.

- Nelson, D. W., and L. E. Sommers. 1982. Total carbon, organic carbon, and organic matter. In *Methods of Soil Analysis, Part 2*. ASA/SSSA. Madison, WI. pp. 574-576.
- Neuman, E. R., J. L. Schrack, and L. P. Gough. 1987. Copper and molybdenum. pp. 215-232. In R. D. Williams and G. E. Schuman (eds.), *Reclaiming Mine Soils and Overburden in the Western United States: Analytical Parameters and Procedures*. Soil Conservation Society of America, Ankeny, Iowa.
- Newton, H. P. and S. J. Toth. 1952. Response of crop plants to I and Br. *Soil Sci.* 73:127-34.
- Nichol, B. E., and L. A. Oliveira. 1995. Effects of aluminum on the growth and distribution of calcium in roots of an aluminum-sensitive cultivar of barley (*Hordeum vulgare*). *Can. J. Bot.* 73:1849-1858.
- Nowakowski, W. 1992. Vanadium bioaccumulation in *Pisum sativum* seedlings. *Biol. Plant.* 35(3):461-465.
- Ouzounidou, G., M. Moustakas, E. P. Eleftheriou. 1997. Physiological and ultrastructural effects of cadmium on wheat (*Triticum aestivum*) leaves. *Arch. Environ. Contam. Toxicol.* 32:154-160.
- Overcash, R. M., J. B. Weber, and M. L. Miles. 1982. *Behavior of organic priority pollutants in the terrestrial system: Di-n-butyl phthalate ester, toluene, and 2,4 dinitrophenol*. UNC-WRRI-82-171. Water Resources Research Institute, Univ. North Carolina.
- Page, A. L., F. T. Bingham, and C. Nelson. 1972. Cadmium absorption and growth of various plant species as influenced by solution cadmium concentration. *J. Environ. Qual.* 1(3):288-91.
- Panda, K. K., M. Lenka, and B. B. Panda. 1992. Monitoring and assessment of mercury pollution in the vicinity of a chlorakali plant. II. Plant-availability, tissue-concentration and genotoxicity of mercury from agricultural soil contaminated with solid waste assessed in barley (*Hordeum vulgare* L.). *Environ. Pollut.* 76:33-42.
- Patel, P. M., and A. Wallace, and R. T. Mueller. 1976. Some effects of copper, cobalt, cadmium, zinc, nickel, and chromium on growth and mineral element concentration in chrysanthemum. *J. Am. Soc. Hortic. Sci.* 101(5):553-556.
- Patterson, W. A., and J. J. Olson. 1983. Effects of heavy metals on radicle growth of selected woody species germinated on filter paper, mineral and organic soil substrates. *Can. J. For. Res.* 13:233-238.
- Peterson, P. J. 1983. Adaptation to toxic metals. In *Metals and Micronutrients: Uptake and Utilization by Plants*. D. A. Robb and W. S. Pierpoint (eds), Academic Press, New York. pp. 51-69.
- Peterson, P. J., L. M. Benson, and R. Zieve. 1981. Metalloids. In *Effects of Heavy Metal Pollution on Plants, Vol. 1. Effects of Trace Metals on Plant Function*. N. W. Lepp (ed), Applied Science Publishers, New Jersey, pp. 279-342.
- Peterson, P. J., and C. A. Girling. 1981. Other trace metals. In *Effect of Heavy Metal Pollution on Plants, Vol 1. Effects of Trace Metals on Plant Function*. N. W. Lepp (ed), Applied Science Publishers, New Jersey. pp. 213-278.

- Pintro, J., J. Barloy, and P. Fallavier. 1996. Aluminum effects on the growth and mineral composition of corn plants cultivated in nutrient solution at low aluminum activity. *J. Plant Nutr.* 19:729-741.
- Pratt, P. F. 1966. Vanadium. In *Diagnostic Criteria for Plants and Soils*. Chapman, H.D. (ed.). Univ. of California, Div. Agric. Sci., Riverside. pp. 480-483.
- Rascio, N., F. D. Vecchia, M. Ferretti, L. Merlo, and R. Ghisi. 1993. Some effects of cadmium on maize plants. *Arch. Environ. Contam. Toxicol.* 25:244-249.
- Reber, H. H. 1989. Threshold levels of cadmium for soil respiration and growth of spring wheat (*Triticum aestivum* L.), and difficulties with their determination. *Biol. Fertil. Soils* 7:152-157.
- Rehab, F. I., and A. Wallace. 1978. Excess trace metal effects on cotton: 6. Nickel and cadmium in Yolo loam soil. *Commun. Soil Sci. Plant Anal.* 9(8):779-784.
- Rehab, F. I., and A. Wallace. 1978. Excess trace metal effects on cotton: 5. Nickel and cadmium in solution culture. *Commun. Soil Sci. Plant Anal.* 9(8):771-778.
- Rolfe, G. L. and F. A. Bazzaz. 1975. Effect of lead contamination on transpiration and photosynthesis of loblolly pine and autumn olive. *Forest Sci.* 21(1):33-35.
- Romney, E. M. and J. D. Childress. 1965. Effects of beryllium in plants and soil. *Soil Sci.* 100(2):210-17.
- Romney, E. M., J. D. A. Wallace, and G. V. Alexander. 1975. Response of bush bean and barley to tin applied to soil and to solution culture. *Plant Soil* 42:585-589.
- Romney, E. M., J. D. Childress, and G. V. Alexander. 1962. Beryllium and the growth of bush beans. *Science* 185:786-87.
- Rosehart, R. G., and J. Y. Lee. 1973. The effect of arsenic trioxide on the growth of white spruce seedlings. *Water Air Soil Pollut.* 2:439-443.
- Saco, D., S. Martin, and M. Alvarez. 1995. Nitrogen metabolism in *Nicotiana rustica* L. grown with molybdenum. I. Vegetative development. *Commun. Soil Sci. Plant Anal.* 26:1719-1732.
- Sadana, U. S. and B. Singh. 1987a. Yield and uptake of cadmium, lead and zinc by wheat grown in soil polluted with heavy metals. *J. Plant Sci. Res.* 3:11-17.
- Sadana, U. S. and B. Singh. 1987b. Effect of zinc application on yield and cadmium content of spinach (*Spinacea oleracea* L.) grown in a cadmium-polluted soil. *Ann. Biol.* 3:59-60.
- Sajwan, K. S., W. H. Ornes, and T. V. Youngblood. 1996. Beryllium phytotoxicity in soybeans. *Water, Air Soil Pollut.* 86:117-124.
- Sasaki, M., M. Kasai, Y. Yamamoto, and H. Matsumoto. 1994. Comparison of the early response to aluminum stress between tolerant and sensitive wheat cultivars: root growth, aluminum content and efflux of K<sup>+</sup>. *J. Plant Nutr.* 17(7):1275-1288.

- Schlegel, H., D. L. Godbold, and A. Huttermann. 1987. Whole plant aspects of heavy metal induced changes in CO<sub>2</sub> uptake and water relations of spruce (*Picea abies*) seedlings. *Physiol. Plantarum*. 69:265-70.
- Schwarzenbach, R. P., P. M. Gschwend, and D. M. Imboden. 1993. *Environmental Organic Chemistry*. John Wiley and Sons, Inc., New York.
- Shacklette, H. T., and J. G. Boerngen. 1984. *Element Concentrations in Soils and Other Surficial Materials of the Conterminous United States*. USGS Prof. Paper 1270. U.S. Govt Printing Office, Washington.
- Sheppard, M. I., T. T. Vandergraaf, D. H. Thibault, and J. A. K. Reid. 1983. Technetium and uranium: Sorption by and plant uptake from peat and sand. *Health Physics* 44(6):635-643.
- Singh, A., N. K. Goyal, and A. P. Gupta. 1991. Effect of cadmium and farm yard manure on the concentration and uptake of zinc by wheat in texturally different soils. *Crop Res.* 4(2):199-205.
- Singh, M., and N. Singh. 1978. Selenium toxicity in plants and its detoxication by phosphorus. *Soil Sci.* 126:255-262.
- Smith, K. L., G. W. Bryan, and J. L. Harwood. 1985. Changes in endogenous fatty acids and lipid synthesis associated with copper pollution in *Fucus*. *J. Exp. Bot.* 36:663-669.
- Smith, S. P. J. Peterson, and K. H. M. Kwan. 1989. Chromium accumulation, transport and toxicity in plants. *Toxicol. Environ. Chem.* 24:241-251.
- Soltanpour, P. N., and S. M. Workman. 1980. Use of NH<sub>4</sub>HCO<sub>3</sub>-DTPA soil test to assess availability and toxicity of selenium to alfalfa plants. *Commun. Soil Sci. Plant Anal.* 11(12):1147-1156.
- Stiborova, M., R. Hromadkova, and S. Leblova. 1986. Effect of ions of heavy metals on the photosynthetic characteristics of maize (*Zea mays* L.). *Biologia.* 41(12):1221-28.
- Stone, E.L., and G. Baird. 1956. Boron level and boron toxicity in red and white pine. *J. Forestry* 54:11-12.
- Strek, J. H. and J. B. Weber. 1980. Absorption and translocation of polychlorinated biphenyls (PCBs) by weeds. *Proc. South. Weed Sci. Soc.* 33:226-232.
- Strek, H.J., and J.B. Weber. 1982a. Behavior of polychlorinated biphenyls (PCBs) in soils and plants. *Environ. Pollut.* 28A:291-312.
- Strek, J. H. and J. B. Weber. 1982b. Adsorption and reduction in bioactivity of polychlorinated biphenyl (Aroclor 1254) to redroot pigweed by soil organic matter and montmorillonite clay. *Soil Sci. Soc. Am. J.* 46:318-22.
- Strickland, R. C., W. R. Chaney, and R. J. Lamoreaux. 1979. Organic matter influences phytotoxicity of cadmium to soybeans. *Plant Soil* 53(3):393-402.
- Suszcynsky, E. M., and J. R. Shann. 1995. Phytotoxicity and accumulation of mercury in tobacco subjected to different exposure routes. *Environ. Toxicol. Chem.* 14(1):61-67.

- Suter, G. W., II, B. E. Sample, D. S. Jones, T. L. Ashwood, and J. M. Loar. 1995. *Approach and Strategy for Performing Ecological Risk Assessments for the U. S. Department of Energy's Oak Ridge Reservation: 1995 Revision*. ES/ER/TM-33/R2. Oak Ridge National Laboratory, Environmental Sciences Division.
- Swaine, D. J. 1978. Selenium: From magma to man. *Trace Subst. Environ. Hlth.* 12:129-134.
- Tiffin, L. O. 1967. Translation of manganese, iron, cobalt and zinc in tomato. *Plant Physiol.* 42:1427-1432.
- Traynor, M. F. and B. D. Knezek. 1973. Effects of nickel and cadmium contaminated soils on nutrient composition of corn plants. *Proc. Annual Conf. on Trace Substances in the Environment.* 7:82-87.
- Trelease, S. F. and H. M. Trelease. 1938. Selenium as a stimulating and possibly essential element for indicator plants. *Am. J. Bot.* 25:372-79.
- Trelease, S. F., A. A. Di Somma, and A. L. Jacobs. 1960. Seleno-amino acid found in *Astragalus bisulcatus*. *Science* 132:618.
- Turner, M. A. 1973. Effect of cadmium treatment on cadmium and zinc uptake by selected vegetable species. *J. Environ. Qual.* 2(1):118-19.
- Turner, M. A. and R. H. Rust. 1971. Effects of chromium on growth and mineral nutrition of soybeans. *Soil Sci. Soc. Am. Proc.* 35:755-58.
- van den Berg, R., C. A. J. Denneman, and J. M. Roels. 1993. Risk assessment of contaminated soil: Proposal for adjusted, toxicologically based Dutch soil clean-up criteria. In *Contaminated Soil '93*. F. Arendt, G. J. Annokkee, R. Bosman, and W.J. van den Brink (eds). Kluwer Academic Pub., Netherlands. pp. 349-364.
- van Gestel, C. A. M., D. M. M. Adema, and E. M. Dirven-van Breemen. 1996. Phytotoxicity of some chloroanilines and chlorophenols, in relation to bioavailability in soil. *Water, Air Soil Pollut.* 88:119-132.
- van Goor, B. J., and D. Wiersma. 1976. Chemical forms of Mn and Zn in phloem exudates. *Physiol. Plant.* 36:213-216.
- Wallace, A. 1979. Excess trace metal effects on calcium distribution in plants. *Commun. Soil Sci. Plant Anal.* 10:473-79.
- Wallace, A., and R. M. Romney. 1971. Some interactions of Ca, Sr, and Ba in plants. *Agronomy J.* 63:245-248.
- Wallace, A. and E. M. Romney. 1977a. Aluminum toxicity in plants grown in solution culture. *Commun. Soil Sci. Plant Anal.* 8(9):791-94.
- Wallace, A. and E. M. Romney. 1977b. Roots of higher plants as a barrier to translocation of some metals to shoots of plants. In *Biological Implications of Metals in the Environment*. Proceeding of the Fifteenth Annual Hanford Life Sciences Symposium, Richland, WA. Tech Info Center, ERDA, Washington, DC. pp. 370-379.

- Wallace, A., R. T. Mueller, and R. A. Wood. 1980. Selenate toxicity effects on bush beans grown in solution culture. *J. Plant Nutr.* 2(1&2):107-109.
- Wallace, A., G. V. Alexander, and F. M. Chaudhry. 1977a. Phytotoxicity of cobalt, vanadium, titanium, silver, and chromium. *Commun. Soil Sci. Plant Anal.* 8(9):751-56.
- Wallace, A., G. V. Alexander, and F. M. Chaudhry. 1977b. Phytotoxicity and some interactions of the essential trace metals iron, manganese, molybdenum, zinc, copper, and boron. *Commun. Soil Sci. Plant Anal.* 8(9):741-50.
- Wallace, A., R. M. Romney, J. W. Cha, S. M. Soufi, and F. M. Chaudhry. 1977c. Lithium toxicity in plants. *Commun. Soil Sci. Plant Anal.* 8(9):773-80.
- Wallace, A., R. M. Romney, J. W. Cha, S. M. Soufi, and F. M. Chaudhry. 1977d. Nickel phytotoxicity in relationship to soil pH manipulation and chelating agents. *Commun. Soil Sci. Plant Anal.* 8(9):757-64.
- Wallace, A., E. M. Romney, G. V. Alexander, R. T. Mueller, S. M. Soufi, and P. M. Patel. 1977e. Some interactions in plants among cadmium, other heavy metals, and chelating agents. *Agronomy J.* 69:18-20.
- Wallihan, E. F. 1966. Iron. In *Diagnostic Criteria for Plants and Soils*. H. D. Chapman (ed.). Univ. of California, Div. Agric. Sci., Riverside. pp. 203-212.
- Wan H. F., R. L. Mikkelsen, and A. L. Page. 1988. Selenium uptake by some agricultural crops from central California soils. *J. Environ. Qual.* 17(2):269-272.
- Wang, W. 1994. Rice seed toxicity tests for organic and inorganic substances. *Environ. Monit. Assess.* 29:101-107.
- Ward, N. I., E. Roberts, and R. R. Brooks. 1979. Silver uptake by seedlings of *Lolium perenne* L. and *Trifolium repens* L. *N. Z. J. Sci.* 22:129-132.
- Warrington, K. 1954. The influence if iron supply on toxic effects of manganese, molybdenum and vanadium on soybeans, peas and flax. *Ann. Appl. Biol.* 41(1):1-22.912.
- Watkins, D. R., J. T. Ammons, J. L. Branson, B. B. Burgoa, et al. 1993. *Final Report on the Background Soil Characterization Project at the Oak Ridge Reservation, Oak Ridge Tennessee, Vols. 1 and 2*. EE/ER/TM-84. Oak Ridge National Laboratory, Environmental Restoration Division.
- Weber, J. B. and E. Mrozek, Jr. 1979. Polychlorinated biphenyls: Phytotoxicity, absorption and translocation by plants, and inactivation by activated carbon. *Bull. Environ. Contam. Toxicol.* 23:412-17.
- Wheeler, D. M. and J. M. Follet. 1991. Effect of aluminum on onions, asparagus and squash. *J. Plant Nutr.* 14(9):897-912.

- White, M. C., R. L. Chaney, and A. M. Decker. 1979. Differential cultivar tolerance in soybean to phytotoxic levels of soil Zn. II. Range of Zn additions and the uptake and translocation of Zn, Mn, Fe, and P. *Agronomy J.* 71:126-131.
- Whitehead, N. E., R. R. Brooks, and P. J. Peterson. 1971. The nature of uranium occurrence in the leaves of *Coprosma australis* (A. Rich) Robinson. *Aust. J. Biol. Sci.* 24:67-73.
- Wierzbicka, M., and D. Antosiewicz. 1993. How lead can easily enter the food chain - a study of plant roots. *Sci. Total Environ. Suppl.*:423-429.
- Wildung, R. E., T. R. Garland, and D. A. Cataldo. 1977. Accumulation of technetium by plants. *Health Physics* 32:314-317.
- Wildung, R. E., K. M. McFadden, and T. R. Garland. 1979. Technetium sources and behavior in the environment. *J. Environ. Qual.* 8(2):156-161.
- Will, M. E., and G. W. Suter, II. 1995. *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Soil and Litter Invertebrates and Heterotrophic Processes*. ES/ER/TM-126/R1. Oak Ridge National Laboratory, Environmental Sciences Division.
- Will, M. E., and G. W. Suter, II. 1995. *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Terrestrial Plants: 1995 Revision*. ES/ER/TM-85/R2. Oak Ridge National Laboratory, Environmental Sciences Division.
- Wong, M. H. and A. D. Bradshaw. 1982. A comparison of the toxicity of heavy metals, using root elongation of rye grass, *Lolium perenne*. *New Phytol.* 92:255-61.
- Wong, M. H. and W. M. Lau. 1985. Root growth of *Cynodon* and *Eleusine indica* collected from motorways at different concentrations of lead. *Environ. Res.* 36:257-67.
- Woolson, E. A., J. H. Axley, and P. C. Kearney. 1971. Correlation between available soil arsenic, estimated by six methods and response of corn (*Zea mays* L.). *Soil Sci. Soc. Am. Proc.* 35:101-105.
- Wu, L., Z. Huang, and R. G. Bureau. 1988. Selenium accumulation and selenium-salt cotolerance in five grass species. *Crop Sci.* 517-522.
- Zavas, T., L. Symeonidis, and S. Karataglis. 1996. Responses to aluminum toxicity effects of two populations of *Piptatherum miliaceum* (L.) Cosson. *J. Agron. and Crop Sci.* 177:25-32.



**APPENDIX A**

**PHYTOTOXICITY DATA DERIVED FROM EXPERIMENTS  
CONDUCTED IN SOIL**

**Table A.1. Phytotoxicity data derived from experiments conducted in soil**

[All chemical concentrations in soils and plants are mg of the element/kg medium; OM = % organic matter in the soil;

CEC = cation exchange capacity in milliequivalents/100 g soil (dry weight)]

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	silt loam			5	white clover			50	seedling establish	Mackay et al. 1990
Antimony		surface soil							5	phytotoxic	Kabata-Pendias & Pendias 1984
Arsenic	NaAsO <sub>2</sub>	sand		1	6	barley	365		2	grain yield	Jiang & Singh 1994
Arsenic	As <sub>2</sub> O <sub>3</sub>	sandy loam				cotton	42		11.2	shoot weight	Deuel & Swoboda 1972
Arsenic	As <sub>2</sub> O <sub>3</sub>	sandy loam				soybean	42		11.2	shoot weight	Deuel & Swoboda 1972
Arsenic	As <sub>2</sub> O <sub>3</sub>	black clay				soybean	42		22.4	shoot weight	Deuel & Swoboda 1972
Arsenic	NaAsO <sub>2</sub>	loam		6	5	barley	365	10	50	grain yield	Jiang & Singh 1994
Arsenic	NaAsO <sub>2</sub>	sand		1	6	ryegrass	365	10	50	grain yield	Jiang & Singh 1994
Arsenic	NaHAsO <sub>4</sub>	sand		1	6	barley	365	10	50	grain yield	Jiang & Singh 1994
Arsenic	As <sub>2</sub> O <sub>3</sub>	black clay				cotton	42	67.2	89.6	shoot weight	Deuel & Swoboda 1972

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Arsenic	NaH <sub>2</sub> AsO <sub>4</sub>	sandy loam			7	corn	28	10	100	fresh weight	Woolson et al. 1971
Arsenic	Al(H <sub>2</sub> AsO <sub>4</sub> ) <sub>3</sub>	loamy sand			7	corn	28	10	100	fresh weight	Woolson et al. 1971
Arsenic	Ca(H <sub>2</sub> AsO <sub>4</sub> ) <sub>2</sub>	loamy sand			7	corn	28	10	100	fresh weight	Woolson et al. 1971
Arsenic	NaAsO <sub>2</sub>	loam		6	5	ryegrass	365	50	250	grain yield	Jiang & Singh 1994
Arsenic	Na <sub>2</sub> HAsO <sub>4</sub>	loam		6	5	ryegrass	365	50	250	grain yield	Jiang & Singh 1994
Arsenic	Na <sub>2</sub> HAsO <sub>4</sub>	sand		1	6	ryegrass	365	50	250	grain yield	Jiang & Singh 1994
Arsenic	Na <sub>2</sub> HAsO <sub>4</sub>	loam		6	5	barley	365	50	250	grain yield	Jiang & Singh 1994
Arsenic	As <sub>2</sub> O <sub>3</sub>					spruce	335		1000	height	Rosehart & Lee 1973
Barium	Ba(NO <sub>3</sub> ) <sub>2</sub>	loam				barley	14		500	plant weight	Chaudhry et al. 1977
Barium	Ba(NO <sub>3</sub> ) <sub>2</sub>	loam				bush beans	14	1000	2000	plant weight	Chaudhry et al. 1977
Beryllium		surface soil							10	phytotoxic	Kabata-Pendias & Pendias 1984

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Beryllium	BeSO <sub>4</sub>	sand			4.9	soybean	14		25LCT	shoot weight	Sajwan et al. 1996
Beryllium	BeSO <sub>4</sub>	loamy sand			5.5	soybean	14		25LCT	shoot weight	Sajwan et al. 1996
Boron	H <sub>3</sub> BO <sub>3</sub>	silt loam	23	6	6	corn	49		0.5	shoot weight	John et al. 1977
Boron	H <sub>3</sub> BO <sub>3</sub>	muck	117	56	5	corn	49	10	50	shoot weight	John et al. 1977
Boron	H <sub>3</sub> BO <sub>3</sub>	silt loam	16	3	6	corn	49	10	50	shoot weight	John et al. 1977
Bromine		surface soil							10	phytotoxic	Kabata-Pendias & Pendias 1984
Cadmium	CdCl <sub>2</sub>	silt loam	8		6	soybean	28		1	shoot weight	Miller et al. 1976
Cadmium	CdCl <sub>2</sub>	sand + peat			6	soybeans	42		1.25	plant weight	Strickland et al. 1979
Cadmium	CdCl <sub>2</sub>	soil + sand				spruce	98	1	2	root & shoot weights	Burton et al. 1984
Cadmium	CdCl <sub>2</sub>	sand + peat	0.4	0.5	6	soybeans	42	1.25	2.5	plant weight	Strickland et al. 1979
Cadmium	CdCl <sub>2</sub>	silty clay loam	31	4	7	radish	26		2.5	root weight	Haghiri 1973
Cadmium	CdCl <sub>2</sub>	silty clay loam	31	4	7	lettuce	37		2.5	plant weight	Haghiri 1973
Cadmium		loamy sand	2		6	corn	28		2.5	shoot weight	Miller et al. 1977
Cadmium		loamy sand		1	8	spinach		2	4	plant weight	Sadana & Singh 1987b

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Cadmium	CdSO4	silt loam	14		8	spinach			4	leaf weight	Bingham et al. 1975
Cadmium	CdSO4	silt loam	14		8	soybean			5	bean weight	Bingham et al. 1975
Cadmium	CdCl2	silty clay loam				sycamore	90		5	leaf weight	Carlson & Bazzaz 1977
Cadmium	CdCl2	silty clay loam	31	4	7	wheat	35	2.5	5	shoot weight	Haghiri 1973
Cadmium	Cd(NO3)2	sand:peat:soil			5	beech				annual ring width	Hagemeyer et al. 1993
Cadmium	CdSO4	silt loam	14		8	curley cress			8	leaf weight	Bingham et al. 1975
Cadmium	CdCl2	sand	6	2	5	black-eyed susan	42		10	germination; root&shoot weights	Miles & Parker 1979a
Cadmium	CdCl2	sand	6	2	5	blazing star	42		10	root & shoot weights	Miles & Parker 1979a
Cadmium	CdCl2	sand	6	2	5	thimbleweed	42		10	shoot weight	Miles & Parker 1979a
Cadmium	CdCl2	sand	6	2	5	bergamot	42		10	root weight	Miles & Parker 1979a
Cadmium	CdCl2	silt loam	7		5	soybean	28	1	10	shoot weight	Miller et al. 1976
Cadmium	CdCl2	silt loam	9		6	soybean	28	1	10	shoot weight	Miller et al. 1976

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Cadmium	CdCl <sub>2</sub>	silt loam	7		7	soybean	28	1	10	shoot weight	Miller et al. 1976
Cadmium	CdCl <sub>2</sub>	loamy sand	2		6	soybean	28	1	10	shoot weight	Miller et al. 1976
Cadmium	CdCl <sub>2</sub>	Brown earth			5	radish	42		10	root & shoot weights	Khan & Frankland 1983
Cadmium	CdCl <sub>2</sub>	Brown earth			5	oats	42		10	root weight	Khan & Frankland 1984
Cadmium		loamy sand		0.9	8	wheat			10	grain yield	Sadana & Singh 1987a
Cadmium	CdCl <sub>2</sub>	sand	12	2.5	8	bluestem	84		10	root & shoot weights	Miles & Parker 1979b
Cadmium	CdCl <sub>2</sub>	silty clay loam	31	4	7	soybean	35	5	10	shoot weight	Haghiri 1973
Cadmium	CdCl <sub>2</sub>	surface soil				soybean		5	10	seeds/plant	Aery & Sakar 1991
Cadmium	CdCl <sub>2</sub>	sand + peat	1	1	6	soybeans	42	5	10	plant weight	Strickland et al. 1979
Cadmium	CdSO <sub>4</sub>	silt loam	14		8	lettuce			13	head weight	Bingham et al. 1975
Cadmium	C <sub>4</sub> H <sub>6</sub> CdO <sub>4</sub>	acid Cambisol		1.7	6	wheat	28	7	14.1	shoot weight	Reber 1989
Cadmium	CdCl <sub>2</sub>	humic sand		3.7	5	tomato	14	3.2	16	fresh shoot weight	Adema & Henzen 1989
Cadmium	CdSO <sub>4</sub>	silt loam	14		8	corn			18	grain yield	Bingham et al. 1975

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Cadmium	CdSO4	silt loam	14		8	carrot			20	tuber weight	Bingham et al. 1975
Cadmium	CdCl2	sandy loam		1.5	6	red oak	112	10	20	plant weight	Dixon 1988
Cadmium	CdCl2	sandy+clay loam	15	1	8	wheat		10	20	grain & straw yields	Singh et al. 1991
Cadmium	CdCl2	sand + peat	1.5	2	6	soybeans	42	10	20	plant weight	Strickland et al. 1979
Cadmium	CdCl2	loamy sand	2	2.1	7	corn	7	15	25	root length	Hassett et al. 1976
Cadmium	CdCl2	sand	6	2.2	5	corn	35		28	plant weight	Traynor & Knezek 1973
Cadmium	CdSO4	silt loam	14		8	turnip			28	tuber weight	Bingham et al. 1975
Cadmium	CdCl2	sand	6	2	5	Ky bluegrass	42	10	30	root & shoot weights	Miles & Parker 1979a
Cadmium	CdCl2	sand	6	2	5	bluestem	42	10	30	root & shoot weights	Miles & Parker 1979a
Cadmium	CdCl2	sand	6	2	5	poison-ivy	42	10	30	root & shoot weights	Miles & Parker 1979a
Cadmium	CdO	alluvial			6	wheat		10	30	grain yield	Muramoto et al. 1990
Cadmium	CdCl2	loam		1.4	8	lettuce	14	3.2	33	fresh shoot weight	Adema & Henzen 1989

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Cadmium	CdCl <sub>2</sub>	silt loam	38	11.8	5	spinach			40	root & leaf weights	John 1973
Cadmium	CdCl <sub>2</sub>	silt loam	38	11.8	5	peas			40	seed, pod, vine weights	John 1973
Cadmium	CdCl <sub>2</sub>	silt loam	38	11.8	5	oats			40	grain yield	John 1973
Cadmium	CdCl <sub>2</sub>	silt loam	38	11.8	5	radish			40	tuber & top weights	John 1973
Cadmium	CdSO <sub>4</sub>	silt loam	14		8	field bean			40	bean weight	Bingham et al. 1975
Cadmium	CdCl <sub>2</sub>	Brown earth			5	wheat	42		50	root weight	Khan & Frankland 1984
Cadmium	CdSO <sub>4</sub>	silt loam	14		8	wheat			50	grain yield	Bingham et al. 1975
Cadmium	CdCl <sub>2</sub> +CdO	Brown earth			5	radish	42		50	root weight	Khan & Frankland 1984
Cadmium	CdSO <sub>4</sub>	silt loam	14		8	radish			96	tuber weight	Bingham et al. 1975
Cadmium	CdCl <sub>2</sub>	humic sand		3.7	5	oats	14	10	97	fresh shoot weight	Adema & Henzen 1989
Cadmium	CdCl <sub>2</sub>	silt loam	21		6	rye		50	100	shoot weight	Carlson & Rolfe 1979
Cadmium	CdCl <sub>2</sub>	surface soils	38	12.9	6	radish	21		100	top & root weights	John et al. 1972b



Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Cadmium	CdO	alluvial			6	rice		30	100	root & shoot weights	Muramoto et al. 1990
Cadmium	CdO	Brown earth			5	wheat	42		100	root weight	Khan & Frankland 1984
Cadmium	CdO	Brown earth			5	radish	42		100	root & shoot weights	Khan & Frankland 1983
Cadmium	CdCl <sub>2</sub>	silt loam	16		7	soybean	28	10	100	shoot weight	Miller et al. 1976
Cadmium	C <sub>4</sub> H <sub>6</sub> CdO <sub>4</sub>	Phaeosem		2.2	7	wheat	28	56.3	113	shoot weight	Reber 1989
Cadmium	CdCl <sub>2</sub>	humic sand		3.7	5	lettuce	14	32	136	fresh shoot weight	Adema & Henzen 1989
Cadmium	CdCl <sub>2</sub>	loam		1.4	8	oats	14	10	159	leaf weight	Adema & Henzen 1989
Cadmium	CdSO <sub>4</sub>	silt loam	14		8	tomato			160	fruit weight	Bingham et al. 1975
Cadmium	CdSO <sub>4</sub>	silt loam	14		8	zucchini			160	fruit weight	Bingham et al. 1975
Cadmium	CdSO <sub>4</sub>	silt loam	14		8	cabbage			170	head weight	Bingham et al. 1975
Cadmium	CdCl <sub>2</sub>	loam		1.4	8	tomato	14	32	171	fresh shoot weight	Adema & Henzen 1989
Cadmium	CdCl <sub>2</sub>	silt loam	38	11.8	5	lettuce		40	200	root & leaf weights	John 1973

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Cadmium	CdCl <sub>2</sub>	silt loam	38	11.8	5	broccoli		40	200	leaf weight	John 1973
Cadmium	CdCl <sub>2</sub>	silt loam	38	11.8	5	cauliflower		40	200	root & leaf weights	John 1973
Cadmium	CdCl <sub>2</sub>	silt loam	38	11.8	5	carrot		40	200	root, tuber,top weight	John 1973
Cadmium		loam			7	cotton	35		300	leaf & stem weights	Rehab & Wallace 1978
Cadmium		loam			7	cotton	35		300	leaf & stem weights	Rehab & Wallace 1978
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	loam		1.4	8	lettuce	14	0.35	1.8	fresh shoot weight	Adema & Henzen 1989
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	loam		1.4	8	tomato	14	3.2	6.8	fresh shoot weight	Adema & Henzen 1989
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	loam		1.4	8	oats	14	3.5	7.4	fresh shoot weight	Adema & Henzen 1989
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	humic sand		3.7	5	lettuce	14		>11	fresh shoot weight	Adema & Henzen 1989
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	humic sand		3.7	5	tomato	14	10	21	fresh shoot weight	Adema & Henzen 1989
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	loam				soybean	3	10	30	fresh shoot weight	Turner & Rust 1971
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	humic sand		3.7	5	oats	14	11	31	fresh shoot weight	Adema & Henzen 1989

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Cobalt		surface soil							25	phytotoxic	Kabata-Pendias & Pendias 1984
Copper	CuSO4	sand	12	2.5	8	bluestem	84		100	root & shoot weights	Miles & Parker 1979b
Copper	CuSO4	sand	6	1.9	5	bluestem	84		100	root & shoot weights	Miles & Parker 1979b
Copper	CuSO4	loam				bush beans	17	100	200	leaf weight	Wallace et al. 1977b
Fluorine		surface soil							200	phytotoxic	Kabata-Pendias & Pendias 1984
Iodine	KI	loam			7	tomato	97	0.4	4	top weight	Newton & Toth 1952
Iodine	KI	sand			7	tomato	97	0.4	4	top weight	Newton & Toth 1952
Iodine	KI	silt loam			7	tomato	97	0.4	4	top weight	Newton & Toth 1952
Iodine	KI	silt loam			7	tomato	97	0.4	4	top weight	Newton & Toth 1952
Lead	PbCl2	silty clay loam				sycamore	90		50	leaf weight	Carlson & Bazzaz 1977
Lead	PbCl2	sandy loam		1.5	6	red oak	112	20	50	plant weight	Dixon 1988
Lead	PbCl2	soil + sand		45.3	3	spruce	98	50	100	root & shoot weights	Burton et al. 1984

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Lead	PbCl <sub>2</sub>	soil:sand:peat				autumn olive	49	80	160	transpiration	Rolfe & Bazzaz 1975
Lead		loamy sand	2		6	corn	31	125	250	plant weight	Miller et al. 1977
Lead	PbCl <sub>2</sub>	sand	12	2.5	8	bluestem	84		450	root & shoot weights	Miles & Parker 1979b
Lead	PbCl <sub>2</sub>	sand	6	1.93	5	bluestem	84		450	root weight	Miles & Parker 1979b
Lead	PbCl <sub>2</sub>	Brown earth			5	oat	42	100	500	root weight	Khan & Frankland 1984
Lead	PbCl <sub>2</sub>	loamy sand	2	2.1	7	corn	7	250	500	root length	Hassett et al. 1976
Lead	PbCl <sub>2</sub>	Brown earth			5	radish	42	100	500	root weight	Khan & Frankland 1983
Lead	PbCl <sub>2</sub>	silty clay loam	46	17	4	lettuce	30		1000	leaf weight	John & van Laerhoven 1972
Lead	Pb(NO <sub>3</sub> ) <sub>2</sub>	silty clay loam	46	17	4	lettuce	30		1000	leaf weight	John & van Laerhoven 1972
Lead	PbCl <sub>2</sub>	Brown earth			5	wheat	42	500	1000	root weight	Khan & Frankland 1984
Lead	PbO	alluvial			6	wheat		300	1000	root & shoot weights	Muramoto et al. 1990
Lead	PbO	Brown earth			5	radish	42		1000	root weight	Khan & Frankland 1983

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Lead	PbCl <sub>2</sub>	silt loam	21		6	rye		1000	5000	shoot weight	Carlson & Rolfe 1979
Lead	PbCl <sub>2</sub>	silt loam	21		6	fescue		1000	5000	shoot weight	Carlson & Rolfe 1979
Lithium	LiSO <sub>4</sub>	surface soil				orange	180		2	phytotoxic	Aldrich et al. 1971
Lithium	LiCl	loam			6	bush beans	10	10	25	leaf weight	Wallace et al. 1977c
Lithium	LiNO <sub>3</sub>	loam			6	cotton	16	25	50	leaf & stem weights	Wallace et al. 1977c
Lithium	Li <sub>2</sub> C <sub>2</sub> O <sub>4</sub>	loam			6	barley	10		500	shoot weight	Wallace. 1979
Manganese	MnSO <sub>4</sub>	loam				bush beans	17		500	stem weight	Wallace et al. 1977b
Mercury		surface soil							0.3	phytotoxic	Kabata-Pendias & Pendias 1984
Mercury	chloralkali waste	clayey sand		0.7	7.4	barley	7	34.9	64.0	Seedling height	Panda et al. 1992
Molybdenum									2	phytotoxic	Kabata-Pendias & Pendias 1984
Nickel	NiSO <sub>4</sub>	loam			6	barley	14		25	shoot weight	Wallace et al. 1977d
Nickel	NiCl <sub>2</sub>	sandy loam		1.5	6	red oak	112	20	50	plant weight	Dixon 1988

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Nickel	NiCl <sub>2</sub>		6	1.4	6	oats	110	20	50	grain & straw weights	Halstead et al. 1969
Nickel	NiSO <sub>4</sub>	loam			6	bush beans	14	25	100	leaf weight	Wallace et al. 1977d
Nickel	NiSO <sub>4</sub>	loam			6	bush beans	14		100	shoot weight	Wallace et al. 1977d
Nickel		loam			7	cotton	35		100	leaf & stem weights	Rehab & Wallace 1978
Nickel		loam			7	cotton	35		100	leaf & stem weights	Rehab & Wallace 1978
Nickel	NiCl <sub>2</sub>		12	4.1	6	oats	110	50	100	straw weight	Halstead et al. 1969
Nickel	NiSO <sub>4</sub>	loam			5	ryegrass	28	90	180	shoot weight	Khalid & Tinsley 1980
Nickel	NiSO <sub>4</sub>	loam			4	corn	14	100	250	shoot weight	Wallace et al. 1977d
Nickel	NiSO <sub>4</sub>	loam			8	bush beans	14	100	250	shoot weight	Wallace et al. 1977d
Nickel	NiSO <sub>4</sub>	loam			6	corn	14	100	250	shoot weight	Wallace et al. 1977d
Nickel	NiSO <sub>4</sub>	loam			8	corn	14	100	250	shoot weight	Wallace et al. 1977d

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Nickel	NiCl <sub>2</sub>	sand	5.7	2.2	5	corn	35	220	294	plant weight	Traynor & Knezek 1973
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	loamy sand	4	18.5	6	sorghrass	42		1	shoot weight	Carlson et al. 1991
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	sand	3	11	5	sorghrass	42		1	shoot weight	Carlson et al. 1991
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	loamy sand	4	18.5	6	sorghrass	42		1	shoot weight	Carlson et al. 1991
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	sand	3	11	7	sorghrass	42		1	shoot weight	Carlson et al. 1991
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	sandy loam		13	7	alfalfa		0.5	1.5	shoot weight	Wan et al. 1988
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	clay loam		15	6	alfalfa		0.5	1.5	shoot weight	Wan et al. 1988
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	clay loam		13	7	alfalfa		0.5	1.5	shoot weight	Wan et al. 1988
Selenium	Na <sub>2</sub> SeO <sub>3</sub>	sand	3	11	5	sorghrass	42	1	2	shoot weight	Carlson et al. 1991
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	silty clay loam		6.5	7	alfalfa		1	2	shoot weight	Soltanpour & Workman 1980
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	silty clay loam		5	8	alfalfa		1	2	shoot weight	Soltanpour & Workman 1980
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	silty clay loam		3.7	8	alfalfa		1	2	shoot weight	Soltanpour & Workman 1980

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	silty clay loam		3.1	8	alfalfa		1	2	shoot weight	Soltanpour & Workman 1980
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	silty clay loam		6.3	7	alfalfa		2	4	shoot weight	Soltanpour & Workman 1980
Selenium	Na <sub>2</sub> SeO <sub>3</sub>	sand	3.25	0.2	7.9	wheat	50		2.5LCT	weight	Singh and Singh 1978
Silver		surface soil							2		Kabata-Pendias & Pendias 1984
Technetium	TcO <sub>4</sub> <sup>-</sup>	silt loam		1.4	7	wheat	30	0.1	1	shoot weight	Wildung et al. 1977
Technetium	TcO <sub>4</sub> <sup>-</sup>	silt loam		1.4	7	soybean	30	0.1	1	shoot weight	Wildung et al. 1977
Thallium		surface soil							1	phytotoxic	Kabata-Pendias & Pendias 1984
Tin		surface soil							50	phytotoxic	Kabata-Pendias & Pendias 1984
Tin	SnCl <sub>2</sub>	loam			6	bush bean	17	50	500	shoot weight	Romney et al. 1975
Uranium	UO <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub>	sand	1.2		6	swiss chard			5	root weight	Sheppard et al. 1983
Uranium	UO <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub>	peat	65	92	3	swiss chard			10	root weight	Sheppard et al. 1983
Vanadium		surface soil							2.5	phytotoxic	EPA 1980



Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Vanadium		surface soil							50	phytotoxic	Kabata-Pendias & Pendias 1984
Zinc	ZnSO4	surface soil				soybean		10	25	seeds/plant	Aery & Sakar 1991
Zinc	ZnSO4	sandy loam			6	soybean			131	leaf weight	White et al. 1979
Zinc	ZnSO4	sandy loam			7	soybean			393	leaf weight	White et al. 1979
Zinc	ZnO	alluvial soil			6	wheat			1000	plant weight & grain yield	Muramoto et al. 1990
Zinc	ZnO	alluvial soil			6	rice			1000	root weight	Muramoto et al. 1990
Zinc	ZnSO4	surface soil				coriander	60		87	root & shoot weights	Lata and Veer 1990
Zinc	ZnSO4	surface soil				spinach	60		87	root & shoot weights	Lata and Veer 1990
Zinc	ZnSO4	sand:peat:soil			5	beech			3.3	annual ring width	Hagemeyer et al. 1993
Zinc	ZnSO4	clay loam				cowpea	30	158	316	shoot weight	Gall & Barnette 1940
Zinc	ZnSO4	clay loam				corn	30	474	632	shoot weight	Gall & Barnette 1940
Zinc	ZnSO4	sandy loam				cowpea	30	112	222	shoot weight	Gall & Barnette 1940

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Zinc	ZnSO4	sandy loam				corn	30	222	334	shoot weight	Gall & Barnette 1940
Zinc	ZnSO4	sand				corn	30	202	404	shoot weight	Gall & Barnette 1940
Zinc	ZnSO4	sand				cowpea	30	81	141	shoot weight	Gall & Barnette 1940
Acenaphthene		loam		2	8	lettuce	14		25	fresh weight shoot	Hulzebos et al. 1993
Biphenyl		loam		2	8	lettuce	14		68	fresh weight shoot	Hulzebos et al. 1993
3-chloroaniline		loam		2	7.5	lettuce	14	ca. 7	23	shoot weight	Van Gestel et al. 1996
3-chlorophenol		loam		2	7.5	lettuce	14	ca. 2	7	shoot weight	Van Gestel et al. 1996
3,4-dichlorophenol		loam		2	7.5	lettuce	7	ca. 8	25	shoot weight	Van Gestel et al. 1996
2,4 Dinitrophenol		Clay			5	Fescue	21	20	40	Fresh weight shoot	Overcash et al. 1982
2,4 Dinitrophenol		Clay			5	Corn	21	20	40	Fresh weight shoot	Overcash et al. 1982
2,4 Dinitrophenol		Clay			5	Soybeans	21		20	Fresh weight shoot	Overcash et al. 1982

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
2,4 Dinitrophenol		Sandy loam			4	Soybean	21	20	40	% seed germination	Overcash et al. 1982
2,4 Dinitrophenol		Sandy loam			6	Fescue	21	60	80	Fresh weight shoot	Overcash et al. 1982
2,4 Dinitrophenol		Sandy loam			4	Corn	21	60	80	% seed germination	Overcash et al. 1982
2,4 Dinitrophenol		Sandy loam			4	Fescue	21	20	40	Fresh weight shoot	Overcash et al. 1982
2,4 Dinitrophenol		Sandy loam			6	Corn	21	20	40	Fresh weight shoot	Overcash et al. 1982
2,4 Dinitrophenol		Sandy loam			4	Soybeans	21	20	40	Fresh weight shoot	Overcash et al. 1982
2,4 Dinitrophenol		Sandy loam			4	Corn	21		20	Fresh weight shoot	Overcash et al. 1982
2,4 Dinitrophenol		Sandy loam			6	Soybeans	21		20	Fresh weight shoot	Overcash et al. 1982
Di-n-butyl phthalate		Sandy loam			6	Corn	21		200	Fresh weight shoot	Overcash et al. 1982
Di-n-butyl phthalate		Sandy loam			6	Soybeans	21		200	Fresh weight shoot	Overcash et al. 1982
Di-n-butyl phthalate		Sandy loam			6	Fescue	21	200	2000	Fresh weight shoot	Overcash et al. 1982

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Di-n-butyl phthalate		Clay			5	Corn	21		200	Fresh weight shoot	Overcash et al. 1982
Di-n-butyl phthalate		Clay			5	Fescue	21	200	2000	Fresh weight shoot	Overcash et al. 1982
Di-n-butyl phthalate		Sandy loam			4	Soybean	21		200	% seed germination	Overcash et al. 1982
Di-n-butyl phthalate		Sandy loam			4	Corn	21		200	Fresh weights root & shoot	Overcash et al. 1982
Diethylphthalate		loam		2	8	lettuce	14		134	fresh weight shoot	Hulzebos et al. 1993
Furan		loam		2	8	lettuce	14		617	fresh weight shoot	Hulzebos et al. 1993
Pentachlorophenol		loam		2	8	lettuce	14		8	fresh weight shoot	Hulzebos et al. 1993
Pentachlorophenol		loam		2	8	lettuce	14		3.2	fresh weight shoot	Hulzebos et al. 1993
Pentachlorophenol		sandy loam		2.2	6.1	turnip	14		10	fresh weight shoot	Gunther and Pestemer 1990
Pentachlorophenol		sandy loam		2.2	6.1	oat	14		20	fresh weight shoot	Gunther and Pestemer 1990
Phenol		loam		2	8	lettuce	14		79	fresh weight shoot	Hulzebos et al. 1993

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Phenol		loam		2	8	lettuce	14		168	fresh weight shoot	Hulzebos et al. 1993
PCB - Aroclor 1254		Sand	1.5	1	5	Soybean	26	10	100	Fresh weight shoot	Weber & Mrozek 1979
PCB - Aroclor 1254		Sand	1.5	1	5	Soybean	16		1000	Fresh weight shoot	Strek & Weber 1980
PCB - Aroclor 1254		Sand	1.5	1	5	Pigweed		40	100	Fresh weight leaves & plant height	Strek & Weber 1980
PCB - Aroclor 1254		Sand	1.5	1	5	Soybean	16		1000	Fresh weight shoot	Strek & Weber 1980
PCB - Aroclor 1254		Sand	1.5	1	5	Pigweed		20	40	Fresh weight leaves & plant height	Strek & Weber 1980
PCB - Aroclor 1254		Sand	1.5	1	5	Soybean	16		1000	Fresh weight leaves & plant height	Strek & Weber 1980
PCB - Aroclor 1254		Sand	1.5	1	5	Beet			1000	Fresh weight leaves & plant height	Strek & Weber 1980
PCB - Aroclor 1254		Sand	1.5	1.4	4	Pigweed	28	50	100	Plant height	Strek & Weber 1982b
Toluene		Clay			5	Soybean	21	2000	20,000	Fresh weight shoot	Overcash et al. 1982

Table A.1 (continued)

Chemical	Chemical form	Soil type	CEC	% OM	pH	Plant species	DUR (D)	Soil NOEC	Soil LOEC	Growth parameter	Reference
Toluene		Clay			5	Corn	21		200	Fresh weight shoot	Overcash et al. 1982
Toluene		Sandy loam			4	Corn		2000	20,000	% germination	Overcash et al. 1982
Toluene		Sandy loam			6	Corn	21	2000	20,000	Fresh weight shoot	Overcash et al. 1982
Toluene		Sandy loam			6	Fescue	21	2000	20,000	Fresh weight shoot	Overcash et al. 1982
Toluene		Sandy loam			6	Soybean	21		200	Fresh weight shoot	Overcash et al. 1982
Toluene		Sandy loam			4	Soybean		200	2000	% germination	Overcash et al. 1982
2,3,5,6-tetrachloroaniline		loam		2	7.5	lettuce	14	ca. 5	17	shoot weight	Van Gestel et al. 1996
2,4,5-trichloroaniline		loam		2	7.5	lettuce	14	ca. 7	23	shoot weight	Van Gestel et al. 1996
2,4,5-trichlorophenol		loam		2	7.5	lettuce	14	ca. 1	4.3	shoot weight	Van Gestel et al. 1996

**APPENDIX B**

**PHYTOTOXICITY DATA DERIVED FROM EXPERIMENTS  
CONDUCTED IN SOLUTION CULTURE**

**Table B.1. Phytotoxicity data derived from experiments conducted in solution culture**

(All chemical concentrations in solutions and plants are mg of the element/L solution;

EXP (D) - Exposure duration in days; LCT - lowest concentration tested)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Aluminum		4.5	barley	4		0.0027LCT	root length	Nichol and Oliveira 1995
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.7	onion	31		0.05 LCT	root & shoot weight	Wheeler and Follet 1991.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.7	asparagus		0.05	0.13	root & shoot weight	Wheeler and Follet 1991.
Aluminum	AlCl <sub>3</sub>	4.5	wheat	5		0.14 LCT	root elongation	Sasaki et al. 1994
Aluminum	AlCl <sub>3</sub>	4.5	wheat	5	0.14	0.27	root elongation	Sasaki et al. 1994
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.7	squash	26	0.13	0.27	root weight	Wheeler and Follet 1991.
Aluminum	AlCl <sub>3</sub>	4.4	corn	7	0.13	0.27	root elongation and weight	Pintro et al. 1996
Aluminum	AlCl <sub>3</sub>	4.4	corn	7	0.27	0.40	root elongation	Pintro et al. 1996
Aluminum	AlCl <sub>3</sub>	4.3	maize	1		0.54 LCT	root elongation	Llugany et al. 1995
Aluminum	KAl(SO <sub>4</sub> ) <sub>2</sub>	7	ryegrass	14		0.63 LCT	length longest root	Wong and Bradshaw 1982.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	lettuce	42	0.54	1.1	air dry weight shoot	McClean and Gilbert 1927.
Aluminum	AlCl <sub>3</sub>	4.3	maize	1	0.54	1.35	root elongation	Llugany et al. 1995
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	beet	126		1.8 LCT	air dry weight plant	McClean and Gilbert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	beet	77		1.8 LCT	air dry weight plant	McClean and Gilbert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	barley	77		1.8 LCT	air dry weight root/shoot	McClean and Gilbert 1927.



Table B.1 (continued)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	rye	63		1.8 LCT	air dry weight root	Mclean and Gilbert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	beet	126		1.8 LCT	air dry weight plant	Mclean and Gilbert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	lettuce	56	0.9	1.8	air dry weight plant	Mclean and Gilbert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4	citrange	60	0.11	2.7	root length	Lin and Myhre 1991.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>		rice	13	0.27	2.7	root & shoot weight	Wallace and Romney 1977a.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>		soybean	13	0.27	2.7	leaf weight	Wallace and Romney 1977a.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	lettuce	56	1.8	2.7	air dry weight plant	Mclean and Gilbert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.5	rye	70		3.6 LCT	air dry weight root	Mclean and Gilbert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	radish	77	1.8	3.6	air dry weight root/shoot	Mclean and Gibert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	carrot	126		3.6 LCT	air dry weight plant	Mclean and Gilbert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	carrot	126		3.6 LCT	air dry weight plant	Mclean and Gilbert 1927.
Aluminum	AlCl <sub>3</sub>	4	Norway Spruce	8	1.4	5.4	root elongation	Godbold & Kettner 1991
Aluminum	AlCl <sub>3</sub>	4.3	barley	30	4	6	root & shoot weight	Macleod and Jackson 1967.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>2</sub>	4.3	turnip	77	3.6	7.2	air dry weight shoot	Mclean and Gilbert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	oat	63	3.6	7.2	air dry weight root/shoot	Mclean and Gilbert 1927.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	cabbage	98		7.2 LCT	air dry weight plant	Mclean and Gilbert 1927.
Aluminum	AlCl <sub>3</sub>	3.5	Douglas fir	279	4	8	shoot weight	Keltjens 1990.

Table B.1 (continued)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Aluminum	AlCl <sub>3</sub> +Al(NO <sub>3</sub> ) <sub>3</sub>	3.8	spruce	21	5.4	8.1	growth rate root	Goransson & Eldhuset 1991
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4	lemon	60	4.8	8.3	fresh weight; root length	Lin and Myhre 1991.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4	orange	60	4.8	8.3	fresh weight; root length	Lin and Myhre 1991.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4	citrumelo	60	4.8	8.3	fresh weight plant	Lin and Myhre 1991.
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4.3	barley	30	8	10	root & shoot weight	Macleod and Jackson 1967.
Aluminum	AlCl <sub>3</sub>	4.5	perennial grass	16	2.2	12.9	root length	Zavas et al. 1996
Aluminum	AlCl <sub>3</sub>	4.5	perennial grass	16	2.2	12.9	root length	Zavas et al. 1996
Aluminum	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	4	orange	60	8.3	24.4	fresh weight; root length	Lin and Myhre 1991.
Aluminum	AlCl <sub>3</sub>	3.5	Douglas fir	279	16	32	root weight & length	Keltjens 1990.
Aluminum	AlCl <sub>3</sub> +Al(NO <sub>3</sub> ) <sub>3</sub>	3.8	pine	21	162	269.8	growth rate shoot	Goransson & Eldhuset 1991
Arsenic	As <sub>2</sub> O <sub>3</sub>		cluster bean			0.001 LCT	root length	Mhatre & Chaphekar 1982.
Arsenic	As <sub>2</sub> O <sub>3</sub>		radish		.001	0.01	root length	Mhatre & Chaphekar 1982.
Arsenic						0.02 LCT		Bowen 1979.
Arsenic	As <sub>2</sub> O <sub>3</sub>		alfalfa		0.1	1	root & shoot lengths	Mhatre & Chaphekar 1982.
Arsenic	As <sub>2</sub> O <sub>3</sub>		mung bean		0.1	1	root & shoot lengths	Mhatre & Chaphekar 1982.
Arsenic	Na <sub>2</sub> HAsO <sub>4</sub>	7.3	mustard	3		5.5 EC50	root length	Fargasova 1994.
Arsenic	Na <sub>2</sub> HAsO <sub>4</sub>	7.3	mustard	3		30 LC50	seed germination	Fargasova 1994.

Table B.1 (continued)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Beryllium		5.3	bush bean	48		0.5 LCT	plant weight	Romney et al. 1962.
Beryllium	BeSO <sub>4</sub>		lettuce	3		0.5 LCT	radicle length	Carlson et al. 1991.
Beryllium	BeSO <sub>4</sub>		turnip	3		0.5 LCT	radicle length	Carlson et al. 1991.
Beryllium	BeCl <sub>2</sub>	5.3	barley	20		2 LCT	plant weight	Romney & Childress 1965.
Beryllium	BeCl <sub>2</sub>	5.3	pea	24		2 LCT	plant weight	Romney & Childress 1965.
Beryllium	BeCl <sub>2</sub>	5.3	lettuce	28		2 LCT	plant weight	Romney & Childress 1965.
Beryllium	BeSO <sub>4</sub>		cabbage	3	1	2.5	radicle length	Carlson et al. 1991.
Beryllium	BeCl <sub>2</sub>	5.3	alfalfa	54	2	4	plant weight	Romney & Childress 1965.
Beryllium	BeSO <sub>4</sub>		radish	3	2.5	5	radicle length	Carlson et al. 1991.
Beryllium	BeSO <sub>4</sub>		wheat	3	10	20	radicle length	Carlson et al. 1991.
Beryllium	BeSO <sub>4</sub>		millet	3	30	40	radicle length	Carlson et al. 1991.
Bismuth						27	phytotoxic	Bowen 1979.
Boron						1		Bowen 1979.
Boron	H <sub>3</sub> BO <sub>3</sub>		bush beans	16	1.1	5.4	root & leaf weights	Wallace et al. 1977b.
Bromine						15	phytotoxic	Martin 1966a.
Cadmium	Cd-acetate		perennial ryegrass	21		0.001LCT	root dry weight	Al-Attar et al. 1988
Cadmium	CdCl <sub>2</sub>	6.3	carrot	35		0.01 LCT	shoot weight	Turner 1973.
Cadmium	Cd(NO <sub>3</sub> ) <sub>2</sub>	6.2	soybeans	21		0.05 LCT	root & leaf weights	Cunningham 1977.

**Table B.1 (continued)**

<b>Chemical</b>	<b>Form</b>	<b>pH</b>	<b>Plant species</b>	<b>DUR (D)</b>	<b>NOEC</b>	<b>LOEC</b>	<b>Growth parameter</b>	<b>Reference</b>
Cadmium		5.2	soybean	21		0.05 LCT		Cunningham et al. 1975.
Cadmium	CdCl <sub>2</sub>		birch	8	0.022	0.056	shoot growth rate	Gussarsson 1996
Cadmium	CdCl <sub>2</sub>	5.5	corn	58	0.01	0.1	plant weight & grain yield	Iwai et al. 1975.
Cadmium	CdSO <sub>4</sub>		bean	21		0.1 LCT	plant weight	Page et al. 1972.
Cadmium	CdSO <sub>4</sub>		turnip	21		0.1 LCT	plant weight	Page et al. 1972.
Cadmium	CdSO <sub>4</sub>		beet	21		0.1 LCT	plant weight	Page et al. 1972.
Cadmium	CdCl <sub>2</sub>	6.3	tomato	14	0.01	0.1	shoot weight	Turner 1973.
Cadmium	CdSO <sub>4</sub>		giant endive	35		0.1 LCT	root & weights	Garate et al. 1993.
Cadmium	CdCl <sub>2</sub>	5.5	wheat	13		0.1 LCT	shoot weight	Jalil et al. 1994
Cadmium	CdSO <sub>4</sub>	4	Norway spruce	7		0.112 LCT	root length	Lamersdorf et al. 1991.
Cadmium	CdSO <sub>4</sub>		chrysanthemum	21		0.112 LCT	root & stem weights	Patel et al. 1976.
Cadmium	CdSO <sub>4</sub>		corn	10		0.112 LCT	fresh plant weight	Stiborova et al. 1986.
Cadmium	CdSO <sub>4</sub>		corn	21	0.25	0.5	plant weight	Page et al. 1972.
Cadmium	CdSO <sub>4</sub>	4.3	Norway Spruce	7		0.56 LCT	root elongation	Godbold & Huttermann 1985
Cadmium	CdCl <sub>2</sub>		lettuce	14		0.84 EC50	fresh shoot weight	Adema and Henzen 1989.
Cadmium	CdCl <sub>2</sub>	5.5	corn	19	0.1	1	plant weight	Iwai et al. 1975.
Cadmium	CdCl <sub>2</sub>	6.3	swiss chard	35	0.1	1	shoot weight	Turner 1973.
Cadmium	CdSO <sub>4</sub>		tomato	21		1 LCT	plant weight	Page et al. 1972.

Table B.1 (continued)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Cadmium	CdSO <sub>4</sub>		pepper	21		1 LCT	plant weight	Page et al. 1972.
Cadmium	CdSO <sub>4</sub>		barley	21		1 LCT	plant weight	Page et al. 1972.
Cadmium	CdSO <sub>4</sub>		lettuce	21		1 LCT	plant weight	Page et al. 1972.
Cadmium	CdCl <sub>2</sub>	6.3	beetroot	35	0.1	1	shoot weight	Turner 1973.
Cadmium	CdSO <sub>4</sub>		sesame	7		1.1 LCT	root growth	Inouhe et al. 1994
Cadmium	CdSO <sub>4</sub>		pea	7		1.1 LCT	root growth	Inouhe et al. 1994
Cadmium	CdSO <sub>4</sub>		radish	7		1.1 LCT	root growth	Inouhe et al. 1994
Cadmium	CdSO <sub>4</sub>		cucumber	7		1.1 LCT	root growth	Inouhe et al. 1994
Cadmium	CdSO <sub>4</sub>		tomato	7		1.1 LCT	root growth	Inouhe et al. 1994
Cadmium	CdSO <sub>4</sub>		Azuki bean	7		1.1 LCT	root growth	Inouhe et al. 1994
Cadmium	CdSO <sub>4</sub>	5.5	cotton			1.12 LCT	plant weight	Rehab and Wallace 1978.
Cadmium	Cd(NO <sub>3</sub> ) <sub>2</sub>	6.5	maize	17		1.12LCT	stalk weight	Kummerova and Brandejsova 1994
Cadmium	CdSO <sub>4</sub>	7	ryegrass	14		1.25 LCT	longest root & shoot	Wong and Bradshaw 1982.
Cadmium	CdCl <sub>2</sub>		rice	6		1.4 EC50	radicle weight	Wang 1994
Cadmium	CdCl <sub>2</sub>	4	corn	12	0.2	2	plant weight	Iwai et al. 1975.
Cadmium	CdCl <sub>2</sub>	5	corn	12	0.2	2	plant weight	Iwai et al. 1975.
Cadmium	CdCl <sub>2</sub>	6	corn	12	0.2	2	plant weight	Iwai et al. 1975.

**Table B.1 (continued)**

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Cadmium	CdCl <sub>2</sub>	6.5	sugar beet	14	0.56	2.25	shoot and whole plant weight	Greger et al. 1991
Cadmium	CdSO <sub>4</sub>		cabbage	21	1	2.5	plant weight	Page et al. 1972.
Cadmium	CdCl <sub>2</sub>		tomato	14	1.1	3 EC50	fresh shoot weights	Adema and Henzen 1989.
Cadmium	CdSO <sub>4</sub>		lettuce	7	1.1	3.4	root growth	Inouhe et al. 1994
Cadmium	CdSO <sub>4</sub>		barley	7	1.1	3.4	root growth	Inouhe et al. 1994
Cadmium	CdCl <sub>2</sub>		broad bean	3	4	6	root length	Misra et al. 1994
Cadmium	CdCl <sub>2</sub>		oat	14		6 EC50	fresh shoot weight	Adema and Henzen 1989.
Cadmium	CdSO <sub>4</sub>		oats	7	3.4	6.8	rootgrowth	Inouhe et al. 1994
Cadmium	CdSO <sub>4</sub>	5	bean	15	0.11	11	root & leaf weights	Wallace 1979.
Cadmium	Cd(NO <sub>3</sub> ) <sub>2</sub>		corn	18	11.2	28.1	root & shoot lengths	Rascio et al. 1993.
Cadmium	Cd(NO <sub>3</sub> ) <sub>2</sub>		wheat	7		29.8LCT	root and shoot length and mass	Ouzounidou et al. 1997
Cadmium	CdCl <sub>2</sub>		maize	5		45 LCT	seed germination, radicle length, & plant weight	El-Enany 1995
Cadmium	CdCl <sub>2</sub>	6.6	mustard	3		48 EC50	root length	Fargasova 1994.
Cadmium	CdCl <sub>2</sub>	6.6	mustard	3		692 LC50	seed germination	Fargasova 1994.
Chromium	CrSO <sub>4</sub>		chrysanthemum	21		0.052 LCT	stem & leaf weights	Patel et al. 1976.
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>		lettuce	14	.004	0.16 EC50	fresh shoot weight	Adema and Henzen 1989.

**Table B.1 (continued)**

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>		tomato	14	0.11	0.29 EC <sub>50</sub>	fresh shoot weight	Adema and Henzen 1989.
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>		bush beans	11		0.54 LCT	leaf weight	Wallace et al. 1977a.
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>		soybean	5	0.5	1	shoot weight	Turner and Rust 1971.
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>		oat	14	0.12	1.4 EC <sub>50</sub>	fresh shoot weight	Adema and Henzen 1989.
Chromium	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	7	ryegrass	14		2.5 LCT	root length	Wong and Bradshaw 1982.
Chromium	K <sub>2</sub> CrO <sub>4</sub>		rice	6		4.8 EC <sub>50</sub>	radicle Weight	Wang 1994
Chromium	CrCl <sub>3</sub> +K <sub>2</sub> CrO <sub>4</sub>	5	cabbage	55	2	10	plant weight	Hara et al. 1976.
Chromium	(NH <sub>4</sub> ) <sub>2</sub> CrO <sub>4</sub>	7.3	mustard	3		46 EC <sub>50</sub>	root length	Fargasova 1994.
Chromium	Cr <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>		rye grass	2.5	10	50	% seed gemination	Breeze 1973.
Chromium	Cr <sub>2</sub> K <sub>2</sub> O <sub>7</sub>		ryegrass	2.5	10	50	% seed germination	Breeze 1973.
Chromium	CrCl <sub>3</sub>		tomato		50	100	root weight and length	Moral et al. 1995
Chromium	(NH <sub>4</sub> ) <sub>2</sub> CrO <sub>4</sub>	7.3	mustard	3		100 LC <sub>50</sub>	seed germination	Fargasova 1994.
Cobalt	CoSO <sub>4</sub>		bush beans	21		0.06 LCT	leaf weight	Wallace et al. 1977a.
Cobalt	CoSO <sub>4</sub>		chrysanthemum	21		0.06 LCT	root weight	Patel et al. 1976.
Cobalt	CoSO <sub>4</sub>	5	honeysuckle	20	1	5	radicle elongation	Patterson & Olson 1983
Cobalt	CoSO <sub>4</sub>	5	paper birch	20	1	5	radicle elongation	Patterson & Olson 1983
Cobalt	CoCl <sub>2</sub>		broad bean	3	8	10	root elongation	Misra et al. 1994
Cobalt	CoSO <sub>4</sub>	5	black spruce	20	5	10	radicle elongation	Patterson & Olson 1983

Table B.1 (continued)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Cobalt	CoSO <sub>4</sub>	5	jack pine	20	10	20	radicle elongation	Patterson & Olson 1983
Cobalt	CoSO <sub>4</sub>	5	red pine	20	10	20	radicle elongation	Patterson & Olson 1983
Cobalt	CoSO <sub>4</sub>	5	white spruce	20	20	50	radicle elongation	Patterson & Olson 1983
Cobalt	CoSO <sub>4</sub>	5	white pine	20	50	100	radicle elongation	Patterson & Olson 1983
Copper	CuSO <sub>4</sub>		ryegrass	14		0.031 LCT	length longest root	Wong and Bradshaw 1982.
Copper	CuSO <sub>4</sub>		corn	10		0.064 LCT	fresh plant weight	Stiborova et al. 1986.
Copper	CuSO <sub>4</sub>		chrysanthemum	21		0.064	root weight	Patel et al. 1976.
Copper	CuCl <sub>2</sub>		rice	6		0.22 EC50	radicle weight	Wang 1994
Copper	CuSO <sub>4</sub>		maize	14	0.45	0.64	shoot length	Mocquot et al. 1996
Copper	CuSO <sub>4</sub>	5	paper birch	20	0.5	1	radicle elongation	Patterson & Olson 1983
Copper	CuSO <sub>4</sub>	6.1	honeysuckle	110		4 LCT	stem dia increase; plant weight	Heale and Ormrod 1982.
Copper	CuSO <sub>4</sub>	6.1	dogwood	110		4 LCT	stem dia increase; plant weight	Heale and Ormrod 1982.
Copper	CuSO <sub>4</sub>	6.1	red pine	110		4 LCT	plant weight	Heale and Ormrod 1982.
Copper	CuSO <sub>4</sub>	5	black spruce	20	1	5	radicle elongation	Patterson & Olson 1983
Copper	CuSO <sub>4</sub>	5	red pine	20	1	5	radicle elongation	Patterson & Olson 1983
Copper	CuSO <sub>4</sub>	5	jack pine	20	5	10	radicle elongation	Patterson & Olson 1983
Copper	CuSO <sub>4</sub>	6.1	maple	110	2	10	plant weight	Heale and Ormrod 1982.



**Table B.1 (continued)**

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Copper	CuSO <sub>4</sub>	5	white spruce	20	10	20	radicle elongation	Patterson & Olson 1983
Copper	CuSO <sub>4</sub>	5	honeysuckle	20	20	50	radicle elongation	Patterson & Olson 1983
Copper	CuSO <sub>4</sub>		rice	4	6.4	64	root length	Gupta and Mukherji 1977.
Copper	CuSO <sub>4</sub>	5	white pine	20	50	100	radicle elongation	Patterson & Olson 1983
Fluorine						5		Bowen 1979.
Iodine	KI	5.8	corn	60	0.1	0.5	shoot weight	Lewis and Powers 1941.
Iodine	KI		tomato	60	0.5	5	shoot weight	Newton and Toth 1952.
Iron						10 LCT		Wallihan 1966.
Iron	FeSO <sub>4</sub>		bush bean	15	11.2	28	root, leaf & stem weights	Wallace et al. 1977b.
Iron	FeSO <sub>4</sub>	5	cabbage	55	10	50	plant weight	Hara et al. 1976.
Lead	Pb(NO <sub>3</sub> ) <sub>2</sub>		cluster bean			0.001 LCT	root length	Mhatre & Chaphekar 1982.
Lead	PbCl <sub>2</sub>	4	Norway Spruce	8		0.021 LCT	root elongation	Godbold & Kettner 1991
Lead	PbCl <sub>2</sub>	4	Norway spruce	7		0.021 LCT	root length	Lamersdorf et al. 1991.
Lead	Pb(NO <sub>3</sub> ) <sub>2</sub>		alfalfa		0.01	0.1	root length	Mhatre & Chaphekar 1982.
Lead	Pb(NO <sub>3</sub> ) <sub>2</sub>		onion	4	0.02	0.21	root growth	Liu et al. 1994
Lead	Pb(NO <sub>3</sub> ) <sub>2</sub>		mung bean		0.1	1	root length	Mhatre & Chaphekar 1982.
Lead	Pb(NO <sub>3</sub> ) <sub>2</sub>		radish		0.1	1	shoot length	Mhatre & Chaphekar 1982.

Table B.1 (continued)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Lead			barley	7	1	2	root length	Wierzbicka & Antosiewicz 1993
Lead			maize	7		1 LCT	root length	Wierzbicka & Antosiewicz 1993
Lead	Pb(NO3)2		ryegrass	14		2.5 LCT	root & shoot lengths	Wong and Bradshaw 1982.
Lead	Pb(C2H3O2)2		rice	6		9.7 EC50	radicle weight	Wang 1994
Lead	Pb(NO3)2		wire grass	14		10 LCT	root length	Wong and Lau 1985.
Lead	Pb(NO3)2		Bermuda grass	14		10 LCT	root length	Wong and Lau 1985.
Lead	Pb(NO3)2		Bermuda grass	14		10 LCT	root length	Wong and Lau 1985.
Lead	Pb(NO3)2		wire grass	14		10 LCT	root length	Wong and Lau 1985.
Lead	Pb(NO3)2		Bermuda grass	14		10 LCT	root length	Wong and Lau 1985.
Lead	PbSO4		french bean	28	5	10	plant weight	Hooper 1937.
Lead	PbSO4		french bean	28	5	10	plant weight	Hooper 1937.
Lead	PbSO4		french bean	28	5	10	plant weight	Hooper 1937.
Lead	Pb(NO3)2		wire grass	14	10	20	root length	Wong and Lau 1985.
Lead	PbSO4		french bean	28	20	30	plant weight	Hooper 1937.
Lead	PbSO4		french bean	28	20	30	plant weight	Hooper 1937.
Lead	Pb(NO3)2		corn	10	20.7	207	fresh plant weight	Stiborova et al. 1986.
Lead	Pb(CH3COO)2	5.5	mustard	3		263 EC50	root length	Fargasova 1994.

**Table B.1 (continued)**

<b>Chemical</b>	<b>Form</b>	<b>pH</b>	<b>Plant species</b>	<b>DUR (D)</b>	<b>NOEC</b>	<b>LOEC</b>	<b>Growth parameter</b>	<b>Reference</b>
Lead	Pb(CH <sub>3</sub> COO) <sub>2</sub>	5.5	mustard	3		1148 LC50	seed germination	Fargasova 1994.
Lithium	LiNO <sub>3</sub>		bush beans	24		3.5 LCT	stem weight	Wallace et al. 1977c.
Manganese	MnSO <sub>4</sub>	7	ryegrass	14		0.75 LCT	length longest root	Wong and Bradshaw 1982.
Manganese	MnSO <sub>4</sub>	4.6	cotton	18		4 LCT	root & leaf weight	Foy et al. 1995
Manganese	MnSO <sub>4</sub>	5	cotton	18		4 LCT	root & leaf weight	Foy et al. 1995
Manganese	MnSO <sub>4</sub>		bush beans	16		5.5 LCT	root, leaf & stem weights	Wallace et al. 1977b.
Manganese	MnSO <sub>4</sub>	5.5	tomato	17	2.8	5.5	plant weight	Le Bot et al. 1990.
Manganese	MnSO <sub>4</sub>	5	cotton	18	4	8	root & leaf weight	Foy et al. 1995
Manganese	MnSO <sub>4</sub>	4.6	cotton	18	8	16	root & leaf weight	Foy et al. 1995
Manganese	MnSO <sub>4</sub>	4.8	wheat	30		30 LCT	root weight	Burke et al. 1990.
Manganese	MnSO <sub>4</sub>	4.8	wheat	30		30 LCT	root weight	Burke et al. 1990.
Manganese	MnSO <sub>4</sub>	4.8	wheat	30		30 LCT	root & shoot weights	Burke et al. 1990.
Manganese	MnSO <sub>4</sub>	4.8	wheat	30		30 LCT	root weight	Burke et al. 1990.
Manganese	MnSO <sub>4</sub>	6	spruce	32	11	44	root length	Langeheinrich et al. 1992.
Manganese	MnSO <sub>4</sub>	6	spruce	32	11	44	growth rate	Langheinrich et al. 1992.
Manganese			potato	32		33.5 LCT	fresh shoot weight	Marsh and Peterson 1990.
Manganese	MnSO <sub>4</sub>	4	spruce	77		44 LCT	height epicotyl	Langheinrich et al. 1992.
Manganese	MnSO <sub>4</sub>	4	spruce	77		44 LCT	height epicotyl	Langheinrich et al. 1992.

**Table B.1 (continued)**

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Manganese	MnSO <sub>4</sub>		bush beans	21	5.5	55	root, leaf & stem weights	Wallace et al. 1977b.
Manganese	MnSO <sub>4</sub>	4.8	wheat	30	30	90	root & shoot weights	Burke et al. 1990.
Manganese	MnCl <sub>2</sub>		rice	6		100 EC50	radicle weight	Wang 1994
Mercury	HgCl <sub>2</sub>	4	Norway spruce	7		0.002 LCT	root length	Lamersdorf et al. 1991.
Mercury	HgCl <sub>2</sub>		alfalfa		.001	0.01	root & shoot lengths	Mhatre & Chaphekar 1982.
Mercury	HgCl <sub>2</sub>		Pennisetum		.001	0.01	root & shoot lengths	Mhatre & Chaphekar 1982.
Mercury	HgCl <sub>2</sub>		mustard		.001	0.01	root length	Mhatre & Chaphekar 1982.
Mercury	HgCl <sub>2</sub>		cluster bean		.001	0.01	root & shoot lengths	Mhatre & Chaphekar 1982.
Mercury	HgCl <sub>2</sub>		sorghum		.001	0.01	root length	Mhatre & Chaphekar 1982.
Mercury	HgCl <sub>2</sub>		radish		.001	0.01	root & shoot lengths	Mhatre & Chaphekar 1982.
Mercury	HgCl <sub>2</sub>	4.3	Norway spruce	7		0.02 LCT	root elongation	Godbold & Huttermann 1985
Mercury	HgCl <sub>2</sub>	4.3	spruce	35		0.02 LCT	needle chlorophyll	Schlegel et al. 1987.
Mercury	HgCl <sub>2</sub>		mung bean		.01	0.1	root & shoot lengths	Mhatre & Chaphekar 1982.
Mercury	HgCl <sub>2</sub>		pea	5	0.1	1	seed germination, root length	Mhatre & Chaphekar 1982.
Mercury	HgCl <sub>2</sub>	6	tobacco	10	0.1	1	root & shoot weight	Suszcynsky & Shann 1995
Mercury	HgCl <sub>2</sub>	7	ryegrass	14		5 LCT	root & shoot lengths	Wong and Bradshaw 1982.
Mercury	HgCl <sub>2</sub>	7.4	mustard	3		9.3 EC50	root length	Fargasova 1994.

**Table B.1 (continued)**

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Mercury	Hg <sub>2</sub> Cl <sub>2</sub>		barley	7	10	50	root length & plant weight	Mukhiya et al. 1983.
Mercury	HgCl <sub>2</sub>		barley	7	10	50	root&shoot length, plant weight	Mukhiya et al. 1983.
Mercury	HgCl <sub>2</sub>	7.4	mustard	3		129 LC50	seed germination	Fargasova 1994.
Mercury	CH <sub>3</sub> HgCl	4	Norway spruce	7		0.0002 LCT	root length	Lamersdorf et al. 1991.
Mercury	CH <sub>3</sub> HgCl	4.3	spruce	35		0.002 LCT	transpiration rate/CO <sub>2</sub> uptake	Schlegel et al. 1987.
Mercury	CH <sub>3</sub> HgCl	4.3	spruce	35		0.02 LCT	needle chlorophyll	Schlegel et al. 1987.
Mercury	C <sub>8</sub> H <sub>8</sub> HgO <sub>2</sub>		barley	7	1	5	shoot length & plant weight	Mukhiya et al. 1983.
Mercury	C <sub>4</sub> H <sub>6</sub> HgO <sub>4</sub>		barley	7	5	10	root length & plant weight	Mukhiya et al. 1983.
Molybdenum						0.5 LCT	phytotoxic	Johnson 1966.
Molybdenum	H <sub>2</sub> MoO <sub>4</sub>	5	bean	14		9.6 LCT	leaf weight	Wallace 1979.
Molybdenum	H <sub>2</sub> MoO <sub>4</sub>		bush beans	14		9.6 LCT	leaf weight	Wallace et al. 1977b.
Nickel	Ni(NH <sub>4</sub> ) <sub>2</sub> (SO <sub>4</sub> ) <sub>2</sub>	7	ryegrass	14		0.13 LCT	length longest root	Wong and Bradshaw 1982.
Nickel	NiSO <sub>4</sub>		chrysanthemum	21	0.06	0.59	stem & leaf weights	Patel et al. 1976.
Nickel	NiCl <sub>2</sub>		rice	6		0.85 EC50	radicle weight	Wang 1994
Nickel	NiSO <sub>4</sub>	5	paper birch	20	0.5	1	radicle elongation	Patterson & Olson 1983

**Table B.1 (continued)**

<b>Chemical</b>	<b>Form</b>	<b>pH</b>	<b>Plant species</b>	<b>DUR (D)</b>	<b>NOEC</b>	<b>LOEC</b>	<b>Growth parameter</b>	<b>Reference</b>
Nickel	NiSO4		lettuce	3	0.5	1	radicle length	Carlson et al. 1991.
Nickel	NiSO4		turnip	3	0.5	1	radicle length	Carlson et al. 1991.
Nickel	NiSO4	5	bush beans	21		1.17 LCT	root & leaf weights	Wallace 1979.
Nickel	NiSO4		cabbage	3	1	2	radicle length	Carlson et al. 1991.
Nickel	NiSO4	6.1	honeysuckle	110		2 LCT	stem dia increase; plant weight	Heale and Ormrod 1982.
Nickel	NiSO4	6.1	red pine	110		2 LCT	stem dia increase; plant weight	Heale and Ormrod 1982.
Nickel	NiSO4	5	jack pine	20	1	5	radicle elongation	Patterson & Olson 1983
Nickel	NiSO4	5	red pine	20	1	5	radicle elongation	Patterson & Olson 1983
Nickel	NiSO4	5	black spruce	20	1	5	radicle elongation	Patterson & Olson 1983
Nickel	NiSO4	5	honeysuckle	20	1	5	radicle elongation	Patterson & Olson 1983
Nickel	NiSO4	6	cotton		0.59	5.9	plant weight	Rehab and Wallace 1978.
Nickel	NiCl2		broad bean	3	6	8	root length	Misra et al. 1994
Nickel	NiSO4		wheat	3	4	8	radicle length	Carlson et al. 1991.
Nickel	NiSO4		radish	3	4	8	radicle length	Carlson et al. 1991.
Nickel	NiSO4	6.1	dogwood	110	2	10	stem dia increase; plant weight	Heale and Ormrod 1982.
Nickel	NiSO4	6.1	maple	110	2	10	plant weight	Heale and Ormrod 1982.
Nickel	NiSO4		millet	3	8	12	radicle length	Carlson et al. 1991.

Table B.1 (continued)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Nickel	NiSO <sub>4</sub>	5	white spruce	20	10	20	radicle elongation	Patterson & Olson 1983
Nickel	NiSO <sub>4</sub>	5	white pine	20	20	50	radicle elongation	Patterson & Olson 1983
Selenium	Na <sub>2</sub> SeO <sub>4</sub>		bermudagrass	21		0.5LCT	root length, shoot height	Wu et al. 1988
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	4.4	bush bean			0.79 LCT	root weight	Wallace et al. 1980.
Selenium	Na <sub>2</sub> SeO <sub>3</sub>		wheat	42		1 LCT	root&shoot weight, plant height	Martin 1937a.
Selenium	Na <sub>2</sub> SeO <sub>3</sub>		buckwheat	42		1 LCT	root&shoot weight, plant height	Martin 1937a.
Selenium	Na <sub>2</sub> SeO <sub>4</sub>		crested wheatgrass	21	0.5	1	root length, shoot height	Wu et al. 1988
Selenium	Na <sub>2</sub> SeO <sub>4</sub>		seaside bentgrass	21	0.5	1	root length, shoot height	Wu et al. 1988
Selenium	Na <sub>2</sub> SeO <sub>4</sub>		buffalograss	21	1	2	root length, shoot height	Wu et al. 1988
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	6.7	Indian mustard	60		2LCT	root and shoot dry weight	Banuelos et al. 1997
Selenium	Na <sub>2</sub> SeO <sub>4</sub>	6.7	Brassica carinata	60		2LCT	root and shoot dry weight	Banuelos et al. 1997
Selenium	Na <sub>2</sub> SeO <sub>3</sub>		milk-vetch		9	27	plant weight	Trelease & Trelease 1938.
Silver	AgNO <sub>3</sub>	5	bush bean	13		0.17 LCT	leaf weight	Wallace 1979.

Table B.1 (continued)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Silver	AgNO <sub>3</sub>		rice	6		0.55 EC50	radicle weight	Wang 1994
Technetium	TcO <sub>4</sub> <sup>-</sup>	5.5	soybean	20	0.04	0.2	fresh weight seedlings	Berlyn et al. 1980.
Technetium	NH <sub>4</sub> TcO <sub>4</sub>		wheat	10	0.03	0.3	shoot weight	Gast et al. 1978.
Technetium	NH <sub>4</sub> TcO <sub>4</sub>		barley	10	0.03	0.3	shoot weight	Gast et al. 1978.
Technetium	NH <sub>4</sub> TcO <sub>4</sub>		oat	10	0.3	1.2	root & shoot weights	Gast et al. 1978.
Technetium	NH <sub>4</sub> TcO <sub>4</sub>		radish	10	0.3	1.2	shoot weight	Gast et al. 1978.
Technetium	NH <sub>4</sub> TcO <sub>4</sub>		corn	10	3	5.8	shoot weight	Gast et al. 1978.
Technetium	NH <sub>4</sub> TcO <sub>4</sub>		soybean	10	5.8	7.8	shoot weight	Gast et al. 1978.
Technetium	TcO <sub>4</sub> <sup>-</sup>	5.5	soybean	14	5	20	fresh weight seedlings	Berlyn et al. 1980.
Tellurium	K <sub>2</sub> TeO <sub>3</sub>		wheat	42		2 LCT	root & shoot weights	Martin 1937b.
Thallium	TlCl <sub>3</sub>	4	spruce	7		0.02 LCT	root length	Lamersdorf et al. 1991.
Thallium	TlNO <sub>3</sub>		perennial ryegrass	21	0.1	0.5	root and shoot weight and length	Al-Attar et al. 1988
Thallium	Tl <sub>2</sub> SO <sub>4</sub>		lettuce	3		0.5 LCT	radicle length	Carlson et al. 1991.
Thallium	TlCl <sub>2</sub>		sunflower			1 LCT	photosynthesis	Carlson et al. 1975
Thallium	TlCl <sub>2</sub>		corn			1 LCT	photosynthesis	Carlson et al. 1975.
Thallium						1 LCT	phytotoxic	Bowen 1979.
Thallium	Tl <sub>2</sub> SO <sub>4</sub>		turnip	3	0.5	1	radicle length	Carlson et al. 1991.
Thallium	Tl <sub>2</sub> SO <sub>4</sub>		wheat	3	2.5	5	radicle length	Carlson et al. 1991.



**Table B.1 (continued)**

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Thallium	Tl <sub>2</sub> SO <sub>4</sub>		millet	3	2.5	5	radicle length	Carlson et al. 1991.
Thallium	Tl <sub>2</sub> SO <sub>4</sub>		radish	3	2.5	5	radicle length	Carlson et al. 1991.
Thallium	Tl <sub>2</sub> SO <sub>4</sub>		cabbage	3	5	7.5	radicle length	Carlson et al. 1991.
Tin	SnCl <sub>2</sub>		bush bean	26	11.9	118.7	shoot weight	Romney et al. 1975.
Titanium	TiCl <sub>3</sub>		bush beans	21		0.069 LCT	leaf weight	Wallace et al. 1977a.
Titanium	TiCl <sub>3</sub>	5	cabbage	55	0.4	4	plant weight	Hara et al. 1976.
Uranium	UO <sub>2</sub>		soybean	6	0.42	42	seedling length	Murthy et al. 1984.
Vandium	NH <sub>4</sub> VO <sub>3</sub>	4.7	lettuce	45	0.1	0.2	plant weight	Gil et al. 1995
Vanadium	NH <sub>4</sub> VO <sub>3</sub>	5	bush beans	14	.051	0.51	root weight	Wallace 1979.
Vanadium	VOSO <sub>4</sub>		lettuce	3	1	2.5	radicle length	Carlson et al. 1991.
Vanadium	VOSO <sub>4</sub>		turnip	3	1	2.5	radicle length	Carlson et al. 1991.
Vanadium	VOSO <sub>4</sub>		cabbage	3	1	2.5	radicle length	Carlson et al. 1991.
Vanadium	VCl <sub>3</sub>	5	cabbage	55	0.4	4	plant weight	Hara et al. 1976.
Vanadium	VOSO <sub>4</sub>		soybean	33	3	6	plant weight	Kaplan et al. 1990.
Vanadium	VOSO <sub>4</sub>		radish	3	7.5	10	radicle length	Carlson et al. 1991.
Vanadium	NH <sub>4</sub> VO <sub>3</sub>		pea	14	10	20	root & shoot weights	Nowakowski 1992.
Vanadium	VOSO <sub>4</sub>		millet	3	50	60	radicle length	Carlson et al. 1991.
Zinc		6	clover	46	.082	0.41	plant weight	Carroll & Loneragan 1968.

Table B.1 (continued)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Zinc		6	barrel medic	46	.082	0.41	plant weight	Carroll & Loneragan 1968.
Zinc		6	lucerne	46	.082	0.41	plant weight	Carroll & Loneragan 1968.
Zinc	ZnSO4	7	ryegrass	14		1.85 LCT	root length	Wong and Bradshaw 1982.
Zinc	ZnSO4		chrysanthemum	21	0.65	6.5	stem weight	Patel et al. 1976.
Zinc	ZnSO4		bush beans	16	0.65	6.5	root & shoot weights	Wallace et al. 1977b.
Zinc	ZnCl2		broad bean	3	8	10	root length	Misra et al. 1994
Zinc	ZnCl2		rice	6		26 EC50	radicle weight	Wang 1994
Aniline		7	chinese cabbage	21		203.5	root length	Feng et al. 1996
Biphenyl			lettuce	16		2.1 EC50	fresh weight shoot	Hulzebos et al. 1993
4-Bromoaniline		7	chinese cabbage	21		102.2	root length	Feng et al. 1996
4-Chloroaniline		7	chinese cabbage	21		39.4	root length	Feng et al. 1996
2-Chlorophenol		7	chinese cabbage	21		58.3	root length	Feng et al. 1996
4-Chlorophenol		7	chinese cabbage	21		47.4	root length	Feng et al. 1996
2-Cresol		7	chinese cabbage	21		54.9	root length	Feng et al. 1996
3,4-Dichloroaniline		7	chinese cabbage	21		14.1	root length	Feng et al. 1996
2,4-Dichlorophenol		7	chinese cabbage	21		17.1	root length	Feng et al. 1996
Furan			lettuce	16		130 EC50	fresh weight shoot	Hulzebos et al. 1993
Heptane			lettuce	16		1.7 EC50	fresh weight shoot	Hulzebos et al. 1993

Table B.1 (continued)

Chemical	Form	pH	Plant species	DUR (D)	NOEC	LOEC	Growth parameter	Reference
Heptane			lettuce	16		47 EC50	fresh weight shoot	Hulzebos et al. 1993
Naphthalene			lettuce	16		13 EC50	fresh weight shoot	Hulzebos et al. 1993
3-Nitroaniline		7	chinese cabbage	21		69.2	root length	Feng et al. 1996
4-Nitroaniline		7	chinese cabbage	21		43.6	root length	Feng et al. 1996
Nitrobenzene			autumn olive	2		8 LCT	photosynthesis, transpiration	McFarlane et al. 1990.
4-Nitrophenol		7	chinese cabbage	21		12.3	root length	Feng et al. 1996
Pentachlorophenol			lettuce	16		0.03 EC50	fresh weight shoot	Hulzebos et al. 1993
Pentachlorophenol			lettuce	16		0.03 EC50	fresh weight shoot	Hulzebos et al. 1993
Phenol			lettuce	16		20 EC50	fresh weight shoot	Hulzebos et al. 1993
Phenol			lettuce	16		14 EC50	fresh weight shoot	Hulzebos et al. 1993
Phenol		7	chinese cabbage	21		125.6	root length	Feng et al. 1996
Styrene			lettuce	16		18 EC50	fresh weight shoot	Hulzebos et al. 1993
Toluene			lettuce	16		16 EC50	fresh weight shoot	Hulzebos et al. 1993
4-Toluidine		7	chinese cabbage	21		102.2	root length	Feng et al. 1996
2,4,6-Trichlorophenol		7	chinese cabbage	21		12.7	root length	Feng et al. 1996
Ortho-xylene			lettuce	16		2 EC50	fresh weight shoot	Hulzebos et al. 1993
Xylene		6	sugar beet	2		100 LCT	root length	Allen et al. 1961.